



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : G01N 33/68, 33/53, C07K 16/00, C12N 15/11, 15/12, 15/00, 15/63, A61K 38/17, 38/16, C12P 21/02		A1	(11) International Publication Number: WO 99/24836
			(43) International Publication Date: 20 May 1999 (20.05.99)
(21) International Application Number: PCT/US98/23435		(72) Inventors; and	
(22) International Filing Date: 4 November 1998 (04.11.98)		(75) Inventors/Applicants (for US only): FENG, Ping [CN/US]; 4 Relda Court, Gaithersburg, MD 20878 (US). CARTER, Kenneth, C. [US/US]; 11601 Brandy Hall Lane, North Potomac, MD 20878 (US). ENDRESS, Gregory, A. [US/US]; 9729 Clagett Farm Drive, Potomac, MD 20854 (US). ROSEN, Craig, A. [US/US]; 22400 Rolling Hill Road, Laytonsville, MD 20882 (US). RUBEN, Steven, M. [US/US]; 18528 Heritage Hills Drive, Olney, MD 20832 (US). JANAT, Fouad [US/US]; 140 High Street #202, Westerly, RI 02891 (US). NI, Jian [CN/US]; 5502 Manorfield Road, Rockville, MD 20853 (US). WEI, Ying-Fei [CN/US]; 1714-C Marina Court, San Mateo, CA 94403 (US). MOORE, Paul, A. [GB/US]; 19005 Leatherbark Drive, Germantown, MD 20874 (US). SOPPET, Daniel, R. [US/US]; 15050 Stillfield Place, Centreville, VA 22020 (US). KYAW, Hla [MM/US]; 520 Sugarbush Circle, Frederick, MD 21703 (US). LaFLEUR, David, W. [CN/US]; 3142 Quesada Street, N.W., Washington, DC 20015 (US). OLSEN, Henrik, S. [DK/US]; 182 Kendrick Place #24, Gaithersburg, MD 20878 (US). SHI, Yanggu [CN/US]; 437 West Side Drive #102, Gaithersburg, MD 20878 (US). EBNER, Reinhard [DE/US]; 9906 Shelburne Terrace #316, Gaithersburg, MD 20878 (US).	
(30) Priority Data:			
60/064,911	7 November 1997 (07.11.97)	US	
60/064,912	7 November 1997 (07.11.97)	US	
60/064,983	7 November 1997 (07.11.97)	US	
60/064,900	7 November 1997 (07.11.97)	US	
60/064,988	7 November 1997 (07.11.97)	US	
60/064,987	7 November 1997 (07.11.97)	US	
60/064,908	7 November 1997 (07.11.97)	US	
60/064,984	7 November 1997 (07.11.97)	US	
60/064,985	7 November 1997 (07.11.97)	US	
06/066,094	17 November 1997 (17.11.97)	US	
06/066,100	17 November 1997 (17.11.97)	US	
06/066,089	17 November 1997 (17.11.97)	US	
06/066,095	17 November 1997 (17.11.97)	US	
06/066,090	17 November 1997 (17.11.97)	US	
(71) Applicant (for all designated States except US): HUMAN GENOME SCIENCES, INC. [US/US]; 9410 Key West Avenue, Rockville, MD 20850 (US).		(74) Agents: HOOVER, Kenley, K. et al.; Human Genome Sciences, Inc., 9410 Key West Avenue, Rockville, MD 20850 (US).	
		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
		Published With international search report. With an indication in relation to deposited biological material furnished under Rule 13bis separately from the description.	
(54) Title: 125 HUMAN SECRETED PROTEINS			
(57) Abstract			
<p>The present invention relates to novel human secreted proteins and isolated nucleic acids containing the coding regions of the genes encoding such proteins. Also provided are vectors, host cells, antibodies, and recombinant methods for producing human secreted proteins. The invention further relates to diagnostic and therapeutic methods useful for diagnosing and treating disorders related to these novel human secreted proteins.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

125 Human Secreted Proteins

Field of the Invention

This invention relates to newly identified polynucleotides and the polypeptides encoded by these polynucleotides, uses of such polynucleotides and polypeptides, and
5 their production.

Background of the Invention

Unlike bacterium, which exist as a single compartment surrounded by a membrane, human cells and other eucaryotes are subdivided by membranes into many functionally distinct compartments. Each membrane-bounded compartment, or
10 organelle, contains different proteins essential for the function of the organelle. The cell uses "sorting signals," which are amino acid motifs located within the protein, to target proteins to particular cellular organelles.

One type of sorting signal, called a signal sequence, a signal peptide, or a leader sequence, directs a class of proteins to an organelle called the endoplasmic reticulum
15 (ER). The ER separates the membrane-bounded proteins from all other types of proteins. Once localized to the ER, both groups of proteins can be further directed to another organelle called the Golgi apparatus. Here, the Golgi distributes the proteins to vesicles, including secretory vesicles, the cell membrane, lysosomes, and the other organelles.

20 Proteins targeted to the ER by a signal sequence can be released into the extracellular space as a secreted protein. For example, vesicles containing secreted proteins can fuse with the cell membrane and release their contents into the extracellular space - a process called exocytosis. Exocytosis can occur constitutively or after receipt of a triggering signal. In the latter case, the proteins are stored in secretory vesicles (or
25 secretory granules) until exocytosis is triggered. Similarly, proteins residing on the cell membrane can also be secreted into the extracellular space by proteolytic cleavage of a "linker" holding the protein to the membrane.

Despite the great progress made in recent years, only a small number of genes encoding human secreted proteins have been identified. These secreted proteins include
30 the commercially valuable human insulin, interferon, Factor VIII, human growth hormone, tissue plasminogen activator, and erythropoietin. Thus, in light of the pervasive role of secreted proteins in human physiology, a need exists for identifying and characterizing novel human secreted proteins and the genes that encode them. This knowledge will allow one to detect, to treat, and to prevent medical disorders by using
35 secreted proteins or the genes that encode them.

Summary of the Invention

The present invention relates to novel polynucleotides and the encoded polypeptides. Moreover, the present invention relates to vectors, host cells, antibodies, and recombinant methods for producing the polypeptides and polynucleotides. Also provided are diagnostic methods for detecting disorders related to the polypeptides, and therapeutic methods for treating such disorders. The invention further relates to screening methods for identifying binding partners of the polypeptides.

Detailed Description

Definitions

The following definitions are provided to facilitate understanding of certain terms used throughout this specification.

In the present invention, "isolated" refers to material removed from its original environment (e.g., the natural environment if it is naturally occurring), and thus is altered "by the hand of man" from its natural state. For example, an isolated polynucleotide could be part of a vector or a composition of matter, or could be contained within a cell, and still be "isolated" because that vector, composition of matter, or particular cell is not the original environment of the polynucleotide.

In the present invention, a "secreted" protein refers to those proteins capable of being directed to the ER, secretory vesicles, or the extracellular space as a result of a signal sequence, as well as those proteins released into the extracellular space without necessarily containing a signal sequence. If the secreted protein is released into the extracellular space, the secreted protein can undergo extracellular processing to produce a "mature" protein. Release into the extracellular space can occur by many mechanisms, including exocytosis and proteolytic cleavage.

As used herein, a "polynucleotide" refers to a molecule having a nucleic acid sequence contained in SEQ ID NO:X or the cDNA contained within the clone deposited with the ATCC. For example, the polynucleotide can contain the nucleotide sequence of the full length cDNA sequence, including the 5' and 3' untranslated sequences, the coding region, with or without the signal sequence, the secreted protein coding region, as well as fragments, epitopes, domains, and variants of the nucleic acid sequence. Moreover, as used herein, a "polypeptide" refers to a molecule having the translated amino acid sequence generated from the polynucleotide as broadly defined.

In the present invention, the full length sequence identified as SEQ ID NO:X was often generated by overlapping sequences contained in multiple clones (contig

analysis). A representative clone containing all or most of the sequence for SEQ ID NO:X was deposited with the American Type Culture Collection ("ATCC"). As shown in Table 1, each clone is identified by a cDNA Clone ID (Identifier) and the ATCC Deposit Number. The ATCC is located at 10801 University Boulevard,
5 Manassas, Virginia 20110-2209, USA. The ATCC deposit was made pursuant to the terms of the Budapest Treaty on the international recognition of the deposit of microorganisms for purposes of patent procedure.

A "polynucleotide" of the present invention also includes those polynucleotides capable of hybridizing, under stringent hybridization conditions, to sequences contained
10 in SEQ ID NO:X, the complement thereof, or the cDNA within the clone deposited with the ATCC. "Stringent hybridization conditions" refers to an overnight incubation at 42°C in a solution comprising 50% formamide, 5x SSC (750 mM NaCl, 75 mM sodium citrate), 50 mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 µg/ml denatured, sheared salmon sperm DNA, followed by washing the
15 filters in 0.1x SSC at about 65°C.

Also contemplated are nucleic acid molecules that hybridize to the polynucleotides of the present invention at lower stringency hybridization conditions. Changes in the stringency of hybridization and signal detection are primarily accomplished through the manipulation of formamide concentration (lower percentages
20 of formamide result in lowered stringency); salt conditions, or temperature. For example, lower stringency conditions include an overnight incubation at 37°C in a solution comprising 6X SSPE (20X SSPE = 3M NaCl; 0.2M NaH₂PO₄; 0.02M EDTA, pH 7.4), 0.5% SDS, 30% formamide, 100 µg/ml salmon sperm blocking DNA; followed by washes at 50°C with 1XSSPE, 0.1% SDS. In addition, to achieve even
25 lower stringency, washes performed following stringent hybridization can be done at higher salt concentrations (e.g. 5X SSC).

Note that variations in the above conditions may be accomplished through the inclusion and/or substitution of alternate blocking reagents used to suppress background in hybridization experiments. Typical blocking reagents include
30 Denhardt's reagent, BLOTTO, heparin, denatured salmon sperm DNA, and commercially available proprietary formulations. The inclusion of specific blocking reagents may require modification of the hybridization conditions described above, due to problems with compatibility.

Of course, a polynucleotide which hybridizes only to polyA+ sequences (such
35 as any 3' terminal polyA+ tract of a cDNA shown in the sequence listing), or to a

complementary stretch of T (or U) residues, would not be included in the definition of "polynucleotide," since such a polynucleotide would hybridize to any nucleic acid molecule containing a poly (A) stretch or the complement thereof (e.g., practically any double-stranded cDNA clone).

5 The polynucleotide of the present invention can be composed of any polyribonucleotide or polydeoxribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. For example, polynucleotides can be composed of single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be
10 single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, the polynucleotide can be composed of triple-stranded regions comprising RNA or DNA or both RNA and DNA. A polynucleotide may also contain one or more modified bases or DNA or RNA backbones modified for stability
15 or for other reasons. "Modified" bases include, for example, tritylated bases and unusual bases such as inosine. A variety of modifications can be made to DNA and RNA; thus, "polynucleotide" embraces chemically, enzymatically, or metabolically modified forms.

 The polypeptide of the present invention can be composed of amino acids joined
20 to each other by peptide bonds or modified peptide bonds, i.e., peptide isosteres, and may contain amino acids other than the 20 gene-encoded amino acids. The polypeptides may be modified by either natural processes, such as posttranslational processing, or by chemical modification techniques which are well known in the art. Such modifications are well described in basic texts and in more detailed monographs,
25 as well as in a voluminous research literature. Modifications can occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini. It will be appreciated that the same type of modification may be present in the same or varying degrees at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. Polypeptides may be
30 branched, for example, as a result of ubiquitination, and they may be cyclic, with or without branching. Cyclic, branched, and branched cyclic polypeptides may result from posttranslation natural processes or may be made by synthetic methods. Modifications include acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a
35 nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cysteine,

formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, pegylation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins
5 such as arginylation, and ubiquitination. (See, for instance, PROTEINS - STRUCTURE AND MOLECULAR PROPERTIES, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York (1993); POSTTRANSLATIONAL COVALENT MODIFICATION OF PROTEINS, B. C. Johnson, Ed., Academic Press, New York, pgs. 1-12 (1983); Seifter et al., Meth Enzymol 182:626-646 (1990);
10 Rattan et al., Ann NY Acad Sci 663:48-62 (1992).)

"SEQ ID NO:X" refers to a polynucleotide sequence while "SEQ ID NO:Y" refers to a polypeptide sequence, both sequences identified by an integer specified in Table 1.

"A polypeptide having biological activity" refers to polypeptides exhibiting
15 activity similar, but not necessarily identical to, an activity of a polypeptide of the present invention, including mature forms, as measured in a particular biological assay, with or without dose dependency. In the case where dose dependency does exist, it need not be identical to that of the polypeptide, but rather substantially similar to the dose-dependence in a given activity as compared to the polypeptide of the present
20 invention (i.e., the candidate polypeptide will exhibit greater activity or not more than about 25-fold less and, preferably, not more than about tenfold less activity, and most preferably, not more than about three-fold less activity relative to the polypeptide of the present invention.)

25 Polynucleotides and Polypeptides of the Invention

FEATURES OF PROTEIN ENCODED BY GENE NO: 1

The translation product of this gene shares sequence homology with
30 transcytosis-associated protein (TAP), which is thought to be important in the docking of transport vesicles with their target membrane. The gene encoding the disclosed cDNA is thought to reside on chromosome 4. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 4.

This gene is expressed primarily in developing brain, other embryonic tissue
35 and placenta.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a

biological sample and for diagnosis of diseases and conditions which include, but are not limited to, developmental and neurodegenerative diseases of the brain as well as other developmental anomalies or fetal deficiencies. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the brain, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. embryonic, cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:136 as residues: Pro-51 to Arg-56, Lys-89 to Gln-94, Glu-144 to Gln-151, Gln-178 to Gln-183, Leu-224 to Gln-229, Tyr-284 to Pro-298, Lys-324 to Lys-334.

The tissue distribution in developing brain and placental tissues and the homology to transcytosis-associated protein (TAP) indicates that polynucleotides and polypeptides corresponding to this gene are useful for a host of conditions which arise as a result of a failure of, or deficiency in, the secretory or endocytic pathway. In addition, the expression in brain would suggest a role in the detection and treatment of brain tumors, developmental and behavioral disorders such as mania, depression, paranoia, addictive behavior and sleep disorders. Furthermore, expression within embryonic tissue and other cellular sources marked by proliferating cells indicates that this protein may play a role in the regulation of cellular division. Similarly, embryonic development also involves decisions involving cell differentiation and/or apoptosis in pattern formation. Thus this protein may also be involved in apoptosis or tissue differentiation and could again be useful in cancer therapy. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:11 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 996 of SEQ ID NO:11, b is an integer of 15 to

1010, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:11, and where b is greater than or equal to a + 14.

5 FEATURES OF PROTEIN ENCODED BY GENE NO: 2

This gene is expressed primarily in human adrenal gland tumor.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, endocrine disorders, particularly adrenal gland tumors. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the adrenal gland, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. endocrine, adrenal gland, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis or treatment of adrenal gland tumors. Furthermore, polynucleotides and polypeptides corresponding to this gene are useful for the detection, treatment, and/or prevention of various endocrine disorders and cancers, particularly Addisoni's disease, Cushingi's Syndrome, and disorders and/or cancers of the pancreas (e.g. diabetes mellitus), adrenal cortex, ovaries, pituitary (e.g., hyper-, hypopituitarism), thyroid (e.g. hyper-, hypothyroidism), parathyroid (e.g. hyper-,hypoparathyroidism) , hypothallamus, and testes. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:12 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of

a-b, where a is any integer between 1 to 1545 of SEQ ID NO:12, b is an integer of 15 to 1559, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:12, and where b is greater than or equal to a + 14.

5

FEATURES OF PROTEIN ENCODED BY GENE NO: 3

When tested against U937 Myeloid cell lines, supernatants removed from cells containing this gene activated the GAS assay. Thus, it is likely that this gene activates myeloid cells, including their progenitors, through the Jak-STAT signal transduction pathway. The gamma activating sequence (GAS) is a promoter element found upstream of many genes which are involved in the Jak-STAT pathway. The Jak-STAT pathway is a large, signal transduction pathway involved in the differentiation and proliferation of cells. Therefore, activation of the Jak-STAT pathway, reflected by the binding of the GAS element, can be used to indicate proteins involved in the proliferation and differentiation of cells.

This gene is expressed primarily in small intestine.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, a variety of gastrointestinal disorders including duodenal ulcers. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the gastrointestinal system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. gastrointestinal, smooth muscle, endothelial, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:138 as residues: Gln-77 to Pro-86.

The tissue distribution in small intestine indicates that the translation product of this gene is useful for the diagnosis and/or treatment of a number of disorders having to do with the gastrointestinal system, and specifically the small intestine, such as obstructions of the ileum, meckel's diverticulum, Crohn's disease, celiac sprue, tropical

sprue, and lymphoma. Protein, as well as, antibodies directed against the protein may show utility as a tissue-specific marker and/or immunotherapy target for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:13 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1575 of SEQ ID NO:13, b is an integer of 15 to 1589, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:13, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 4

The translation product of this gene shares sequence homology with the mouse astrotactin protein, which is thought to be important in supporting neuronal migration along glial fibers. Additionally, astrotactin is thought to act as a ligand for neuron-glial binding during neuronal migration. The gene encoding the disclosed cDNA is thought to reside on chromosome 1. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 1.

This gene is expressed primarily in brain tissue from a patient with Alzheimer's disease.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, neural or CNS disorders, particularly neurodegenerative disorders such as Alzheimer's disease. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the central nervous system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. brain, cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the

expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:139 as residues: Gln-43 to Trp-53, Arg-69 to Ser-76.

5 The tissue distribution in brain combined with the homology to mouse astrotactin indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and/or treatment of CNS diseases, such as Alzheimer's disease. Moreover, polynucleotides and polypeptides corresponding to this gene are useful for the detection/treatment of neurodegenerative disease states, behavioural
10 disorders, or inflammatory conditions such as Parkinsons Disease, Huntingtons Disease, Tourette Syndrome, meningitis, encephalitis, demyelinating diseases, peripheral neuropathies, neoplasia, trauma, congenital malformations, spinal cord injuries, ischemia and infarction, aneurysms, hemorrhages, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, panic disorder, learning disabilities, ALS,
15 psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and preception. In addition, elevated expression of this gene product in regions of the brain indicates that it plays a role in normal neural function. Potentially, this gene product is involved in synapse formation, neurotransmission, learning, cognition, homeostasis, or neuronal differentiation or survival. Moreover, the
20 gene or gene product may also play a role in the treatment and/or detection of developmental disorders associated with the developing embryo, sexually-linked disorders, or disorders of the cardiovascular system. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

25 Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:14 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome.

30 Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1241 of SEQ ID NO:14, b is an integer of 15 to 1255, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:14, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 5

The translation product of this gene shares sequence homology with transporter protein, which is thought to be important in metabolic and respiratory functions.

5 This gene is expressed primarily in T-cell lymphoma and dendritic cells, and to a lesser extent in placenta.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, haemopoietic disorders, particularly cancer including T-cell lymphoma and disorders associated with embryogenesis. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. immune, cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:140 as residues: Thr-87 to Trp-94.

The tissue distribution in T-cell lymphoma and dendritic cells and the homology to transporter protein indicates that polynucleotides and polypeptides corresponding to this gene are useful for treatment and diagnosis of haemopoietic disorders such as cancer, particularly T-cell lymphoma and disorders associated with embryogenesis. Furthermore, this gene product may play a role in the survival, proliferation, and/or differentiation of hematopoietic lineages. Expression of this gene product in T cells and primary dendritic cells also strongly indicates a role for this protein in immune function and immune surveillance.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:15 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of

a-b, where a is any integer between 1 to 1177 of SEQ ID NO:15, b is an integer of 15 to 1191, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:15, and where b is greater than or equal to a + 14.

5

FEATURES OF PROTEIN ENCODED BY GENE NO: 6

This gene is expressed primarily in the liver, and to a lesser extent, in testis.

Therefore, polynucleotides and polypeptides of the invention are useful as
10 reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, hepatic, reproductive, or endocrine disorders, particularly hepatoma or male infertility. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s)
15 or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune and hematopoietic systems, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. hepatic, reproductive, endocrine, testical, immune, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, seminal fluid, plasma, urine, synovial fluid and
20 spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:141 as residues: Ser-21 to Trp-34, Cys-68 to Gly-89, Cys-122 to Phe-133.

25 The tissue distribution in liver tissue indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of liver disorders, particularly those affecting the immune and hematopoietic systems such as hepatomas. Furthermore, the protein product of this gene would also be useful for the detection and treatment hepatoblastoma, jaundice, hepatitis, or liver metabolic
30 diseases and conditions that are attributable to the differentiation of hepatocyte progenitor cells. Furthermore, the expression within testis indicates that the protein may show utility in the treatment and/or detection of a variety of reproductive disorders such as male infertility, impotence, and may even be useful as a contraceptive. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker
35 and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ

ID NO:16 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome.

- Accordingly, preferably excluded from the present invention are one or more
 5 polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1172 of SEQ ID NO:16, b is an integer of 15 to 1186, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:16, and where b is greater than or equal to a + 14.

10

FEATURES OF PROTEIN ENCODED BY GENE NO: 7

- The translation product of this gene shares sequence homology with urokinase receptor which is thought to be important in cell matrix remodeling and cell movement.
 15 In specific embodiments, polypeptides of the invention comprise the following amino acid sequence:

- FYIADHSFTARPTLRMFRISAVVATDKMTFTSGGTLFGDGCASSVA GEVMNC
 QTVLCILWTPFVFCPSIAVIIIICVFTSKALEAIWKWCRVERRPHIIEVDVLGKCP
 AF (SEQ ID NO:261), RPTLRMFRISAVVATDKMTFTSGGT (SEQ ID NO:262),
 20 PSIAVIIIICVFTSKALEAIWKWCRVER (SEQ ID NO:263), TSVSFHHRYSKSS
 DRPAHKVS (SEQ ID NO:264). Polynucleotides encoding these polypeptides are also encompassed by the invention.

This gene is expressed primarily in fetal lung, breast, and Hodgkin's Lymphoma II.

- 25 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, pulmonary, reproductive, immune, or hematopoietic disorders, particularly cell growth and differentiation conditions. Similarly, polypeptides and
 30 antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the fetal lung, breast, and tissues involved in Hodgkin's Lymphoma II expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g. Pulmonary,
 35 immune, reproductive, and cancerous and wounded tissues) or bodily fluids (e.g.lymph, amniotic fluid, pulmonary surfactant or sputum, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual

having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:142 as residues: Asn-32 to Asp-38, Thr-40 to Phe-46, Asn-53 to Gln-74, Ser-84 to Ile-91, Cys-95 to Glu-100, Ser-109 to Cys-121.

The tissue distribution in proliferating and differentiating tissues, combined with the homology to a urokinase receptor indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of cell growth and differentiation disorders, particularly of the lung, renal, breast, immune and endothelial tissues. Moreover, the expression within fetal tissue and other cellular sources marked by proliferating cells indicates that this protein may play a role in the regulation of cellular division, and may show utility in the diagnosis and treatment of cancer and other proliferative disorders. Similarly, developmental tissues rely on decisions involving cell differentiation and/or apoptosis in pattern formation. Thus this protein may also be involved in apoptosis or tissue differentiation and could again be useful in cancer therapy. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:17 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1168 of SEQ ID NO:17, b is an integer of 15 to 1182, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:17, and where b is greater than or equal to a + 14.

30

FEATURES OF PROTEIN ENCODED BY GENE NO: 8

The translation product of this gene shares sequence homology with cell adhesion molecules, which are implicated in cell migration, axonal guidance and fasciculation, and growth and tumorigenesis. When tested against U937 cell lines, supernatants removed from cells containing this gene activated the GAS (gamma activating sequence) promoter element. Thus, it is likely that this gene activates myeloid

35

cells, including their progenitors, through the JAK-STAT signal transduction pathway. GAS is a promoter element found upstream of many genes which are involved in the Jak-STAT pathway. The Jak-STAT pathway is a large, signal transduction pathway involved in the differentiation and proliferation of cells. Therefore, activation of the Jak-STAT pathway, reflected by the binding of the GAS element, can be used to indicate proteins involved in the proliferation and differentiation of cells. In specific embodiments, polypeptides of the invention comprise the following amino acid sequence:

RHNDFNKLSYTECNNMNKRMAKPEKKKGSVKSSLGIFLGPNCHLISSLFLFS
 10 VSLYPFATQF PFHYVLIFIIQAFGLCLPLTERQEAKSGLGGLCPDYTWPC
 PCLLVSCLSLRL (SEQ ID NO:265), CEVFSWHFPWSKLSPHLFLVSFLCIPL
 SLCHTV SFSLCSNIYNPGLRTMLAPHRETGGQVWAGWALSRLHVALPMSLG
 VLSLPAPTVTVVRMEGGDWKVCEQL GQCTYSHRMTK (SEQ ID NO:266),
 KRMAKPEKKKGSVKSSLGIFLG (SEQ ID NO:267), and/or YNPGLRTMLA
 15 PHRETGGQVWAGWALSRLHVA (SEQ ID NO:268). Polynucleotides encoding these polypeptides are also encompassed by the invention.

This gene is expressed primarily in the meninges, melanocytes, and to a lesser extent, in breast.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, neurodegenerative disease states and behavioral disorders, in addition to integumentary or reproductive disorders, particularly of the breast. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the nervous system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g. neural, integumentary, breast, reproductive, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:143 as residues: Asn-71 to Asp-79.

The tissue distribution in meninges combined with the homology to cell adhesion molecules and the detected GAS biological activity indicates that

polynucleotides and polypeptides corresponding to this gene are useful for the treatment and/or detection of neurodegenerative disease states and behavioural disorders such as Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder and panic disorder.

5 Moreover, the expression within melanocytes and breast tissue indicates polynucleotides and polypeptides corresponding to this gene are useful for the treatment, diagnosis, and/or prevention of various skin disorders including congenital disorders (i.e. nevi, moles, freckles, Mongolian spots, hemangiomas, port-wine syndrome), integumentary tumors (i.e. keratoses, Bowen's disease, basal cell carcinoma, squamous cell carcinoma, malignant melanoma, Paget's disease, mycosis fungoides, and Kaposi's sarcoma), injuries and inflammation of the skin (i.e. wounds, rashes, prickly heat disorder, psoriasis, dermatitis), atherosclerosis, urticaria, eczema, photosensitivity, autoimmune disorders (i.e. lupus erythematosus, vitiligo, dermatomyositis, morphea, scleroderma, pemphigoid, and pemphigus), keloids, striae, 10 erythema, petechiae, purpura, and xanthelasma. In addition, such disorders may predispose increased susceptibility to viral and bacterial infections of the skin (i.e. cold sores, warts, chickenpox, molluscum contagiosum, herpes zoster, boils, cellulitis, erysipelas, impetigo, tinea, athlete's foot, and ringworm). Moreover, the protein product of this gene may also be useful for the treatment or diagnosis of various 20 connective tissue disorders such as arthritis, trauma, tendonitis, chondromalacia and inflammation, autoimmune disorders such as rheumatoid arthritis, lupus, scleroderma, and dermatomyositis as well as dwarfism, spinal deformation, and specific joint abnormalities as well as chondrodysplasias (i.e. spondyloepiphyseal dysplasia congenita, familial osteoarthritis, Atelosteogenesis type II, metaphyseal chondrodysplasia type Schmid). Furthermore, This protein may show utility in 25 modulating the immune systems response to various degenerative neural conditions based upon the detected GAS biological activity. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

30 Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:18 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome.

35 Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1157 of SEQ ID NO:18, b is an integer of 15

to 1171, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:18, and where b is greater than or equal to a + 14.

5 FEATURES OF PROTEIN ENCODED BY GENE NO: 9

This gene is expressed primarily in fetal liver and spleen, and infant brain.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune, hematopoietic, neural, and developmental disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune and developmental systems, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g. Immune, hematopoietic, neural, developmental, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, amniotic fluid, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:144 as residues: Thr-187 to Lys-192, Asn-255 to Leu-262.

The tissue distribution of this gene in fetal liver spleen indicates a key role in the development of the immune system. Thus this gene could be used in the treatment and/or detection of immune disorders including arthritis, asthma, immunodeficiency diseases and leukemia. Expression in infant brain also indicates a role in the treatment and/or detection of neurodegenerative disease states and behavioural disorders such as Alzheimers Disease, Parkinsons Disease, Huntintons Disease, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder and panic disorder. Moreover, expression within fetal tissue and other cellular sources marked by proliferating cells indicates that this protein may play a role in the regulation of cellular division, and may show utility in the diagnosis and treatment of cancer and other proliferative disorders. Similarly, developmental tissues rely on decisions involving cell differentiation and/or apoptosis in pattern formation. Thus this protein may also be involved in apoptosis or tissue differentiation and could again be useful in cancer therapy. Protein, as well as,

antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:19 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1323 of SEQ ID NO:19, b is an integer of 15 to 1337, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:19, and where b is greater than or equal to a + 14.

15 FEATURES OF PROTEIN ENCODED BY GENE NO: 10

This gene is expressed primarily in breast, and to a lesser extent in liver.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, breast cancer, hepatoblastoma, hepatitis, liver metabolic diseases and conditions that are attributable to the differentiation of hepatocyte progenitor cells. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the breast and liver, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. breast, liver, cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:145 as residues: Gln-29 to Gly-38, Lys-57 to Asp-62.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the detection and treatment of liver disorders and cancers (e.g. hepatoblastoma, jaundice, hepatitis, liver metabolic diseases), and conditions that are attributable to the differentiation of hepatocyte progenitor cells. In

addition, the expression in breast would suggest a possible role in the detection and treatment of breast cancer.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:20 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1148 of SEQ ID NO:20, b is an integer of 15 to 1162, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:20, and where b is greater than or equal to a + 14.

15 FEATURES OF PROTEIN ENCODED BY GENE NO: 11

This gene is expressed primarily in brain, and to a lesser extent in retina.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, developmental, degenerative and behavioral diseases of the brain such as depression, schizophrenia, Alzheimer's disease, Parkinson's disease, Huntington's disease, specific brain tumors, aphasia, mania, depression, dementia, paranoia, addictive behavior and sleep disorders as well as conditions that affect vision and function of the eye such as retinoblastoma and cataracts. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the brain and eye, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g., brain, retina, cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:146 as residues: Pro-46 to Gln-60, Pro-68 to Gly-75, Leu-78 to Ala-86, Gln-93 to Asp-98.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of developmental, degenerative and behavioral diseases, and conditions of the brain such as aphasia, depression, schizophrenia, Alzheimer's disease, Parkinson's disease, Huntington's disease, specific brain tumors, mania, depression, dementia, paranoia, addictive behavior and sleep disorders. In addition, the expression in retina would also suggest a role for this gene product in the diagnosis and treatment of conditions that affect vision and function of the eye such as retinoblastoma, myopia, hyperopia and cataracts.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:21 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1823 of SEQ ID NO:21, b is an integer of 15 to 1837, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:21, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 12

The gene encoding the disclosed cDNA is thought to reside on chromosome 8. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 8. One embodiment of this gene comprises polypeptides of the following amino acid sequence:

MSPYASQGFPLPPYPPQEANRSITSLSVADTVSSSTTSHTTAKPAAPSFVLSN
LPLPIPTVDASIPTSQNGFGYKMPDVPDAFPELSELSVSQLTDMNEQEEVLLEQF
LTLPLQLKQIITDKDDL VKSIEELARKNLLLEPSLEAKRQTVLDKYELLTQMKSTF
EKKMQRQHELSECSASALQARLKVAAHEAEEEESDNIAEDFLEGKMEIDDFLSS
FMEKRTICHCRRAKEEKLQQAAMHSQFHAPL (SEQ ID NO:269), LPPYPPQE
ANRSITSLSVADTVS (SEQ ID NO:270), TAKPAAPSFVLSNLPLPIPTVDASIP
(SEQ ID NO:271), PDVPDAFPELSELSVSQLTDMNEQE (SEQ ID NO:272), QFLTL
PQLKQIITDKDDL VKSIEELARKN (SEQ ID NO:273), RQTVLDKYELLTQ MKS
TFEKKMQRQ (SEQ ID NO:274), ASALQARLKVAAHEAEEEESDNIAEDFLE (SEQ
ID NO:275), and/or MEKRTICHCRRAKEEKLQQAAMHSQF (SEQ ID NO: 276).

An additional embodiment is the polynucleotides encoding these polypeptides.

This gene is expressed primarily in breast and placenta, and to a lesser extent in a variety of other tissues and cell types.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, breast and endometrial cancers as well as prenatal disorders and deficiencies. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the breast and reproductive system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. breast, placental, cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the detection and treatment of breast cancer, ovarian and other endometrial cancers, infertility and pre-natal disorders. Furthermore, the tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for treating female infertility. The protein product is likely involved in preparation of the endometrium of implantation and could be administered either topically or orally. Alternatively, this gene could be transfected in gene-replacement treatments into the cells of the endometrium and the protein products could be produced. Similarly, these treatments could be performed during artificial insemination for the purpose of increasing the likelihood of implantation and development of a healthy embryo. In both cases this gene or its gene product could be administered at later stages of pregnancy to promote healthy development of the endometrium.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:22 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1040 of SEQ ID NO:22, b is an integer of 15

to 1054, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:22, and where b is greater than or equal to a + 14.

5 FEATURES OF PROTEIN ENCODED BY GENE NO: 13

This gene is expressed primarily in anergic T-cells.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune or hematopoietic disorders, particularly autoimmune disorders such as lupus. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:148 as residues: Lys-49 to Gln-57, Arg-63 to Ala-69.

The tissue distribution in T-cells indicates that the polypeptides or polynucleotides are useful for treatment, prophylaxis, and diagnosis of immune and autoimmune diseases, such as lupus, transplant rejection, allergic reactions, arthritis, asthma, immunodeficiency diseases, leukemia, and AIDS. The expression observed predominately in hematopoietic cells also indicates that the polynucleotides or polypeptides are important in treating and/or detecting hematopoietic disorders, such as graft versus host reaction, graft versus host disease, transplant rejection, myelogenous leukemia, bone marrow fibrosis, and myeloproliferative disease. The polypeptides or polynucleotides are also useful to enhance or protect proliferation, differentiation, and functional activation of hematopoietic progenitor cells (e.g., bone marrow cells), useful in treating cancer patients undergoing chemotherapy or patients undergoing bone marrow transplantation. The polypeptides or polynucleotides are also useful to increase the proliferation of peripheral blood leukocytes, which can be used in the combat of a range of hematopoietic disorders, including immunodeficiency diseases, leukemia,

and septicemia. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:23 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1052 of SEQ ID NO:23, b is an integer of 15 to 1066, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:23, and where b is greater than or equal to a + 14.

15 FEATURES OF PROTEIN ENCODED BY GENE NO: 14

The translation product of this gene shares sequence homology with a drought-induced protease inhibitor from soybean. As a result, the protein product of this gene may show utility in the treatment and/or prevention of a variety of proliferative disorders (e.g. for inhibition of key proteolytic events during cellular metabolism of the tumor which may lead to cessation of mitosis) or for the treatment of degenerative conditions where the inhibition of aberrant proteolysis may lead to cessation of degeneration and ultimately in immune protection.

This gene is expressed primarily in the kidney.

25 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, disorders of the kidney. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the urogenital system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. kidney, cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:149 as residues: Glu-48 to Arg-56, Ser-61 to Gly-66.

The tissue distribution in kidney tissue combined with the homology to a protease inhibitor indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of disorders affecting the kidney. Furthermore, this gene or gene product could be used in the treatment and/or detection of kidney diseases including renal failure, nephritis, renal tubular acidosis, proteinuria, pyuria, edema, pyelonephritis, hydronephritis, nephrotic syndrome, crush syndrome, glomerulonephritis, hematuria, renal colic and kidney stones, in addition to Wilms Tumor Disease, and congenital kidney abnormalities such as horseshoe kidney, polycystic kidney, and Falconi's syndrome. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:24 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 914 of SEQ ID NO:24, b is an integer of 15 to 928, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:24, and where b is greater than or equal to a + 14.

25

FEATURES OF PROTEIN ENCODED BY GENE NO: 15

When tested against U937 cell lines, supernatants removed from cells containing this gene activated the GAS (gamma activating sequence) promoter element. Thus, it is likely that this gene activates myeloid cells, including their progenitors, through the JAK-STAT signal transduction pathway. GAS is a promoter element found upstream of many genes which are involved in the Jak-STAT pathway. The Jak-STAT pathway is a large, signal transduction pathway involved in the differentiation and proliferation of cells. Therefore, activation of the Jak-STAT pathway, reflected by the binding of the GAS element, can be used to indicate proteins involved in the proliferation and differentiation of cells. In specific embodiments, polypeptides of the invention comprise the following amino acid sequence:

TRPVFLSMTPLKGIVSVLPQVFLCAYMAAFNSINGNRSYTCKPLERSLLMAGA
VASSTFLGVIPQFVQ (SEQ ID NO:277), PLKGIVSVLPQVFLCAYMAA (SEQ ID
NO:278), and/or AFNSINGNRSYTCKPLERSLL (SEQ ID NO:279). Polynucleotides
encoding these polypeptides are also encompassed by the invention. The gene encoding
5 the disclosed cDNA is believed to reside on chromosome 10. Accordingly,
polynucleotides related to this invention are useful as a marker in linkage analysis for
chromosome 10.

This gene is expressed primarily in B cell and T cell lymphomas.

Therefore, polynucleotides and polypeptides of the invention are useful as
10 reagents for differential identification of the tissue(s) or cell type(s) present in a
biological sample and for diagnosis of diseases and conditions which include, but are
not limited to, immune or hematopoietic disorders, particularly B cell and T cell
lymphomas, infections, multiple myeloma, immunodeficiencies, and inflammatory
conditions. Similarly, polypeptides and antibodies directed to these polypeptides are
15 useful in providing immunological probes for differential identification of the tissue(s)
or cell type(s). For a number of disorders of the above tissues or cells, particularly
immune or hematopoietic disorders, such as B- and T-cell lymphomas, expression of
this gene at significantly higher or lower levels may be routinely detected in certain
tissues and cell types (e.g. Immune, hematopoietic, and cancerous and wounded
20 tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal
fluid) or another tissue or cell sample taken from an individual having such a disorder,
relative to the standard gene expression level, i.e., the expression level in healthy tissue
or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID
25 NO:150 as residues: Phe-85 to Gly-96, Glu-133 to Thr-143.

The tissue distribution in B- and T-cell lymphomas, combined with the detected
GAS biological activity indicates that polynucleotides and polypeptides corresponding
to this gene are useful for the diagnosis and treatment of a variety of immune disorders,
particularly proliferative conditions such as cancer and leukemias. In addition,
30 polynucleotides and polypeptides corresponding to this gene are useful for the treatment
and diagnosis of hematopoietic related disorders such as anemia, pancytopenia,
leukopenia, thrombocytopenia or leukemia since stromal cells are important in the
production of cells of hematopoietic lineages. The uses include bone marrow cell ex
vivo culture, bone marrow transplantation, bone marrow reconstitution, radiotherapy or
35 chemotherapy of neoplasia. The gene product may also be involved in lymphopoiesis,
therefore, it can be used in immune disorders such as infection, inflammation, allergy,
immunodeficiency etc. In addition, this gene product may have commercial utility in the

expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

- 5 Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:25 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome.
- 10 Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 952 of SEQ ID NO:25, b is an integer of 15 to 966, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:25, and where b is greater than or equal to a + 14.

15

FEATURES OF PROTEIN ENCODED BY GENE NO: 16

- 20 The protein product of this gene was found to have homology to the Poly(A) polymerase of *Bos taurus*, which is known to be important in the creation of the 3' poly(A) tail of mRNA's. The gene encoding the disclosed cDNA is believed to reside on chromosome 14. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 14.

 This gene is expressed primarily in brain, and to a lesser extent, in prostate.

- 25 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, neural disorders, such as neurodegenerative disease states and behavioral conditions, in addition to reproductive disorders, particularly of the prostate. Similarly,
- 30 polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the nervous system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g. neural, reproductive, and cancerous and wounded
- 35 tissues) or bodily fluids (e.g. lymph, serum, seminal fluid, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a

disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:151 as residues: Glu-47 to Ser-52.

5 The tissue distribution in brain indicates that polynucleotides and polypeptides corresponding to this gene are useful for the detection/treatment of neurodegenerative disease states and behavioural disorders such as Alzheimers Disease, Parkinsons Disease, Huntingtons Disease, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder and panic disorder. Moreover, expression of the gene in prostate
10 indicates that polynucleotides and polypeptides corresponding to this gene are useful for the detection or treatment of prostate disorders including benign prostate hyperplasia, prostate cancer, and metabolic disorders. The homology to the PAP polyA polymerase indicates that the protein product of this gene, antibodies directed to this protein, or the gene encoding this protein via a gene therapy approach, may show utility as a
15 preventative therapy for proliferative conditions. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ
20 ID NO:26 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of
25 a-b, where a is any integer between 1 to 1132 of SEQ ID NO:26, b is an integer of 15 to 1146, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:26, and where b is greater than or equal to a + 14.

30 **FEATURES OF PROTEIN ENCODED BY GENE NO: 17**

This gene is expressed primarily in epididymus.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a
35 biological sample and for diagnosis of diseases and conditions which include, but are not limited to, diseases of the reproductive organs. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes

for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the reproductive system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. reproductive, cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:152 as residues: Met-1 to Pro-6, Glu-58 to Cys-63, Glu-65 to Gly-72, Thr-74 to Val-87.

The tissue distribution in epididymus indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of disorders of the epididymus and reproductive organs. Furthermore, the tissue distribution indicates that the protein product of this gene is useful for the treatment and diagnosis of conditions concerning proper testicular function (e.g. endocrine function, sperm maturation), as well as cancer. Therefore, this gene product is useful in the treatment of male infertility and/or impotence. This gene product is also useful in assays designed to identify binding agents as such agents (antagonists) are useful as male contraceptive agents. Similarly, the protein is believed to be useful in the treatment and/or diagnosis of testicular cancer. The testes are also a site of active gene expression of transcripts that may be expressed, particularly at low levels, in other tissues of the body. Therefore, this gene product may be expressed in other specific tissues or organs where it may play related functional roles in other processes, such as hematopoiesis, inflammation, bone formation, and kidney function, to name a few possible target indications.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:27 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 788 of SEQ ID NO:27, b is an integer of 15 to 802, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:27, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 18

5 This gene is expressed primarily in synovium and rhabdomyosarcoma.

 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, muscular skeletal system and cancer. Similarly, polypeptides and
10 antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the muscular skeletal system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. musculo-skeletal, cancerous and wounded tissues) or bodily
15 fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

 Preferred epitopes include those comprising a sequence shown in SEQ ID
20 NO:153 as residues: Trp-30 to Val-35, Lys-44 to Arg-49.

 The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of disorders of the muscular skeletal system and cancer. Furthermore, the expression of this gene product in synovium would suggest a role in the detection and treatment of disorders and
25 conditions affecting the skeletal system, in particular osteoporosis as well as disorders afflicting connective tissues (e.g. arthritis, trauma, tendonitis, chondromalacia and inflammation), such as in the diagnosis or treatment of various autoimmune disorders such as rheumatoid arthritis, lupus, scleroderma, and dermatomyositis as well as dwarfism, spinal deformation, and specific joint abnormalities as well as
30 chondrodysplasias (ie. spondyloepiphyseal dysplasia congenita, familial osteoarthritis, Atelosteogenesis type II, metaphyseal chondrodysplasia type Schmid). Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

 Many polynucleotide sequences, such as EST sequences, are publicly available
35 and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:28 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the

scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1155 of SEQ ID NO:28, b is an integer of 15
5 to 1169, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:28, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 19

10

The gene encoding the disclosed cDNA is thought to reside on chromosome 5. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 5.

This gene is expressed primarily in fetal liver/spleen, and to a lesser extent, in
15 tonsils.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune, hematopoietic, or hepatic disorders, particularly multiple
20 myeloma, immunodeficiencies, and cancers. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the hepatic system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell
25 types (e.g. immune, cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, bile, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

30 Preferred epitopes include those comprising a sequence shown in SEQ ID NO:154 as residues: Asp-27 to Ser-36.

Expression of this gene at either the RNA or protein level could be used as a diagnostic indicator of hepatic cancer. Furthermore, the tissue distribution in fetal liver and tonsil tissue indicates that polynucleotides and polypeptides corresponding to this
35 gene are useful for the diagnosis and treatment of a variety of immune system disorders. Moreover, the protein product of this gene may play a role in the regulation of the proliferation; survival; differentiation; and/or activation of potentially all

hematopoietic cell lineages, including blood stem cells. This gene product may be involved in the regulation of cytokine production, antigen presentation, or other processes that may also suggest a usefulness in the treatment of cancer (e.g. by boosting immune responses). Since the gene is expressed in cells of lymphoid origin, the gene or protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues. Therefore it may be also used as an agent for immunological disorders including arthritis, asthma, immune deficiency diseases such as AIDS, leukemia, rheumatoid arthritis, inflammatory bowel disease, sepsis, acne, and psoriasis. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:29 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1452 of SEQ ID NO:29, b is an integer of 15 to 1466, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:29, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 20

This gene is expressed primarily in human brain.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, disorders or diseases of the central nervous system. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the brain, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. brain, cancerous and wounded tissues) or bodily fluids (e.g.

lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

5 The tissue distribution in brain tissue indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and the treatment of CNS disorders. Furthermore, polynucleotides and polypeptides corresponding to this gene are useful for the detection/treatment of neurodegenerative disease states and behavioural disorders such as Alzheimers Disease, Parkinsons Disease, Huntingtons
10 Disease, Tourette Syndrome, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, panic disorder, learning disabilities, ALS, psychoses, autism, and altered bahaviors, including disorders in feeding, sleep patterns, balance, and preception. In addition, the gene or gene product may also play a role in the treatment and/or detection of developmental disorders associated with the developing embryo,
15 sexually-linked disorders, or disorders of the cardiovascular system. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ
20 ID NO:30 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of
25 a-b, where a is any integer between 1 to 1212 of SEQ ID NO:30, b is an integer of 15 to 1226, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:30, and where b is greater than or equal to a + 14.

30 **FEATURES OF PROTEIN ENCODED BY GENE NO: 21**

In specific embodiments, polypeptides of the invention comprise the following amino acid sequence:

MSDFEKVDISVHQHIHVGPLLLMTTESWGPSCAPSPALLSGHTAASFHTLGG
35 VLGCPPYHKFYSS AHTSDHRKETNKVEEGRWVDVTRSLGNFNFRRKFCC
VSELLICGIFLDSSWKLQINSNDCKVL (SEQ ID NO:280), VGPLLLMTTESW
GPSCAPSPALLSGHTAAS (SEQ ID NO:281), and/or ETNKVEEGRWVDVTRS

LGNFNFRRKFF (SEQ ID NO:282). Polynucleotides encoding these polypeptides are also encompassed by the invention.

This gene is expressed primarily in fetal spleen or liver, adult spleen, and to a lesser extent, in activated T-cells.

5 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune or hematopoietic disorders, particularly abnormal proliferation or activation of hematopoietic cells, particularly of T-cells and their progenitors.

10 Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g. Immune, hematopoietic, and

15 cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

20 Preferred epitopes include those comprising a sequence shown in SEQ ID NO:156 as residues: Arg-19 to Phe-24, Ala-44 to Asp-51, Glu-60 to Ile-66.

The tissue distribution in spleen tissues and T-cells indicates that polynucleotides and polypeptides corresponding to this gene are useful for modulating or detecting the abnormal proliferation or activation of T-cells and immune cell

25 precursor cells. Moreover, expression within fetal spleen indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of hematopoietic related disorders such as anemia, pancytopenia, leukopenia, thrombocytopenia or leukemia since stromal cells are important in the production of cells of hematopoietic lineages. The uses include bone marrow cell ex vivo culture,

30 bone marrow transplantation, bone marrow reconstitution, radiotherapy or chemotherapy of neoplasia. The gene product may also be involved in lymphopoiesis, therefore, it can be used in immune disorders such as infection, inflammation, allergy, immunodeficiency etc. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the

35 differentiation and/or proliferation of various cell types. Similarly, This gene product may be involved in the regulation of cytokine production, antigen presentation, or other processes that may also suggest a usefulness in the treatment of cancer (e.g. by

boosting immune responses). Since the gene is expressed in cells of lymphoid origin, the natural gene product may be involved in immune functions. Therefore it may be also used as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's disease, scleroderma and tissues. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:31 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1080 of SEQ ID NO:31, b is an integer of 15 to 1094, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:31, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 22

This gene is expressed primarily in the amygdala.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, developmental, degenerative and behavioral diseases of the brain such as depression, schizophrenia, Alzheimer's disease, Parkinson's disease, Huntington's disease, specific brain tumors, aphasia, mania, depression, dementia, paranoia, addictive behavior and sleep disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential

identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the brain, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. brain, cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:157 as residues: Pro-94 to Ala-107.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of developmental, degenerative and behavioral diseases and conditions of the brain such as aphasia, depression, schizophrenia, Alzheimer's disease, Parkinson's disease, Huntington's disease, specific brain tumors, mania, depression, dementia, paranoia, addictive behavior and sleep disorders.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:32 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1023 of SEQ ID NO:32, b is an integer of 15 to 1037, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:32, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 23

30

The translation product of this gene shares sequence homology with octaprenyltransferase, which is thought to be important in cellular respiration and metabolism. When tested against fibroblast cell lines, supernatants removed from cells containing this gene activated the EGR1 assay. Thus, it is likely that this gene activates fibroblast cells through a signal transduction pathway. Early growth response 1 (EGR1) is a promoter associated with certain genes that induces various tissues and cell types upon activation, leading the cells to undergo differentiation and proliferation. The

gene encoding the disclosed cDNA is thought to reside on chromosome 4. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 4.

5 This gene is expressed primarily in synovium, liver cells, dendritic cells and stromal cells.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, metabolic and respiratory disorders, immune disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the metabolic processes and immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. immune, liver, cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

20 Preferred epitopes include those comprising a sequence shown in SEQ ID NO:158 as residues: Asp-54 to Asn-69, His-176 to Asp-181, Phe-194 to Trp-201, Ser-220 to Pro-225, Arg-248 to Trp-253, Trp-276 to Ile-288.

The tissue distribution and homology to octaprenyltransferase indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of metabolic and respiratory disorders. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:33 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1362 of SEQ ID NO:33, b is an integer of 15 to 1376, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:33, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 24

5 This gene is expressed primarily in activated T cells and in the spleen from a patient suffering from lymphocytic leukemia.

 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune or hematopoietic disorders, particularly immunodeficiencies, multiple myeloma, and leukemias. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

 The tissue distribution in T-cells and spleen tissue indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis or treatment of leukemia. Furthermore, the tissue distribution indicates that the polypeptides or polynucleotides are useful for treatment, prophylaxis, and diagnosis of immune and autoimmune diseases, such as lupus, transplant rejection, allergic reactions, arthritis, asthma, immunodeficiency diseases, leukemia, and AIDS. The expression observed predominantly in hematopoietic cells also indicates that the polynucleotides or polypeptides are important in treating and/or detecting hematopoietic disorders, such as graft versus host reaction, graft versus host disease, transplant rejection, myelogenous leukemia, bone marrow fibrosis, and myeloproliferative disease. The polypeptides or polynucleotides are also useful to enhance or protect proliferation, differentiation, and functional activation of hematopoietic progenitor cells (e.g., bone marrow cells), useful in treating cancer patients undergoing chemotherapy or patients undergoing bone marrow transplantation. The polypeptides or polynucleotides are also useful to increase the proliferation of peripheral blood leukocytes, which can be used in the combat of a range of hematopoietic disorders, including immunodeficiency diseases, leukemia,

and septicemia. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:34 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1206 of SEQ ID NO:34, b is an integer of 15 to 1220, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:34, and where b is greater than or equal to a + 14.

15 FEATURES OF PROTEIN ENCODED BY GENE NO: 25

This gene is expressed primarily in bone marrow.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune or hematopoietic disorders, particularly disorders afflicting stem cell or myeloid progenitors, and in particular multiple myeloma, immunodeficiencies, or SCID. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune and hematopoietic systems, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. immune, cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in bone marrow indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of disorders affecting the immune and hematopoietic systems. In addition, The protein product of this gene is useful for the diagnosis and/or treatment of hematopoietic disorders. Furthermore, this gene product is primarily expressed in hematopoietic cells

and tissues, suggesting that it plays a role in the survival, proliferation, and/or differentiation of hematopoietic lineages. This is particularly supported by the expression of this gene product in bone marrow, which is a primary sites of definitive hematopoiesis. The uses include bone marrow cell ex vivo culture, bone marrow
5 transplantation, bone marrow reconstitution, radiotherapy or chemotherapy of neoplasia. The gene product may also be involved in lymphopoiesis, therefore, it can be used in immune disorders such as infection, inflammation, allergy, immunodeficiency etc. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation
10 and/or proliferation of various cell types. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ
15 ID NO:35 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of
20 a-b, where a is any integer between 1 to 1332 of SEQ ID NO:35, b is an integer of 15 to 1346, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:35, and where b is greater than or equal to a + 14.

25 **FEATURES OF PROTEIN ENCODED BY GENE NO: 26**

This gene is expressed primarily in the cells of the immune system.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a
30 biological sample and for diagnosis of diseases and conditions which include, but are not limited to, disorders of the immune systems, such as AIDS. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of
35 this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or

another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:161 as residues: His-17 to Ser-24, Glu-53 to Asn-58, Glu-66 to Lys-72.

The tissue distribution in immune cells indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of a variety of immune system disorders. Further, the expression of this gene product indicates a role in the regulation of the proliferation; survival; differentiation; and/or activation of potentially all hematopoietic cell lineages, including blood stem cells. This gene product may be involved in the regulation of cytokine production, antigen presentation, or other processes that may also suggest a usefulness in the treatment of cancer (e.g. by boosting immune responses). Since the gene is expressed in cells of lymphoid origin, the gene or protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues. Therefore it may be also used as an agent for immunological disorders including arthritis, asthma, immune deficiency diseases such as AIDS, leukemia, rheumatoid arthritis, inflammatory bowel disease, sepsis, acne, and psoriasis. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:36 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1012 of SEQ ID NO:36, b is an integer of 15 to 1026, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:36, and where b is greater than or equal to a + 14.

35

FEATURES OF PROTEIN ENCODED BY GENE NO: 27

The translation product of this gene shares sequence homology with glucan synthetase which is thought to be important in modifying carbohydrate moieties on extracellular molecules.

5 This gene is expressed primarily in T-cells.

 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune or hematopoietic disorders, particularly autoimmune diseases
10 and inflammation. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g. Immune,
15 hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

20 Preferred epitopes include those comprising a sequence shown in SEQ ID NO:162 as residues: Gly-33 to Leu-39, Thr-69 to Ser-77, Arg-102 to Thr-109.

 The tissue distribution in T-cells combined with the homology to glucan synthetase indicates that polynucleotides and polypeptides corresponding to this gene are useful for modifying the response and production of active cytokines by T cells, in
25 modulating cell-cell interactions, or cell-tissue interactions, and in inflammatory conditions. Alternatively, this gene product may be involved in the regulation of cytokine production, antigen presentation, or other processes that may also suggest a usefulness in the treatment of cancer (e.g. by boosting immune responses). Since the gene is expressed in cells of lymphoid origin, the natural gene product may be involved
30 in immune functions. Therefore it may be also used as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues; such as
35 host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's disease,

scleroderma and tissues. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:37 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 818 of SEQ ID NO:37, b is an integer of 15 to 832, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:37, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 28

In specific embodiments, polypeptides of the invention comprise the following amino acid sequence:

GRGDKPRQDRPASLRLKGPPSCQAPASHSSTLSSHCPCSLFACGSVWPGSLGS
GIFARLSQLLPSPASWG WDFLTLRQAQQMLGPSLCPGHSTSAHQHYGAYVLP
RDLCSFLLTSTVQGTAPLKNSRVTCIGSQVPLC (SEQ ID NO:283), AEVTSPA
KTDLQVFVSRDLPHARPLPLTAAPFPLIVPVPFLPVDLFGQGPWGQEYLQDSAS
SFPAQPLGA GTFSPCGRHNRCDPVSAQVTAQVHISTMGPMSCPETSAPSC
SHPQFRARRPSRTPEPVSSAPSKCLFV YDVPLL (SEQ ID NO:284), SLRLKGP
PSCQAPASHSSTLSSHCPCSLFA (SEQ ID NO:285), QQMLGPSLCPGHS TSAH
QHYGAYVLPRDLC (SEQ ID NO:286), DLQVFVSRDLPHARPLPLTAAPF PLIV
PVPF (SEQ ID NO:287), AQVHISTMGPMSCPETSAPSCSHPQFRARRP SRTPE
SPV (SEQ ID NO:288), and/or QAPPRQTCKSSSQGTSL (SEQ ID NO:289).
Polynucleotides encoding these polypeptides are also encompassed by the invention.

This gene is expressed primarily in endometrial tumors, fetal spleen, and to a lesser extent, in activated monocytes and T-cells.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are

not limited to, reproductive, immune, hematopoietic disorders, particularly pregnancy defects. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the reproductive and immune systems, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g. reproductive, endometrial, immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, amniotic fluid, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:163 as residues: Ser-66 to Thr-75.

The tissue distribution in endometrial tissue indicates that the protein product of this gene could be used in the treatment and/or detection of pregnancy associated disorders including miscarriage, and endometriosis. Alternatively, expression in hematopoietic cells indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and/or detection of immune system related diseases including arthritis, asthma, immunodeficiency diseases and leukemia. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:38 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 692 of SEQ ID NO:38, b is an integer of 15 to 706, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:38, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 29

In specific embodiments, polypeptides of the invention comprise the following amino acid sequence:

AALRPSGSLAGPEWPWQHWCGCWREHXVKPQQVDLHSARLWAAPAAVGPA
 HAGGSPGMPPGGTAPHARRH SLPSPTAQSHLWHVHGLRQRGPKAVPLDLAQ
 LVTTTTPLFXLALSALLGRRHHPLQLAAMGPLCLGAAC SLAGEFRTPT
 GCGFLAATCLRGLKSVQQSALLQEERLDAVTLLYATSLPSFCLLAGAALVLEA
 5 GVAPP TAGDSRLWACILLSCLLSVLYNLASFLLALTSALT VHVLGNLTVV
 GNLILSRLLFGSRLSALS YVGIA LTLSGMFLYHNCEFVASWAARRGLW
 RRDQPSKGL (SEQ ID NO:290), GQPSGPPAAWPGPSGHGSTGVAAGGSTXSSL
 NKWIFTVHGFGRP LLLSALHMLVAALACHRGARRP (SEQ ID NO:291), WPGPS
 GHGSTGVAAGGSTXSS (SEQ ID NO:292), EWPWQHWCGCWREHXVKPQQVD
 10 LHSA (SEQ ID NO:293), QQSALLQEERLDAVTLLYATSLPSFCLL (SEQ ID
 NO:294), ACILLSCLLSVLYNLASFLLALTSAL (SEQ ID NO:295), and/or
 SLNKWIFTVHGFGRP LLLSAL (SEQ ID NO:296). Polynucleotides encoding these
 polypeptides are also encompassed by the invention.

This gene is expressed primarily in brain tissue from a patient suffering from
 15 Alzheimer's disease (spongy change), and to a lesser extent, in human umbilical vein
 and human pancreas tumor tissue.

Therefore, polynucleotides and polypeptides of the invention are useful as
 reagents for differential identification of the tissue(s) or cell type(s) present in a
 biological sample and for diagnosis of diseases and conditions which include, but are
 20 not limited to, developmental, immune, metabolic, digestive or neural disorders, such
 as Alzheimer's disease, in addition to cancers and tumors. Similarly, polypeptides and
 antibodies directed to these polypeptides are useful in providing immunological probes
 for differential identification of the tissue(s) or cell type(s). For a number of disorders
 of the above tissues or cells, particularly of the immune and secretory systems,
 25 expression of this gene at significantly higher or lower levels may be routinely detected
 in certain tissues and cell types (e.g. developmental, immune, metabolic, digestive,
 cancerous and wounded tissues) or bodily fluids (e.g. lymph, bile, amniotic fluid,
 serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample
 taken from an individual having such a disorder, relative to the standard gene
 30 expression level, i.e., the expression level in healthy tissue or bodily fluid from an
 individual not having the disorder.

The tissue distribution in brain tissue indicates that polynucleotides and
 polypeptides corresponding to this gene are useful for diagnosis and treatment of
 Alzheimer's disease, and immune and secretory system disorders such as cancers.
 35 Moreover, polynucleotides and polypeptides corresponding to this gene are useful for
 the detection/treatment of neurodegenerative disease states, behavioural disorders, or
 inflammatory conditions such as Parkinsons Disease, Huntingtons Disease, Tourette

Syndrome, meningitis, encephalitis, demyelinating diseases, peripheral neuropathies, neoplasia, trauma, congenital malformations, spinal cord injuries, ischemia and infarction, aneurysms, hemorrhages, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and preception. In addition, elevated expression of this gene product in regions of the brain indicates that it plays a role in normal neural function. Potentially, this gene product is involved in synapse formation, neurotransmission, learning, cognition, homeostasis, or neuronal differentiation or survival. Moreover, the gene or gene product may also play a role in the treatment and/or detection of developmental disorders associated with the developing embryo, sexually-linked disorders, or disorders of the cardiovascular system. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:39 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome.

Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1333 of SEQ ID NO:39, b is an integer of 15 to 1347, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:39, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 30

This gene is expressed primarily in neutrophils.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, infection and inflammation. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell

types (e.g. immune, cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:165 as residues: Asn-43 to Ala-49.

The tissue distribution in neutrophils indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of infection and inflammation related immune diseases. Furthermore, the gene product may also be involved in lymphopoiesis, therefore, it can be used in immune disorders such as infection, inflammation, allergy, immunodeficiency, etc. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Additionally, expression of this gene product in neutrophils also strongly indicates a role for this protein in immune function and immune surveillance.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:40 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1453 of SEQ ID NO:40, b is an integer of 15 to 1467, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:40, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 31

The translation product of this gene shares sequence homology with Ly6C antigen, which is thought to be important in T-cell activation. In specific embodiments, polypeptides of the invention comprise the following amino acid sequence:

KSTLSAAVVATILRTLA (SEQ ID NO:297), GDHSEQCLIKEMGARERRFCKAR
 GYRDTG REAQAKAGGRRGSQWNESQCS SQRPRPAKEVRKTRPRAGVGRGP
 ALLQLSLLQQVVLVYVRPSLRLVWLKA S (SEQ ID NO:298), MERGEYGGWG
 TYGSLDLGSQLCTVRSSGPCGSLHWGQH RSPISGPDNPSSSR GQQSIGSK

VGSPRSQWRSWKEVGRDPEKGE (SEQ ID NO:299), QAKAGGRRGSQWNESQ
CSSQRPR (SEQ ID NO:300), VGRGPALLQL SLLQQVVLYVRPSLRL (SEQ ID
NO:301), YGSLDLGSQ LCTVRSSGPGSL (SEQ ID NO:302), and/or KVGSPSR
SQWRSWKEVGRDP (SEQ ID NO:303). Polynucleotides encoding these polypeptides
5 are also encompassed by the invention.

This gene is expressed primarily in bone cancer, fetal brain, lung, and adipose
tissue.

Therefore, polynucleotides and polypeptides of the invention are useful as
reagents for differential identification of the tissue(s) or cell type(s) present in a
10 biological sample and for diagnosis of diseases and conditions which include, but are
not limited to, skeletal, developmental, pulmonary, or metabolic disorders, particular
disorders in the immune responses to the above conditions, such as in autoimmunities.
Similarly, polypeptides and antibodies directed to these polypeptides are useful in
providing immunological probes for differential identification of the tissue(s) or cell
15 type(s). For a number of disorders of the above tissues or cells, particularly of the
immune system, expression of this gene at significantly higher or lower levels may be
routinely detected in certain tissues or cell types (e.g. skeletal, developmental,
pulmonary, metabolic, and cancerous and wounded tissues) or bodily fluids (e.g.
lymph, amniotic fluid, pulmonary surfactant or sputum, serum, plasma, urine, synovial
20 fluid and spinal fluid) or another tissue or cell sample taken from an individual having
such a disorder, relative to the standard gene expression level, i.e., the expression level
in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID
NO:166 as residues: Gln-37 to Gln-45, Phe-76 to Leu-83, Thr-89 to Thr-105.

25 The tissue distribution combined with the homology to the Ly6C T-cell
activation antigen indicates that polynucleotides and polypeptides corresponding to this
gene are useful for the diagnosis and intervention of immune related disorders. The
tissue distribution in tissues particularly active in immune reaction, for example bone
cancer, indicate that this gene may also be involved in T-cell activation. Thus the gene
30 product can be used either for the development of immune suppressants, or modulators,
for immune responses. Moreover, the expression within brain tissue indicates that the
protein is useful for the treatment and/or prevention of neurodegenerative disorders,
particularly, but not limited to, Alzheimer's or Parkinson's disease. Alternatively, the
expression within fetal tissue and other cellular sources marked by proliferating cells
35 indicates that this protein may play a role in the regulation of cellular division, and may
show utility in the diagnosis and treatment of cancer and other proliferative disorders.
Similarly, developmental tissues rely on decisions involving cell differentiation and/or

apoptosis in pattern formation. Thus this protein may also be involved in apoptosis or tissue differentiation and could again be useful in cancer therapy. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

- 5 Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:41 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome.
- 10 Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 900 of SEQ ID NO:41, b is an integer of 15 to 914, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:41, and where b is greater than or equal to a + 14.

15

FEATURES OF PROTEIN ENCODED BY GENE NO: 32

- 20 The gene encoding the disclosed cDNA is thought to reside on chromosome 12. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 12.

 This gene is expressed primarily in brain, keratinocytes and fibroblasts.

- Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, disorders of the brain and epidermal system. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the epidermal and neural systems,
- 25 expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. skin, brain, cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily
- 30 fluid from an individual not having the disorder.

35

 The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of diseases of the

neural and epidermal systems. Furthermore, the tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the detection/treatment of neurodegenerative disease states and behavioural disorders such as Alzheimers Disease, Parkinsons Disease, Huntingtons Disease, Tourette Syndrome, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, panic disorder, learning disabilities, ALS, psychoses, autism, and altered bahaviors, including disorders in feeding, sleep patterns, balance, and perception. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues. Additionally, the tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment, diagnosis, and/or prevention of various skin disorders including congenital disorders (i.e. nevi, moles, freckles, Mongolian spots, hemangiomas, port-wine syndrome), integumentary tumors (i.e. keratoses, Boweni's disease, basal cell carcinoma, squamous cell carcinoma, malignant melanoma, Pageti's disease, mycosis fungoides, and Kaposii's sarcoma), injuries and inflammation of the skin (i.e. wounds, rashes, prickly heat disorder, psoriasis, dermatitis), atherosclerosis, uticaria, eczema, photosensitivity, autoimmune disorders (i.e. lupus erythematosus, vitiligo, dermatomyositis, morphea, scleroderma, pemphigoid, and pemphigus), keloids, striae, erythema, petechiae, purpura, and xanthelasma. Moreover, such disorders may predispose increased susceptibility to viral and bacterial infections of the skin (i.e. cold sores, warts, chickenpox, molluscum contagiosum, herpes zoster, boils, cellulitis, erysipelas, impetigo, tinea, athletes foot, and ringworm).

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:42 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1117 of SEQ ID NO:42, b is an integer of 15 to 1131, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:42, and where b is greater than or equal to a + 14.

35 FEATURES OF PROTEIN ENCODED BY GENE NO: 33

The translation product of this gene shares sequence homology with a sodium dependent sulfate transporter which is thought to be important in sulfate uptake by cells. The gene encoding the disclosed cDNA is thought to reside on chromosome 7.

Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 7. One embodiment of this gene comprises polypeptides of

the following amino acid sequence:
MPQSLSSLASSSSSFQRXKPCFGKKNDGENQEHS LGTEPIITWKDFQKTMPWE
IVILVGGGYALASGSKSSGLSTWIGNQMLSSLPPWAVTLLACILVSIVTEFVS
NPATITIFLPILCSLSETLHINPLYTLIPVTMCISFAVMLPVGNNPPNAIVFSYGHQC
10 IKDMVKAGLG VNVIGLVIVMVAINTWGVSLFHLDTYPAWARVSNITDQA (SEQ
ID NO:304), NDGENQEHS LGTEPIITWKDFQK (SEQ ID NO:305), IGNQMLSSLSS
LPPWAVTLLACILV (SEQ ID NO:306), ATITIFLPILCSLSETLHINPLYTLIP (SEQ
ID NO:307), LPVGNNPPNAIVFSYGHQCQIKDMVKAG (SEQ ID NO:308), and/or
LVIVMVAINTWGVSLFHLDTYPAWARVSN (SEQ ID NO:309). An additional
15 embodiment is the polynucleotides encoding these polypeptides.

This gene is expressed primarily in placenta, and to a lesser extent, in infant brain and spinal cord.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a
20 biological sample and for diagnosis of diseases and conditions which include, but are not limited to, metabolic, reproductive, or central nervous system disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the central nervous
25 system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. CNS, reproductive, metabolic, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, amniotic fluid, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression
30 level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in placental and neural tissues, combined with the homology to a sodium dependent sulfate transporter indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment of metabolic
35 disorders involving sodium and sulfate metabolism and CNS disorders involving neuronal signalling abnormalities. Moreover, polynucleotides and polypeptides corresponding to this gene are useful for the detection/treatment of neurodegenerative

disease states, behavioural disorders, or inflammatory conditions such as Alzheimers Disease, Parkinsons Disease, Huntingtons Disease, Tourette Syndrome, meningitis, encephalitis, demyelinating diseases, peripheral neuropathies, neoplasia, trauma, congenital malformations, spinal cord injuries, ischemia and infarction, aneurysms, hemorrhages, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and preception. In addition, elevated expression of this gene product in regions of the brain indicates that it plays a role in normal neural function. Potentially, this gene product is involved in synapse formation, neurotransmission, learning, cognition, homeostasis, or neuronal differentiation or survival. Moreover, the gene or gene product may also play a role in the treatment and/or detection of developmental disorders associated with the developing embryo, sexually-linked disorders, or disorders of the cardiovascular system. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:43 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1319 of SEQ ID NO:43, b is an integer of 15 to 1333, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:43, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 34

Contact of cells with supernatant expressing the product of this gene increases the permeability of bovine chondrocyte cells to calcium. Thus, it is likely that the product of this gene is involved in a signal transduction pathway that is initiated when the product of this gene binds a receptor on the surface of the chondrocyte cell. Thus, polynucleotides and polypeptides have uses which include, but are not limited to, activating chondrocyte cells.

This gene is expressed primarily in CD34 positive cells.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune or reproductive disorders, particularly diseases related to lymphocytes. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. bone, immune, hematopoietic, reproductive, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, amniotic fluid, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:169 as residues: Leu-26 to Arg-32, Asn-40 to Ser-46.

The tissue distribution in CD34 positive cells indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis or treatment of the diseases of the immune system particularly those related to T lymphocytes. Furthermore, the tissue distribution, as well as the detected calcium flux biological activity data, suggest that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and/or treatment of bone and hematopoietic disorders. The ability of the translation product of this gene to induce a calcium flux in chondrocytes indicates that it may play a role in the survival, proliferation, and/or growth of bone. Therefore, it may be useful in influencing bone mass in such conditions as osteoporosis. More generally, as evidenced by expression in CD34 positive cells, this gene may play a role in the survival, proliferation, and/or differentiation of hematopoietic cells, and may be of use in the augmentation of the numbers of stem cells and committed progenitors. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:44 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more

polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 990 of SEQ ID NO:44, b is an integer of 15 to 1004, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:44, and where b is greater than or equal to a + 14.

5

FEATURES OF PROTEIN ENCODED BY GENE NO: 35

The gene encoding the disclosed cDNA is thought to reside on chromosome 9. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 9.

This gene is expressed primarily in the brain, and to a lesser extent, in liver.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, disorders affecting the brain, central nervous system, or liver, including cancer. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune, hematopoietic, or central nervous systems, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. brain, liver, immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, bile, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in brain and liver tissue indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of disorders affecting the immune, hematopoietic, or central nervous systems. Furthermore, polynucleotides and polypeptides corresponding to this gene are useful for the detection/treatment of neurodegenerative disease states and behavioural disorders such as Alzheimers Disease, Parkinsons Disease, Huntingtons Disease, Tourette Syndrome, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and preception. In addition, the gene or gene product may also play a role in the treatment and/or detection of developmental disorders associated with the developing embryo. Alternatively, the

expression within hepatic tissue indicates polynucleotides and polypeptides corresponding to this gene are useful for the detection and treatment of liver disorders and cancers (e.g. hepatoblastoma, jaundice, hepatitis, liver metabolic diseases and conditions that are attributable to the differentiation of hepatocyte progenitor cells).

- 5 Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:45 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1480 of SEQ ID NO:45, b is an integer of 15 to 1494, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:45, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 36

20

When tested against U937 and Jurkat cell lines, supernatants removed from cells containing this gene activated the GAS (gamma activating sequence) promoter element. Thus, it is likely that this gene activates myeloid and T-cells through the JAK-STAT signal transduction pathway. GAS is a promoter element found upstream of many genes which are involved in the Jak-STAT pathway. The Jak-STAT pathway is a large, signal transduction pathway involved in the differentiation and proliferation of cells. Therefore, activation of the Jak-STAT pathway, reflected by the binding of the GAS element, can be used to indicate proteins involved in the proliferation and differentiation of cells. In specific embodiments, polypeptides of the invention comprise the following amino acid sequence:

ETCPSNGIELRQAPTSLYILLHHIQPTPTHMLGRSYVLPAFSXNXEHGGLPNQI
 PKGDRNGNIRHSRIT FPCSSSTLQPESH LGFIRSKLHGLVRPGKDLRGRRSL
 QLSKHSLSSTCYMLRWET YKQVSYTAV (SEQ ID NO:310), QRHQENDKRN VH
 RFLHTCVHMPMCTHTHTQAVLSTWEGQFSNVASFTSLKRIPLSII YHSSHSP
 35 RRFVKVCQLRQEKALELTEVYVSASLKLQLYHLHCHFHTAV (SEQ ID NO:311),
 RQAPTSLYILLHHIQPTPTHMLG (SEQ ID NO:312), SHLGFIRSKLHGLVRPG
 KDLRGRRS (SEQ ID NO:313), RNVHRFLHTCVHMPMCTHTHTQ (SEQ ID

NO:314), and/or QLRQEKALELTEVYVSASLKLQLYH (SEQ ID NO:315).

Polynucleotides encoding these polypeptides are also encompassed by the invention.

This gene is expressed primarily in neutrophils.

Therefore, polynucleotides and polypeptides of the invention are useful as
5 reagents for differential identification of the tissue(s) or cell type(s) present in a
biological sample and for diagnosis of diseases and conditions which include, but are
not limited to, diseases of the immune system, particularly neutropenia, cancer,
inflammatory diseases and allergies. Similarly, polypeptides and antibodies directed to
these polypeptides are useful in providing immunological probes for differential
10 identification of the tissue(s) or cell type(s). For a number of disorders of the above
tissues or cells, particularly of the immune system, expression of this gene at
significantly higher or lower levels may be routinely detected in certain tissues and cell
types (e.g. Immune, hematopoietic, and cancerous and wounded tissues) or bodily
fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another
15 tissue or cell sample taken from an individual having such a disorder, relative to the
standard gene expression level, i.e., the expression level in healthy tissue or bodily
fluid from an individual not having the disorder.

The tissue distribution in neutrophils, combined with the detected GAS
biological activity indicates that polynucleotides and polypeptides corresponding to this
20 gene are useful for treatment/diagnosis of diseases of the immune system since
expression is primarily in neutrophils, and may be useful as a growth factor for the
differentiation or proliferation of neutrophils for the treatment of neutropenia following
chemotherapy or may be useful in the treatment of immune dysfunction or anti-
inflammatory by inhibiting infiltration of neutrophils to the site of injury or distress.
25 Protein, as well as, antibodies directed against the protein may show utility as a tumor
marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available
and accessible through sequence databases. Some of these sequences are related to SEQ
ID NO:46 and may have been publicly available prior to conception of the present
30 invention. Preferably, such related polynucleotides are specifically excluded from the
scope of the present invention. To list every related sequence is cumbersome.
Accordingly, preferably excluded from the present invention are one or more
polynucleotides comprising a nucleotide sequence described by the general formula of
a-b, where a is any integer between 1 to 1152 of SEQ ID NO:46, b is an integer of 15
35 to 1166, where both a and b correspond to the positions of nucleotide residues shown
in SEQ ID NO:46, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 37

5 In specific embodiments, polypeptides of the invention comprise the following amino acid sequence: PRVRGRKEPGCLGPGRAGGDSQKEIGSWQQM (SEQ ID NO:316), LSKGNRIMAADDDNGDGTSLFDVFSASPLKNNDEGSLDIYA GLDSAVSDSA SKSCVPSRNCLDLYEELTEEGTAKEATYNDLQVEYGKCQ LQMKELMKKFKEIQTQNFSLINENQSLKKN ISALIKTARVEINRKDEEI
10 SNLHQKIVLSFHIFEIHKLQGHILQLKQKILNLDLHIWMIVQRLITRAKS DVSKD VHHSTSLPNLEKEGKPHSDKRSTSHLPTSVEKHCTNGVWSRSHYQVGEGSSN EDSRRGRKDIRHS QFNRGTERVRKDLSTGCGDGEPRILEASQRLQGTS (SEQ ID NO:317), NRIMAADDDNGDGTSLFDVFSASPLKN (SEQ ID NO:318), CLDLY EEILTEEGTAKEATYNDL (SEQ ID NO:319), DEEISNLHQKIVLSFHIFEIHKLQG
15 (SEQ ID NO:320), EKEGKPHSDKRSTSHLPTSVEK (SEQ ID NO:321), and/or TERVRKDLSTGCGDGEPRILEASQRL (SEQ ID NO:322). Polynucleotides encoding these polypeptides are also encompassed by the invention.

This gene is expressed primarily in activated T cells.

20 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune and inflammatory disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders
25 of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression
30 level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

35 The tissue distribution in T-cells indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of immune and inflammatory disorders. Furthermore, expression of this gene product in tonsils indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. This gene product may be involved in the regulation of cytokine production, antigen presentation, or other

processes that may also suggest a usefulness in the treatment of cancer (e.g. by boosting immune responses). Since the gene is expressed in cells of lymphoid origin, the natural gene product may be involved in immune functions. Therefore it may be also used as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's disease, scleroderma and tissues. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:47 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1522 of SEQ ID NO:47, b is an integer of 15 to 1536, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:47, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 38

Contact of cells with supernatant expressing the product of this gene has been shown to increase the permeability of the plasma membrane of chondrocytes to calcium. Thus it is likely that the product of this gene is involved in a signal transduction pathway that is initiated when the product binds a receptor on the surface of the plasma membrane of both chondrocytes, in addition to other cell lines or tissue cell types. Thus, polynucleotides and polypeptides have uses which include, but are not limited to, activating chondrocytes. Binding of a ligand to a receptor is known to alter intracellular levels of small molecules, such as calcium, potassium and sodium, as well as alter pH

and membrane potential. Alterations in small molecule concentration can be measured to identify supernatants which bind to receptors of a particular cell. In specific embodiments, polypeptides of the invention comprise the following amino acid sequence:

- 5 KSYFRTMGGTKRGIKKLVNVCLKHPKNTSLSQQLVFAKINKILISKTTK
STNLKGLKCLPPLSVSIHPTFIYYKHNTTLRIVFGTYFDFFPYRKNKDQAFEGE
DWESSLNVS DAW (SEQ ID NO:323), TKRGIKKLVNVCLKHPKNTSLS (SEQ ID
NO:324), and/or SIHPTFIYYKHNTTLRIVFGTYFDFF (SEQ ID NO:325).

Polynucleotides encoding these polypeptides are also encompassed by the invention.

- 10 The gene encoding the disclosed cDNA is believed to reside on chromosome 3.
Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 3.

This gene is expressed primarily in resting T-cells, and to a lesser extent, in retina and placenta.

- 15 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune, reproductive, or eye disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes
20 for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g. Immune, hematopoietic, eye, reproductive, and cancerous and wounded tissues) or bodily fluids (e.g.lymph, serum, amniotic fluid, plasma, urine,
25 synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

- Preferred epitopes include those comprising a sequence shown in SEQ ID
30 NO:173 as residues: Met-1 to Pro-12.

- The tissue distribution of this gene predominantly in T-cells and placenta, combined with the detected calcium flux activity indicates that the gene could be important for the treatment or detection of immune or hematopoietic disorders including arthritis, asthma, immunodeficiency diseases and leukemia. Expression of the gene at
35 high levels in the retina indicates a role in the treatment and/or detection of eye disorders including color blindness, blindness, vision defects, and light sensitivity. Protein, as

well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:48 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1024 of SEQ ID NO:48, b is an integer of 15 to 1038, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:48, and where b is greater than or equal to a + 14.

15 FEATURES OF PROTEIN ENCODED BY GENE NO: 39

This gene is expressed primarily in brain.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, developmental, degenerative and behavioral diseases of the brain such as depression, schizophrenia, Alzheimer's disease, Parkinson's disease, Huntington's disease, specific brain tumors, aphasia, mania, depression, dementia, paranoia, addictive behavior and sleep disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the brain, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g., brain, cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:174 as residues: Pro-35 to Met-42.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of developmental,

degenerative and behavioral diseases and conditions of the brain such as aphasia, depression, schizophrenia, Alzheimer's disease, Parkinson's disease, Huntington's disease, specific brain tumors, mania, depression, dementia, paranoia, addictive behavior and sleep disorders.

- 5 Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:49 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome.
- 10 Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of $a-b$, where a is any integer between 1 to 1162 of SEQ ID NO:49, b is an integer of 15 to 1176, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:49, and where b is greater than or equal to $a + 14$.

15

FEATURES OF PROTEIN ENCODED BY GENE NO: 40

- The gene encoding the disclosed cDNA is thought to reside on chromosome 17.
- 20 Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 17.

This gene is expressed primarily in synovium.

- Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, disorders of the muscular-skeletal system. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the muscular skeletal system, expression of
- 25 this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. synovium, cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an
- 30 individual not having the disorder.
- 35

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:175 as residues: Pro-15 to Cys-29, Gly-40 to Tyr-54, Pro-72 to His-79.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of disorders of the muscular skeletal system. Furthermore, the expression of this gene product in synovium would suggest a role in the detection and treatment of disorders and conditions affecting the skeletal system, in particular osteoporosis, as well as disorders afflicting connective tissues (e.g. arthritis, trauma, tendonitis, chondromalacia and inflammation), such as in the diagnosis or treatment of various autoimmune disorders such as rheumatoid arthritis, lupus, scleroderma, and dermatomyositis as well as dwarfism, spinal deformation, and specific joint abnormalities as well as chondrodysplasias (ie. spondyloepiphyseal dysplasia congenita, familial osteoarthritis, Atelosteogenesis type II, metaphyseal chondrodysplasia type Schmid). Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:50 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 717 of SEQ ID NO:50, b is an integer of 15 to 731, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:50, and where b is greater than or equal to a + 14.

25

FEATURES OF PROTEIN ENCODED BY GENE NO: 41

The translation product of this gene shares sequence homology with Enoyl-CoA hydratase, which is an RNA binding protein with intrinsic enzymatic activity thought to be important in metabolic disorders. The gene encoding the disclosed cDNA is thought to reside on chromosome 1. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 1.

This gene is expressed primarily in fetal liver.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, metabolic disorders, liver disorders and cancer. Similarly, polypeptides

and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the hepatic and metabolic systems, expression of this gene at significantly higher or lower levels may be routinely
5 detected in certain tissues or cell types (e.g. liver, cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

10 Preferred epitopes include those comprising a sequence shown in SEQ ID NO:176 as residues: Pro-10 to Arg-15, Leu-96 to Ser-103, Gly-172 to Pro-178, Gln-213 to Asp-218, Asn-268 to Leu-275, Arg-282 to Phe-289.

The tissue distribution and homology to Enoyl-CoA hydratase indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment
15 and diagnosis of metabolic and liver diseases and cancer. Furthermore, the tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the detection and treatment of liver disorders and cancers (e.g. hepatoblastoma, jaundice, hepatitis, liver metabolic diseases and conditions that are attributable to the differentiation of hepatocyte progenitor cells). Protein, as well as,
20 antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:51 and may have been publicly available prior to conception of the present
25 invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1423 of SEQ ID NO:51, b is an integer of 15
30 to 1437, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:51, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 42

35

This gene is expressed primarily in rhabdomyosarcoma tissue.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, disorders of the muscular skeletal system and cancer. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the muscular skeletal system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. musculo-skeletal, cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for treatment and diagnosis of disorders of the muscular skeletal system and cancer. Furthermore, the tissue distribution indicates a role in the detection and treatment of disorders and conditions affecting the musculo-skeletal system, in particular rhabdomyosarcomas as well as related cancers. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:52 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1355 of SEQ ID NO:52, b is an integer of 15 to 1369, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:52, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 43

This gene is expressed primarily in neutrophils.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a

biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune or hematopoietic disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in neutrophils indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of aberrant immune responses to foreign antigens. Furthermore, expression of this gene product in neutrophils indicates a role in the regulation of the proliferation; survival; differentiation; and/or activation of potentially all hematopoietic cell lineages, including blood stem cells. This gene product may be involved in the regulation of cytokine production, antigen presentation, or other processes that may also suggest a usefulness in the treatment of cancer (e.g. by boosting immune responses). Since the gene is expressed in cells of lymphoid origin, the gene or protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues. Therefore it may be also used as an agent for immunological disorders including arthritis, asthma, immune deficiency diseases such as AIDS, leukemia, rheumatoid arthritis, inflammatory bowel disease, sepsis, acne, and psoriasis. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:53 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome.

Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1023 of SEQ ID NO:53, b is an integer of 15

to 1037, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:53, and where b is greater than or equal to a + 14.

5 FEATURES OF PROTEIN ENCODED BY GENE NO: 44

This gene is expressed primarily in neutrophils induced with IL-1 and LPS.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune or hematopoietic disorders, particularly in aberrant neutrophil responses to infection. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:179 as residues: Lys-36 to Cys-42.

The tissue distribution in neutrophils indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of a lack of immune response to infection. Furthermore, expression of this gene product in neutrophils indicates a role in the regulation of the proliferation; survival; differentiation; and/or activation of potentially all hematopoietic cell lineages, including blood stem cells. This gene product may be involved in the regulation of cytokine production, antigen presentation, or other processes that may also suggest a usefulness in the treatment of cancer (e.g. by boosting immune responses). Since the gene is expressed in cells of lymphoid origin, the gene or protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues. Therefore it may be also used as an agent for immunological disorders including arthritis, asthma, immune deficiency diseases such as AIDS, leukemia, rheumatoid arthritis, inflammatory bowel disease, sepsis, acne, and psoriasis. In addition, this gene product may have commercial utility in the expansion

of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

- 5 Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:54 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome.
- 10 Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1359 of SEQ ID NO:54, b is an integer of 15 to 1373, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:54, and where b is greater than or equal to a + 14.

15

FEATURES OF PROTEIN ENCODED BY GENE NO: 45

This gene is expressed primarily in brain.

- 20 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, CNS disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification
- 25 of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the central nervous system (CNS), expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. brain, cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken
- 30 from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

- The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of disorders of the
- 35 central nervous system. Furthermore, elevated expression of this gene product within the frontal cortex of the brain indicates that it may be involved in neuronal survival; synapse formation; conductance; neural differentiation, etc. Such involvement may

impact many processes, such as learning and cognition. It may also be useful in the treatment of such neurodegenerative disorders as schizophrenia; ALS; or Alzheimer's.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:55 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1333 of SEQ ID NO:55, b is an integer of 15 to 1347, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:55, and where b is greater than or equal to a + 14.

15 FEATURES OF PROTEIN ENCODED BY GENE NO: 46

This gene is expressed primarily in spleen.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune or hematopoietic disorders, particularly those affecting the spleen, such as in T- and B-cell maturation and their resulting efficacy in the immune response. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. immune, hematopoietic, spleen, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:181 as residues: Ser-20 to Ser-34, Thr-40 to Ser-46.

The tissue distribution in spleen indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of disorders affecting the spleen and immune system. Furthermore, this gene may play a role in the

survival, proliferation, and/or differentiation of hematopoietic cells in general, and may be of use in the augmentation of the numbers of stem cells and committed progenitors. This gene product may be involved in the regulation of cytokine production, antigen presentation, or other processes that may also suggest a usefulness in the treatment of cancer (e.g. by boosting immune responses). Since the gene is expressed in cells of lymphoid origin, the natural gene product may be involved in immune functions. Therefore it may be also used as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's disease, scleroderma and tissues. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:56 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 808 of SEQ ID NO:56, b is an integer of 15 to 822, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:56, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 47

In specific embodiments, polypeptides of the invention comprise the following amino acid sequence:

QRPHPQPWXPMTLMGTGIPVFAHKMLPFDPPCHLSCTHINPKPXXPQGDEQK
SQGTEEWCDREGKKRSI (SEQ ID NO:326), PMTLMGTGIPVFAHKMLPFDP

(SEQ ID NO:327), PPCHLSCTHINPKPXXPQGDE (SEQ ID NO:328), EQKSQGT
EEWCDREGKKRRSI (SEQ ID NO:329), DEWGAGRRMEWEDNLPLEFSCPVT
KLLSVPSWTPLDAQMLLLFFPSLSHHSSVPWLFCSGPCGXXGLGFI (SEQ ID
NO:330), EWEDNLPLEFSCPVTKLLSVP (SEQ ID NO:331), PSWTPLDAQM
5 LLLFFPSLSHH (SEQ ID NO:332), and/or HSSVPWLFCSGPCGXXGLGFI (SEQ
ID NO:333). Polynucleotides encoding these polypeptides are also encompassed by the
invention.

This gene is expressed primarily in neutrophils.

Therefore, polynucleotides and polypeptides of the invention are useful as
10 reagents for differential identification of the tissue(s) or cell type(s) present in a
biological sample and for diagnosis of diseases and conditions which include, but are
not limited to, diseases of the immune system, including neutropenia, cancer,
inflammatory diseases and allergies. Similarly, polypeptides and antibodies directed to
these polypeptides are useful in providing immunological probes for differential
15 identification of the tissue(s) or cell type(s). For a number of disorders of the above
tissues or cells, particularly of the immune system, expression of this gene at
significantly higher or lower levels may be routinely detected in certain tissues and cell
types (e.g. Immune, hematopoietic, and cancerous and wounded tissues) or bodily
fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another
20 tissue or cell sample taken from an individual having such a disorder, relative to the
standard gene expression level, i.e., the expression level in healthy tissue or bodily
fluid from an individual not having the disorder.

The tissue distribution in neutrophils indicates that polynucleotides and
polypeptides corresponding to this gene are useful for treatment/diagnosis of diseases
25 of the immune system since expression is primarily in neutrophils, and may be useful
as a growth factor for the differentiation or proliferation of neutrophils for the treatment
of neutropenia following chemotherapy or may be useful in the treatment of immune
dysfunction or anti-inflammatory by inhibiting infiltration of neutrophils to the site of
injury or distress. Protein, as well as, antibodies directed against the protein may show
30 utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available
and accessible through sequence databases. Some of these sequences are related to SEQ
ID NO:57 and may have been publicly available prior to conception of the present
invention. Preferably, such related polynucleotides are specifically excluded from the
35 scope of the present invention. To list every related sequence is cumbersome.
Accordingly, preferably excluded from the present invention are one or more
polynucleotides comprising a nucleotide sequence described by the general formula of

a-b, where a is any integer between 1 to 522 of SEQ ID NO:57, b is an integer of 15 to 536, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:57, and where b is greater than or equal to a + 14.

5

FEATURES OF PROTEIN ENCODED BY GENE NO: 48

This gene is expressed primarily in prostate, brain and T-cells.

Therefore, polynucleotides and polypeptides of the invention are useful as
10 reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, disorders of the reproductive, CNS and immune system. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a
15 number of disorders of the above tissues or cells, particularly of the reproductive, CNS and immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. immune, brain, prostate, cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual
20 having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:183 as residues: Asp-26 to Gly-32, Ile-37 to Trp-44.

25

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of disorders of the reproductive, CNS and immune systems. Furthermore, the tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the detection/treatment of neurodegenerative disease states and behavioural disorders such
30 as Alzheimers Disease, Parkinsons Disease, Huntingtons Disease, Tourette Syndrome, mania, dementia, paranoia, obsessive compulsive disorder, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and perception. Additionally, the tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for
35 the diagnosis and/or treatment of hematopoietic disorders. This gene product is primarily expressed in hematopoietic cells and tissues, suggesting that it plays a role in the survival, proliferation, and/or differentiation of hematopoietic lineages. Expression

of this gene product in T cells strongly indicates a role for this protein in immune function and immune surveillance. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

- 5 Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:58 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome.
- 10 Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1248 of SEQ ID NO:58, b is an integer of 15 to 1262, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:58, and where b is greater than or equal to a + 14.

15

FEATURES OF PROTEIN ENCODED BY GENE NO: 49

This gene is expressed primarily in frontal cortex of schizophrenics.

- 20 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, CNS diseases and Schizophrenia. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for
- 25 differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the central nervous system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. brain, cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample
- 30 taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

- The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of disorders of the
- 35 CNS and schizophrenia. Furthermore, the tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and/or treatment of disorders of the brain and nervous system. Elevated

expression of this gene product within the frontal cortex of the brain indicates that it may be involved in neuronal survival; synapse formation; conductance; neural differentiation, etc. Such involvement may impact many processes, such as learning and cognition. It may also be useful in the treatment of such neurodegenerative disorders as schizophrenia; ALS; or Alzheimer's.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:59 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1255 of SEQ ID NO:59, b is an integer of 15 to 1269, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:59, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 50

This gene is expressed primarily in the testes.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, reproductive or endocrine disorders, particularly for male infertility and testicular cancer. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the male reproductive system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. reproductive, testicular, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, seminal fluid, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:185 as residues: His-62 to Ser-74, Leu-99 to Gln-104.

The tissue distribution in testes indicates that polynucleotides and polypeptides corresponding to this gene are useful for treating male infertility. The protein product is likely involved in sperm development and could be administered by injection or related techniques. Alternatively, this gene could be transfected in gene-replacement treatments into the cells of the testes and the protein products could be produced. The presence of expression of this gene at either the RNA or protein level could be used as a diagnostic in testicular cancer. Furthermore, the tissue distribution indicates that the protein product of this gene is useful for the treatment and diagnosis of conditions concerning proper testicular function (e.g. endocrine function, sperm maturation), as well as cancer. Therefore, this gene product is useful in the treatment of male infertility and/or impotence. This gene product is also useful in assays designed to identify binding agents as such agents (antagonists) are useful as male contraceptive agents. Similarly, the protein is believed to be useful in the treatment and/or diagnosis of testicular cancer. The testes are also a site of active gene expression of transcripts that may be expressed, particularly at low levels, in other tissues of the body. Therefore, this gene product may be expressed in other specific tissues or organs where it may play related functional roles in other processes, such as hematopoiesis, inflammation, bone formation, and kidney function, to name a few possible target indications.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:60 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1815 of SEQ ID NO:60, b is an integer of 15 to 1829, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:60, and where b is greater than or equal to a + 14.

30

FEATURES OF PROTEIN ENCODED BY GENE NO: 51

In specific embodiments, polypeptides of the invention comprise the following amino acid sequence: QGLSHIFWMNEQTLK (SEQ ID NO:334). Polynucleotides encoding these polypeptides are also encompassed by the invention.

This gene is expressed primarily in activated T-cells.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune disorders, particularly acute inflammatory conditions or autoimmune disease. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g. Immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in activated T-cells indicates that polynucleotides and polypeptides corresponding to this gene are useful for modulating the response of activated T-cells to treat inflammation or autoimmune diseases. The expression of this gene product indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. This gene product may be involved in the regulation of cytokine production, antigen presentation, or other processes that may also suggest a usefulness in the treatment of cancer (e.g. by boosting immune responses). Since the gene is expressed in cells of lymphoid origin, the natural gene product may be involved in immune functions. Therefore it may be also used as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's disease, scleroderma and tissues. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ

ID NO:61 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome.

Accordingly, preferably excluded from the present invention are one or more

- 5 polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1098 of SEQ ID NO:61, b is an integer of 15 to 1112, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:61, and where b is greater than or equal to a + 14.

10

FEATURES OF PROTEIN ENCODED BY GENE NO: 52

When tested against U937 cell lines, supernatants removed from cells containing this gene activated the GAS (gamma activating sequence) promoter element.

- 15 Thus, it is likely that this gene activates myeloid cells, including their progenitors, through the JAK-STAT signal transduction pathway. GAS is a promoter element found upstream of many genes which are involved in the Jak-STAT pathway. The Jak-STAT pathway is a large, signal transduction pathway involved in the differentiation and proliferation of cells. Therefore, activation of the Jak-STAT pathway, reflected by the
- 20 binding of the GAS element, can be used to indicate proteins involved in the proliferation and differentiation of cells. In specific embodiments, polypeptides of the invention comprise the following amino acid sequence:

TLVCLGVSSEEGSCPRDVTGPGCCFSLTLTGF (SEQ ID NO:335), ADLIVLWH
HHPLWPQHLALPSSGASHDH VELTVYPKTVAASWLLELSRPPIFCLFTXPALT
25 XHGLDRVAALVECTIWXXXGMWYRRRYSQCFRDRSI RDVFPEAVMLQQH
LRHLAVATYRCRRRSPCKAPTVEEAEGGKPRAVPSGTGFQKHGQEPGGSTSP
HWFVG HLQLLVLSVNNRQLFVQGRAGYLEMTGLPCPKLLLTLLRGLT
PGVGHGLCAYRRGCLAWRLDXAS (SEQ ID NO:336), ILWRQAPEAPHCSQDSV
SSSPRLQEDLAHVTQVTRHPHFRSLPSAWCSHSSLLPVSLPRHALATKSPNMX
30 XSSPILHLIQFTGQISS PLGGXVQPPGQTASPICTQPM SHPRRQASQQCEQ
QLWTGQTSHLQIPCPALNKELPVVDTQDKELQMSPE PMWGCGPSRLLPM
LLESCA (SEQ ID NO:337), MLQQHLRHLAVATYRCRRRSPCKAPTVEEAEGGK
(SEQ ID NO:338), VTQVTRHPHFRSLPSAWCSHSSLLPVSLP (SEQ ID NO:339),
and/or GQTASPICTQPM SHPRRQASQQCEQQLW (SEQ ID NO:340).

- 35 Polynucleotides encoding these polypeptides are also encompassed by the invention.

This gene is expressed primarily in activated T-cells.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune disorders, particularly autoimmune diseases and inflammation.

5 Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g. Immune, hematopoietic, and
10 cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

15 Preferred epitopes include those comprising a sequence shown in SEQ ID NO:187 as residues: Ser-25 to Lys-33.

The tissue distribution in neutrophils, combined with the detected GAS biological activity indicates that polynucleotides and polypeptides corresponding to this gene are useful for modulating the response of activated T-cells and other cells of the
20 immune system involved in inflammation and autoimmune diseases. Similarly, this gene product may be involved in the regulation of cytokine production, antigen presentation, or other processes that may also suggest a usefulness in the treatment of cancer (e.g. by boosting immune responses). Since the gene is expressed in cells of lymphoid origin, the natural gene product may be involved in immune functions.
25 Therefore it may be also used as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-
30 graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's disease, scleroderma and tissues. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the
35 differentiation and/or proliferation of various cell types. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:62 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of $a-b$, where a is any integer between 1 to 1660 of SEQ ID NO:62, b is an integer of 15 to 1674, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:62, and where b is greater than or equal to $a + 14$.

FEATURES OF PROTEIN ENCODED BY GENE NO: 53

In specific embodiments, polypeptides of the invention comprise the following amino acid sequence:
 FITLRLGPKNMAGVLWRHSNLQTPHYISWCPLLNYRETGNCLLHVSG FLNSR
 LLANCSGEASGKVIQTLLWPGEISAVA (SEQ ID NO:341), KIRTFLLFSGHRLFST
 QGQSLTVKAHTAF MLIVKNLRYFIAFKFLMGISDSSEIGLVMQPLQKPHTV
 ILIRGIEFLSPGGVLP (SEQ ID NO:342), MAGVLWRHSNLQTPHYISWCPLLNYR
 (SEQ ID NO:343), and/or YFIAFKFLMGISDSSEIGLVMQPLQKPHT (SEQ ID
 NO:344). Polynucleotides encoding these polypeptides are also encompassed by the invention. The gene encoding the disclosed cDNA is believed to reside on chromosome 12. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 12.

This gene is expressed primarily in spleen, and to a lesser extent, in bone marrow and B-cells.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune and hematopoietic disorders, particularly multiple myeloma, immunodeficiencies, and infections. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune and hematopoietic disorders, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g. Immune, hematopoietic, and cancerous and wounded

tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

5 The tissue distribution of this gene predominantly in hematopoietic cell types and immune tissues indicates that the gene could be important for the treatment or detection of immune or hematopoietic disorders including arthritis, asthma, immunodeficiency diseases and leukemia. Moreover, this gene product may be involved in the regulation of cytokine production, antigen presentation, or other
10 processes that may also suggest a usefulness in the treatment of cancer (e.g. by boosting immune responses). Since the gene is expressed in cells of lymphoid origin, the natural gene product may be involved in immune functions. Therefore it may be also used as an agent for immunological disorders including arthritis, asthma,
15 immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia,
20 rheumatoid arthritis, Sjogren's disease, scleroderma and tissues. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

25 Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:63 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome.

30 Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of $a-b$, where a is any integer between 1 to 1031 of SEQ ID NO:63, b is an integer of 15 to 1045, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:63, and where b is greater than or equal to $a + 14$.

FEATURES OF PROTEIN ENCODED BY GENE NO: 54

The translation product of this gene shares very weak sequence homology with follicle-stimulating hormone beta subunit, which is thought to be important in hormonal regulation. When tested against K562 leukemia cell lines, supernatants removed from cells containing this gene activated the ISRE assay. Thus, it is likely that this gene activates leukemia cells through the Jak-STAT signal transduction pathway. The interferon-sensitive response element is a promoter element found upstream of many genes which are involved in the Jak-STAT pathway. The Jak-STAT pathway is a large, signal transduction pathway involved in the differentiation and proliferation of cells. Therefore, activation of the Jak-STAT pathway, reflected by the binding of the ISRE element, can be used to indicate proteins involved in the proliferation and differentiation of cells. The gene encoding the disclosed cDNA is thought to reside on chromosome 4. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 4.

This gene is expressed primarily in adult brain and adipocytes.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, endocrine diseases. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the endocrine system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. brain, cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in brain and homology to follicle stimulating hormone indicates that polynucleotides and polypeptides corresponding to this gene are useful as a hormone for the diagnosis and treatment of endocrine disorders. The brain is a major site for secreting various hormones that regulate a wide range of body physiology. The secretory molecule encoded by this gene has very weak homology with FSH, and further indicates that it may serve as an endocrine. Endocrines can often be used in hormonal treatment of pathological disorders or change of physiology under certain circumstances such as in the treatment of reproductive disorders.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:64 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1037 of SEQ ID NO:64, b is an integer of 15 to 1051, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:64, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 55

The translation product of this gene shares homology with a number of a C. elegans proteases, which are thought to be important in programmed cell death.

This gene is expressed primarily in activated T-cells and to a lesser extent in human stomach.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune disorders or stomach diseases. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. immune, cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:190 as residues: Lys-41 to Arg-47, Asp-125 to Lys-139, Ser-177 to Glu-185.

The tissue distribution in activated T-cells and stomach indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of immune disorders, transplantation or stomach disease.

Particularly, the expression of the gene by activated T-cells can be used for the development of therapeutic agents as immune suppressants or immune modulators.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:65 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1168 of SEQ ID NO:65, b is an integer of 15 to 1182, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:65, and where b is greater than or equal to a + 14.

15 FEATURES OF PROTEIN ENCODED BY GENE NO: 56

The translation product of this gene shares sequence homology with CD53 tetraspan transmembrane molecule which is thought to be important in leukocyte activation. The gene encoding the disclosed cDNA is thought to reside on chromosome 7. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 7.

This gene is expressed primarily in KMH2 and activated T-cells, and to a lesser extent in tonsils.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, infection, inflammation and other immune disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. immune, cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:191 as residues: Lys-99 to Arg-107.

The tissue distribution and homology to CD53 indicates that polynucleotides and polypeptides corresponding to this gene are useful for diagnosis and development of therapeutic agents for immune disorders including infection, allergy, inflammation, transplantation and immune deficiencies. Furthermore, expression of this gene product in tonsils indicates a role in the regulation of the proliferation; survival; differentiation; and/or activation of potentially all hematopoietic cell lineages, including blood stem cells. This gene product may be involved in the regulation of cytokine production, antigen presentation, or other processes that may also suggest a usefulness in the treatment of cancer (e.g. by boosting immune responses). Since the gene is expressed in cells of lymphoid origin, the gene or protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues. Therefore it may be also used as an agent for immunological disorders including arthritis, asthma, immune deficiency diseases such as AIDS, leukemia, rheumatoid arthritis, inflammatory bowel disease, sepsis, acne, and psoriasis. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Expression of this gene product in T cells strongly indicates a role for this protein in immune function and immune surveillance. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:66 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 661 of SEQ ID NO:66, b is an integer of 15 to 675, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:66, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 57

The gene encoding the disclosed cDNA is thought to reside on chromosome 17. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 17.

5 This gene is expressed primarily in fetal liver and to a lesser extent in neutrophils and keratinocytes.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, inflammation, autoimmune and skin defects. Similarly, polypeptides and
10 antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. liver, cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum,
15 plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID
20 NO:192 as residues: Pro-41 to Gln-50.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the study and treatment of inflammatory, general immune, and skin disorders. Furthermore, the tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the
25 diagnosis and/or treatment of hematopoietic disorders. This gene product is primarily expressed in hematopoietic cells and tissues, suggesting that it plays a role in the survival, proliferation, and/or differentiation of hematopoietic lineages. This is particularly supported by the expression of this gene product in fetal liver, which is a primary site of definitive hematopoiesis. Expression of this gene product in neutrophils
30 also strongly indicates a role for this protein in immune function and immune surveillance.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:67 and may have been publicly available prior to conception of the present
35 invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more

polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1091 of SEQ ID NO:67, b is an integer of 15 to 1105, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:67, and where b is greater than or equal to a + 14.

5

FEATURES OF PROTEIN ENCODED BY GENE NO: 58

This gene is expressed primarily in induced neutrophils.

10 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune and haemopoietic disorders. Similarly, polypeptides and
15 antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the haemopoietic and immune systems, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. immune, cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another
20 tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of disorders of the
25 haemopoietic and immune systems. This gene product may be involved in the regulation of cytokine production, antigen presentation, or other processes that may also suggest a usefulness in the treatment of cancer (e.g. by boosting immune responses). Since the gene is expressed in cells of lymphoid origin, the gene or protein, as well as, antibodies directed against the protein may show utility as a tumor marker
30 and/or immunotherapy targets for the above listed tissues. Expression of this gene product in neutrophils also strongly indicates a role for this protein in immune function and immune surveillance.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ
35 ID NO:68 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome.

Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1265 of SEQ ID NO:68, b is an integer of 15 to 1279, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:68, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 59

10 This gene is expressed primarily in the endometrium.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of female infertility or reproductive disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in
15 providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the reproductive system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. reproductive, endometrium, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, amniotic
20 fluid, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in endometrium indicates that polynucleotides and
25 polypeptides corresponding to this gene are useful for treating female infertility. The protein product may show utility in the preparation of the endometrium of implantation and could be administered either topically or orally. Alternatively, this gene could be transfected in gene-replacement treatments into the cells of the endometrium and the protein products could be produced. Similarly, these treatments could be performed
30 during artificial insemination for the purpose of increasing the likelihood of implantation and development of a healthy embryo. In both cases this gene or its gene product could be administered at later stages of pregnancy to promote healthy development of the endometrium. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

35 Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:69 and may have been publicly available prior to conception of the present

invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome.

Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1624 of SEQ ID NO:69, b is an integer of 15 to 1638, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:69, and where b is greater than or equal to a + 14.

10 FEATURES OF PROTEIN ENCODED BY GENE NO: 60

This gene is expressed primarily in the cells of the immune system, such as eosinophils, T-cells, dendritic cells, and tonsils.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune or hematopoietic disorders, such as AIDS, inflammatory conditions, multiple myeloma, or SCID. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types or cell type (e.g. immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in various immune cells and tissues indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of immune system disorders, such as AIDS. Furthermore, expression of this gene product in tonsils and other immune cells indicates a role in the regulation of the proliferation; survival; differentiation; and/or activation of potentially all hematopoietic cell lineages, including blood stem cells. This gene product may be involved in the regulation of cytokine production, antigen presentation, or other processes that may also suggest a usefulness in the treatment of cancer (e.g. by boosting immune responses). Since the gene is expressed in cells of lymphoid origin, the gene or protein, as well as, antibodies directed against the protein may show utility

as a tumor marker and/or immunotherapy targets for the above listed tissues. Therefore it may be also used as an agent for immunological disorders including arthritis, asthma, immune deficiency diseases such as AIDS, leukemia, rheumatoid arthritis, inflammatory bowel disease, sepsis, acne, and psoriasis. In addition, this gene product
5 may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available
10 and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:70 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more
15 polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 873 of SEQ ID NO:70, b is an integer of 15 to 887, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:70, and where b is greater than or equal to a + 14.

20

FEATURES OF PROTEIN ENCODED BY GENE NO: 61

The translation product of this gene shares homology with human stannin, which is thought to play a role in the toxic effects of organotins. Moreover, the protein
25 product of this gene may also show utility in the treatment, and/or prevention of a variety of defects in calcium regulation and metabolism.

This gene is expressed primarily in GM-CSF treated macrophages.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a
30 biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune disorders, particularly in the treatment or amelioration of aberrant immune response to tumor or foreign antigens, and in phagocytosis. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell
35 type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. immune, and cancerous and

wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

5 Preferred epitopes include those comprising a sequence shown in SEQ ID NO:196 as residues: Gly-43 to Gly-55.

The tissue distribution in macrophages indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of immune disorders. Furthermore, the tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and/or
10 treatment of hematopoietic disorders. This gene product is primarily expressed in hematopoietic cells and tissues, suggesting that it plays a role in the survival, proliferation, and/or differentiation of hematopoietic lineages. Expression of this gene product in macrophage also strongly indicates a role for this protein in immune function
15 and immune surveillance. The protein product may even serve to stimulate the immune response, or may be used to inhibit such a response which may be useful during host versus graft disease or autoimmune disorders. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

20 Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:71 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome.

25 Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 850 of SEQ ID NO:71, b is an integer of 15 to 864, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:71, and where b is greater than or equal to a + 14.

30

FEATURES OF PROTEIN ENCODED BY GENE NO: 62

This gene is expressed primarily in activated monocytes.

35 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are

not limited to, immune or hematopoietic disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene

5 at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily

10 fluid from an individual not having the disorder.

The tissue distribution in monocytes indicates that polynucleotides and polypeptides corresponding to this gene are useful for diagnosing and/or treating immune or hematopoietic disorders. This gene product is primarily expressed in hematopoietic cells and tissues, suggesting that it plays a role in the survival,

15 proliferation, and/or differentiation of hematopoietic lineages. Expression of this gene product in monocytes also strongly indicates a role for this protein in immune function and immune surveillance. Moreover, polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of hematopoietic related disorders such as anemia, pancytopenia, leukopenia, thrombocytopenia or leukemia since stromal

20 cells are important in the production of cells of hematopoietic lineages. The uses include bone marrow cell ex vivo culture, bone marrow transplantation, bone marrow reconstitution, radiotherapy or chemotherapy of neoplasia. The gene product may also be involved in lymphopoiesis, therefore, it can be used in immune disorders such as infection, inflammation, allergy, immunodeficiency etc. In addition, this gene product

25 may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available

30 and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:72 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more

35 polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1203 of SEQ ID NO:72, b is an integer of 15

to 1217, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:72, and where b is greater than or equal to a + 14.

5 FEATURES OF PROTEIN ENCODED BY GENE NO: 63

This gene is expressed primarily in activated monocytes.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune or hematopoietic disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:198 as residues: Met-1 to Gly-6.

The tissue distribution in monocytes indicates that polynucleotides and polypeptides corresponding to this gene are useful for diagnosing and/or treating immune or hematopoietic disorders. This gene product is primarily expressed in hematopoietic cells and tissues, suggesting that it plays a role in the survival, proliferation, and/or differentiation of hematopoietic lineages. Expression of this gene product in monocytes also strongly indicates a role for this protein in immune function and immune surveillance. Moreover, polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of hematopoietic related disorders such as anemia, pancytopenia, leukopenia, thrombocytopenia or leukemia since stromal cells are important in the production of cells of hematopoietic lineages. The uses include bone marrow cell ex vivo culture, bone marrow transplantation, bone marrow reconstitution, radiotherapy or chemotherapy of neoplasia. The gene product may also be involved in lymphopoiesis, therefore, it can be used in immune disorders such as infection, inflammation, allergy, immunodeficiency etc. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors

of various blood lineages, and in the differentiation and/or proliferation of various cell types. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available
5 and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:73 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more
10 polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1703 of SEQ ID NO:73, b is an integer of 15 to 1717, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:73, and where b is greater than or equal to a + 14.

15

FEATURES OF PROTEIN ENCODED BY GENE NO: 64

When tested against K562 leukemia cell lines, supernatants removed from cells containing this gene activated the ISRE assay. Thus, it is likely that this gene activates
20 leukemia cells through the Jak-STAT signal transduction pathway. The interferon-sensitive response element is a promoter element found upstream of many genes which are involved in the Jak-STAT pathway. The Jak-STAT pathway is a large, signal transduction pathway involved in the differentiation and proliferation of cells. Therefore, activation of the Jak-STAT pathway, reflected by the binding of the ISRE
25 element, can be used to indicate proteins involved in the proliferation and differentiation of cells. The gene encoding the disclosed cDNA is thought to reside on chromosome 3. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 3.

This gene is expressed primarily in spleen from a chronic lymphocytic leukemia
30 patient.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune or hematopoietic disorders, particularly leukemias. Similarly,
35 polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system,

expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. spleen, immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in leukemia cells combined with the detected ISRE biological activity in K562 cell lines indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and/or treatment of chronic lymphocytic leukemia. Furthermore, since the gene is expressed in cells of lymphoid origin, the gene or protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues. Therefore it may be also used as an agent for immunological disorders including arthritis, asthma, immune deficiency diseases such as AIDS, rheumatoid arthritis, inflammatory bowel disease, sepsis, acne, and psoriasis. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Therefore it may be also used as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's disease, scleroderma and tissues. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:74 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of

a-b, where a is any integer between 1 to 1262 of SEQ ID NO:74, b is an integer of 15 to 1276, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:74, and where b is greater than or equal to a + 14.

5

FEATURES OF PROTEIN ENCODED BY GENE NO: 65

This gene is expressed primarily in neutrophils.

Therefore, polynucleotides and polypeptides of the invention are useful as
10 reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune or hematopoietic disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders
15 of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the
20 standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in neutrophils indicates that polynucleotides and polypeptides corresponding to this gene are useful for diagnosis and treatment of neutrophils inactivation and other immune system disorders. Furthermore,
25 polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and/or treatment of hematopoietic disorders. This gene product is primarily expressed in hematopoietic cells and tissues, suggesting that it plays a role in the survival, proliferation, and/or differentiation of hematopoietic lineages. Expression of this gene product in neutrophils also strongly indicates a role for this protein in immune
30 function and immune surveillance. Therefore it may be also used as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such
35 as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's disease,

scleroderma and tissues. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or

5 immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:75 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the

10 scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1130 of SEQ ID NO:75, b is an integer of 15 to 1144, where both a and b correspond to the positions of nucleotide residues shown

15 in SEQ ID NO:75, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 66

20 This gene is expressed primarily in neutrophils.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune or hematopoietic disorders, particularly neutropenia. Similarly,

25 polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. immune, hematopoietic, and cancerous and

30 wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in neutrophils indicates that polynucleotides and

35 polypeptides corresponding to this gene are useful for diagnosis and treatment of immune system disorders. Furthermore, expression of this gene product in neutrophils also strongly indicates a role for this protein in immune function and immune

surveillance. The protein may also be useful in the inhibition of neutrophil activation which may show utility in host-versus-graft disease and autoimmune disorders. Therefore it may be also used as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid
5 arthritis, granulomatous disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug
10 induced hemolytic anemia, rheumatoid arthritis, Sjogren's disease, scleroderma and tissues. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy
15 targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:76 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the
20 scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 904 of SEQ ID NO:76, b is an integer of 15 to 918, where both a and b correspond to the positions of nucleotide residues shown in
25 SEQ ID NO:76, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 67

30 When tested against U937 myeloid cell lines, supernatants removed from cells containing this gene activated the GAS promoter element. Thus, it is likely that this gene activates myeloid cells, and their progenitors, through the Jak-STAT signal transduction pathway. The gamma activating sequence (GAS) is a promoter element found upstream of many genes which are involved in the Jak-STAT pathway. The Jak-
35 STAT pathway is a large, signal transduction pathway involved in the differentiation and proliferation of cells. Therefore, activation of the Jak-STAT pathway, reflected by

the binding of the GAS element, can be used to indicate proteins involved in the proliferation and differentiation of cells.

This gene is expressed primarily in neutrophils.

Therefore, polynucleotides and polypeptides of the invention are useful as
5 reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune or hematopoietic disorders, such as neutropenia. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a
10 number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a
15 disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:202 as residues: Asp-23 to Trp-29.

The tissue distribution in neutrophils, combined with the detected GAS
20 biological activity in myeloid cell lines indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of immune system disorders. Furthermore, the tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and/or treatment of hematopoietic disorders. This gene product is primarily expressed in hematopoietic cells
25 and tissues, suggesting that it plays a role in the survival, proliferation, and/or differentiation of hematopoietic lineages. Expression of this gene product in neutrophils also strongly indicates a role for this protein in immune function and immune surveillance. The protein product of this gene may show utility in the inhibition of neutrophil activation which may show utility in host-versus-graft disease and in
30 autoimmune disorders. Therefore it may be also used as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as
35 host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lens tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's disease,

scleroderma and tissues. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:77 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1051 of SEQ ID NO:77, b is an integer of 15 to 1065, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:77, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 68

This gene is expressed primarily in neutrophils induced with IL-1 and LPS.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune or hematopoietic disorders, such as neutropenia. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in neutrophils indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of inactive immune response to foreign antigens. Furthermore, the tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for

the diagnosis and/or treatment of hematopoietic disorders. This gene product is primarily expressed in hematopoietic cells and tissues, suggesting that it plays a role in the survival, proliferation, and/or differentiation of hematopoietic lineages. Expression of this gene product in neutrophils also strongly indicates a role for this protein in

5 immune function and immune surveillance. Therefore it may be also used as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and

10 tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's disease, scleroderma and tissues. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood

15 lineages, and in the differentiation and/or proliferation of various cell types. The protein product of this gene may also show utility in the inactivation of neutrophils which may show utility in host-versus-graft disease or in autoimmune disorders, for example. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

20 Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:78 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome.

25 Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1112 of SEQ ID NO:78, b is an integer of 15 to 1126, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:78, and where b is greater than or equal to a + 14.

30

FEATURES OF PROTEIN ENCODED BY GENE NO: 69

The translation product of this nucleotide sequence shares homology with a

35 number of cysteine proteinases. Contact of cells with supernatant expressing the product of this gene increases the permeability of TF-1 Myeloid cells to calcium. Thus, it is likely that the product of this gene is involved in a signal transduction pathway that

is initiated when the product of this gene binds a receptor on the surface of the myeloid cell. Thus, polynucleotides and polypeptides have uses which include, but are not limited to, activating myeloid cells.

This gene is expressed primarily in tissue from an ovarian tumor.

5 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, reproductive disorders, particularly ovarian cancer. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing
10 immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the reproductive system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. reproductive, immune, hematopoietic, ovarian, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum,
15 amniotic fluid, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

 The homology to proteins of the cysteine proteinase family, tissue distribution
20 in ovarian tissues, combined with the detected calcium flux activity in myeloid cells indicates that the protein product of this gene may show utility in the treatment, and/or prevention of a variety of reproductive disorders, such as in ovarian cancer, or even in the modulation of the immune response to. Thus, it is useful for diagnosis and treatment of ovarian cancer. Furthermore, the biological activity data, when compared
25 to the tissue distribution, suggest that the translation product of this gene could be useful in activating the immune system to respond to cancerous growths, particularly those involving ovarian cancer. Protein, as well as, antibodies directed against the protein may show utility as a tissue-specific marker and/or immunotherapy target for the above listed tissues.

30 Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:79 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome.
35 Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 970 of SEQ ID NO:79, b is an integer of 15 to

984, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:79, and where b is greater than or equal to a + 14.

5 FEATURES OF PROTEIN ENCODED BY GENE NO: 70

This gene is expressed primarily in anergic T-cells.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune or hematopoietic disorders, such as autoimmune disorders including lupus. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:205 as residues: Ser-26 to Lys-34.

The tissue distribution in T-cells indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of a variety of immune system disorders. Expression of this gene product in T-cells indicates a role in the regulation of the proliferation; survival; differentiation; and/or activation of potentially all hematopoietic cell lineages, including blood stem cells. This gene product may be involved in the regulation of cytokine production, antigen presentation, or other processes that may also suggest a usefulness in the treatment of cancer (e.g. by boosting immune responses). Expression of this gene product in T cells also strongly indicates a role for this protein in immune function and immune surveillance. Therefore it may be also used as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-

host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's disease, scleroderma and tissues. In addition, this gene product may have commercial utility in the expansion of stem cells and committed
5 progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ
10 ID NO:80 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of
15 a-b, where a is any integer between 1 to 1233 of SEQ ID NO:80, b is an integer of 15 to 1247, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:80, and where b is greater than or equal to a + 14.

20 FEATURES OF PROTEIN ENCODED BY GENE NO: 71

This gene shares homology with the human adult heart neutral calponin, which is implicated in the regulation and modulation of smooth muscle contraction. It is capable of binding to actin, calmodulin, troponin C, and tropomyosin. The interaction
25 of calponin with actin inhibits the actomyosin Mg-ATPase activity. Therefore, the protein product of this gene may be beneficial as a vasoconstrictor or vasodilator, a muscle relaxor, treatment for tetanus stimuli, or for the treatment of various cardiovascular disorders. The gene encoding the disclosed cDNA is thought to reside on chromosome 19. Accordingly, polynucleotides related to this invention are useful as
30 a marker in linkage analysis for chromosome 19.

This gene is expressed primarily in adrenal gland tumor and human 12 week embryo. Furthermore, the gene is expressed in cardiomyopathy tissue.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a
35 biological sample and for diagnosis of diseases and disorders: endocrine, developmental, cardiovascular disorders, particularly diseases involving abnormal cellular proliferation such as cancers particularly of the adrenal gland, and disorders

involving heart muscle, such as cardiomyopathy. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the adrenal gland, heart, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. heart, muscle, endocrine, developmental, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of abnormal cellular proliferation, such as tumors. Alternatively, given the tissue distribution and the homology to human adult heart neutral calponin, it indicates that the translation product of this gene is useful for detecting, identifying, and/or treating disorders involving the degeneration of the regulation and modulation of smooth muscle contraction, such as is seen with cardiomyopathies. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:81 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 932 of SEQ ID NO:81, b is an integer of 15 to 946, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:81, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 72

This gene is expressed primarily in human bone and 9 week embryo.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are

not limited to, skeletal, immune, hemopoietic, or developmental disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the hematopoietic tissues, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. immune, bone, cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, amniotic fluid, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:207 as residues: Ala-22 to Lys-36.

The tissue distribution in bone and embryonic tissues indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis or the treatment of hemopoietic diseases. Furthermore, it may be useful in influencing bone mass in such conditions as osteoporosis. More generally, this gene may play a role in the survival, proliferation, and/or differentiation of hematopoietic cells in general, and may be of use in augmentation of the numbers of stem cells and committed progenitors.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:82 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1378 of SEQ ID NO:82, b is an integer of 15 to 1392, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:82, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 73

This gene is expressed primarily in T cells.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are

not limited to, disorder of the immune or hematopoietic systems, particularly immunodeficiencies or inflammatory conditions, such as AIDS, SCID, leukemias, or multiple myeloma. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:208 as residues: Asp-26 to Leu-36, Leu-42 to Phe-50.

The tissue distribution in T-cells indicates that polynucleotides and polypeptides corresponding to this gene are useful for treatment of disorders of the immune system such as AIDS. Furthermore, this gene product may be involved in the regulation of cytokine production, antigen presentation, or other processes that may also suggest a usefulness in the treatment of cancer (e.g. by boosting immune responses). Since the gene is expressed in cells of lymphoid origin, the gene or protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues. Therefore it may be also used as an agent for immunological disorders including arthritis, asthma, immune deficiency diseases such as AIDS, leukemia, rheumatoid arthritis, inflammatory bowel disease, sepsis, acne, and psoriasis. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Expression of this gene product in T cells also strongly indicates a role for this protein in immune function and immune surveillance. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:83 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of

a-b, where a is any integer between 1 to 1141 of SEQ ID NO:83, b is an integer of 15 to 1155, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:83, and where b is greater than or equal to a + 14.

5

FEATURES OF PROTEIN ENCODED BY GENE NO: 74

When tested against U937 cell lines, supernatants removed from cells containing this gene activated the GAS (gamma activating sequence) promoter element. Thus, it is likely that this gene activates myeloid cells, including progenitors, through the JAK-STAT signal transduction pathway. GAS is a promoter element found upstream of many genes which are involved in the Jak-STAT pathway. The Jak-STAT pathway is a large, signal transduction pathway involved in the differentiation and proliferation of cells. Therefore, activation of the Jak-STAT pathway, reflected by the binding of the GAS element, can be used to indicate proteins involved in the proliferation and differentiation of cells. In specific embodiments, polypeptides of the invention comprise the following amino acid sequence:

DVLLPLLYLLVRKHINRAGIGNTFQGGANCI (SEQ ID NO:345), MCCCLCCT
 SWSGSTSTERVSGTRFREVPTASCSSAPAPSELGSSLSVAAAALLSLPPRARLA
 LPRLPRL PSQENLRNPKGPQGNFQAPGAFVLSSSVA (SEQ ID NO:346), CAAA
 SAVPPGPEAHQQSGYREHVSGRCLHHRPLHPRRPNSALLSLLLLLLFSASH
 QEPGWHSQGSRAF QARRISGIPRDPRGTSKHLELLSFLVLWHRCCLPGG RXF
 CESLXQGRSACLLHQKPPLMLSAPLGEQLP TQLLLPPRSSGSKFXRYQRPGP
 RVGVHLHKGSSEIREAGGPQLWPQC PHPVDLDVLR TTQHCLQSEGPTS VH
 LSSV (SEQ ID NO:347), EVEEAELAAALPMEPRASIAGASGAADMHFCPAXGTH
 RXA YPQEGSTYATELERTKAPGAWKFPWGPLGFLRFSWLGRRGSLGSAS
 RALGGRLRRAAAATEREEPSSDGA GAEDHDAVGTS LKRVDPTRS VDVLPD
 QEVQQRQQHI (SEQ ID NO:348), RRISGIPRDPRGTSKHLELLSFLVLWHRCCL
 (SEQ ID NO:349), and/or RTKAPGAWKFPWGPLGFLRFSWLGRRGSL (SEQ ID
 NO:350). Polynucleotides encoding these polypeptides are also encompassed by the invention.

This gene is expressed primarily in smooth muscle.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, disorders of smooth muscle tissue, particularly vascular disorders, such as vasculitis, microvascular disease, atherosclerosis, stroke, aneurysm, and

embolism. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of smooth muscle tissue, expression of this gene at significantly higher or lower levels
 5 may be routinely detected in certain tissues and cell types (e.g. smooth muscle, vascular, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the
 10 disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:209 as residues: Ser-23 to Glu-54.

The tissue distribution in smooth muscle, combined with the detected GAS biological activity indicates that polynucleotides and polypeptides corresponding to this
 15 gene are useful for diagnosis and treatment of vascular or cardiopulmonary disorders. In addition, the protein may show utility in the modulation of the immune system in response to various vascular disorders, particularly in the early stages of atherosclerosis, embolism, thrombosis, and stroke. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy
 20 targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:84 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the
 25 scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1359 of SEQ ID NO:84, b is an integer of 15 to 1373, where both a and b correspond to the positions of nucleotide residues shown
 30 in SEQ ID NO:84, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 75.

35 In specific embodiments, polypeptides of the invention comprise the following amino acid sequence: PRLAQLRLLSL (SEQ ID NO:351),
 QSDFREMNQTNSTSNAAKAREAQQGRGRD REAIFSSSALEHLVCYLQAYKHT

LLFIRSLNEHGLQQLLFQWRDGLFGNWFYFRIPILLFFTGFHCYHLSC PHLPC
 AQRQSSRGTVPYVLCPPHHHLHHYSWFPFLIPVLHTLPKLQPKFHGRPEQPL
 NLLQVKPTSGTI ASAEQVWVK (SEQ ID NO:352). VCYLQAYKHTLLFIRSLNEH
 GLQQLLFQW (SEQ ID NO:353), and/or VPYVLCPPHHHLHHYSWFPFLIPVLH

5 TLPKL (SEQ ID NO:354). Polynucleotides encoding these polypeptides are also encompassed by the invention. The gene encoding the disclosed cDNA is believed to reside on chromosome 1. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 1.

10 This gene is expressed primarily in brain, ulcerative colitis, pancreas tumor, placenta, and to a lesser extent, in thyroid, bone marrow stromal cells, B-cell lymphoma, and hemangiopericytoma.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, tumors and degenerative conditions involving infiltration by the immune system, particularly in soft-tissues, in addition to, neural, gastrointestinal, metabolic, reproductive, endocrine, and hematopoietic, or immune disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g. neural, gastrointestinal, metabolic, reproductive, endocrine, hematopoietic, immune disorders, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, bile, amniotic fluid, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:210 as residues: Lys-33 to Arg-51, Gly-64 to Gly-74.

30 The tissue distribution in brain tissues indicates that polynucleotides and polypeptides corresponding to this gene are useful for treating the secondary effects of immune system involvement in diseases such as pancreatic tumors, ulcerative colitis, and Alzheimer's disease. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ

ID NO:85 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome.

Accordingly, preferably excluded from the present invention are one or more
5 polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1244 of SEQ ID NO:85, b is an integer of 15 to 1258, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:85, and where b is greater than or equal to a + 14.

10

FEATURES OF PROTEIN ENCODED BY GENE NO: 76

When tested against PC12 cell lines, supernatants removed from cells containing this gene activated the EGR1 (early growth response gene 1) promoter
15 element. Thus, it is likely that this gene activates sensory neuron cells through the EGR1 signal transduction pathway. EGR1 is a separate signal transduction pathway from Jak-STAT, genes containing the EGR1 promoter are induced in various tissues and cell types upon activation, leading the cells to undergo differentiation and proliferation. In specific embodiments, polypeptides of the invention comprise the
20 following amino acid sequence: ESERAVVYLITGALFIVSSCVLCFLPSSRRE (SEQ ID NO:355). Polynucleotides encoding these polypeptides are also encompassed by the invention. The gene encoding the disclosed cDNA is believed to reside on chromosome 12. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 12.

25 This gene is expressed primarily in activated T cells, tonsils, and activated monocytes.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are
30 not limited to, immune and inflammatory disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the activated T cells, tonsils and activated monocytes, expression of this gene at significantly higher or lower levels may be
35 routinely detected in certain tissues and cell types (e.g. Immune, hematopoietic, neural, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an

individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in T-cells and immune tissues or cell types, combined
5 with the detected EGR biological activity indicates that polynucleotides and polypeptides corresponding to this gene are useful for diagnosis and treatment of immune and inflammatory disorders. Moreover, this gene product may be involved in the regulation of cytokine production, antigen presentation, or other processes that may also suggest a usefulness in the treatment of cancer (e.g. by boosting immune
10 responses). Since the gene is expressed in cells of lymphoid origin, the natural gene product may be involved in immune functions. Therefore it may be also used as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis,
15 hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's disease, scleroderma and tissues. In addition, this gene
20 product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available
25 and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:86 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more
30 polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1304 of SEQ ID NO:86, b is an integer of 15 to 1318, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:86, and where b is greater than or equal to a + 14.

35

FEATURES OF PROTEIN ENCODED BY GENE NO: 77

When tested against fibroblast cell lines, supernatants removed from cells containing this gene activated the EGR1 assay. Thus, it is likely that this gene activates fibroblast cells through a signal transduction pathway. Early growth response 1 (EGR1) is a promoter associated with certain genes that induces various tissues and cell types upon activation, leading the cells to undergo differentiation and proliferation. The gene encoding the disclosed cDNA is thought to reside on chromosome 16. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 16.

10 This gene is expressed primarily in eosinophils and activated T-cells and to a lesser extent in lung and thymus stromal cells.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. immune, cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

25 Preferred epitopes include those comprising a sequence shown in SEQ ID NO:212 as residues: Met-1 to Trp-10.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of immune disorders, including infection, allergy, inflammation, graft rejection and immunodeficiency. Furthermore, the tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and/or treatment of hematopoietic disorders. This gene product is primarily expressed in hematopoietic cells and tissues, suggesting that it plays a role in the survival, proliferation, and/or differentiation of hematopoietic lineages. Expression of this gene product in T cells and eosinophils also strongly indicates a role for this protein in immune function and immune surveillance.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:87 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 964 of SEQ ID NO:87, b is an integer of 15 to 978, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:87, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 78

One embodiment of this gene comprises polypeptides of the following amino acid sequence:
 MWVXGEEVLGSHAASPAFLHRCFSEESCFSIPEVEGYVVVLQPDAPQILLSGTA
 HFARPAVDFEGTNGVPLFPDLQITCSISHQVEAKKDESWQGTVTDTMSDEIVH
 NLDGCEISLVGDDLDPERESLLLDTTSLQQRGLELTNTSAYLTIAGVESITVYEEI
 LRQARYRLRHGAALYTRKFRLSCSEMNGRYSSENEFIVEVNVLHSMNRVAHPS
 HVLSXQQFLHRGHQPPPEMAGHSLASSHRNSST (SEQ ID NO:356), LGSHAA
 SPAFLHRCFSEESCFSI (SEQ ID NO:357), GYVVVLQPDAPQILLSGTAHFARP
 AVDFE (SEQ ID NO:358), ITCSISHQVEAKKDESWQGTVTDTMRM (SEQ ID
 NO:359), NLDGCEISLVGDDLDPERESLLLDTTSLQ (SEQ ID NO:360), SAYLTI
 AGVESITVYEEILRQAR (SEQ ID NO:361), RLSCSEMNGRYSSENEFIVEVNVLH
 SM (SEQ ID NO:362), and/or QQFLHRGHQPPPEMAGHSLASSHRN (SEQ ID
 NO:363). An additional embodiment is the polynucleotides encoding these polypeptides.

This gene is expressed primarily in brain and spleen.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, brain afflictions such as depression, schizophrenia, Alzheimer's disease, Parkinson's disease, Huntington's disease, specific brain tumors, aphasia, mania, depression, dementia, paranoia, addictive behavior and sleep disorders, as well as immune disorders such as leukemias, lymphomas, AIDS, arthritis and inflammation. Similarly, polypeptides and antibodies directed to these polypeptides are useful in

providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the central nervous and immune systems, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g., brain, cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

10 Preferred epitopes include those comprising a sequence shown in SEQ ID NO:213 as residues: Gly-36 to Leu-44.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of developmental, degenerative and behavioral diseases and conditions of the brain such as aphasia, depression, schizophrenia, Alzheimer's disease, Parkinson's disease, Huntington's disease, specific brain tumors, mania, depression, dementia, paranoia, addictive behavior and sleep disorders. In addition, the expression in spleen would suggest a possible role in the detection and treatment of immune disorders including: leukemias, lymphomas, auto-immunities, immunodeficiencies (e.g. AIDS), immuno-suppressive conditions (transplantation) and hematopoietic disorders as well as conditions of general microbial infection, inflammation or cancer.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:88 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1849 of SEQ ID NO:88, b is an integer of 15 to 1863, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:88, and where b is greater than or equal to a + 14.

35 FEATURES OF PROTEIN ENCODED BY GENE NO: 79

When tested against K562 leukemia cell lines, supernatants removed from cells containing this gene activated the ISRE assay. Thus, it is likely that this gene activates

leukemia cells through the Jak-STAT signal transduction pathway. The interferon-sensitive response element is a promoter element found upstream of many genes which are involved in the Jak-STAT pathway. The Jak-STAT pathway is a large, signal transduction pathway involved in the differentiation and proliferation of cells.

5 Therefore, activation of the Jak-STAT pathway, reflected by the binding of the ISRE element, can be used to indicate proteins involved in the proliferation and differentiation of cells. One embodiment of this gene comprises polypeptides of the following amino acid sequence: MADSETFISLE

ECRGHKRARKRTSMETALALEKLFPKQCQVLGIVTPGIVVXPMGSGSNRPQEI
 10 EIGESGFALLFPQIEGI KIQPFHFIDPKPNLTLEHQLTEVGLLDNPELRVVLV
 FGYNCKVGASNYLQQVVSTFSDMNIILAGGQV DNLSSLTSEKNPLDID AS
 GVVGLSFSGHRIQSATVLLNEDVSDEKTAEAAMQRLKAANIPEHNTIGFMFA
 CVGRGFQYYRAKGNVEADAFRKFPPSVPLFGFFGNGEIGCDRIVTGNFILRKC
 NEVKDDDLFHSYTTIMA LIHLGSSK (SEQ ID NO:364), HKRARKRTSMETAL
 15 ALEKLFP (SEQ ID NO:365), MGSGSNRPQEI EIGESGFALLFPQ (SEQ ID
 NO:366), FHFIDPKPNLTLEHQLTEVGL (SEQ ID NO:367), FGYNCKVGASN
 YLQQVVSTFSD (SEQ ID NO:368), TSEKNPLDIDASGVVGLSFS (SEQ ID
 NO:369), NEDVSDEKTAEAAMQRLKAANIPEHN (SEQ ID NO:370, YYRAKGNV
 EADAFRKFPPSVPLFGF (SEQ ID NO:371), and/or ICDRIVTGNFILRKCNE
 20 VKDDDLFH (SEQ ID NO:372). An additional embodiment is the polynucleotides
 encoding these polypeptides.

This gene is expressed primarily in endothelial cells and to a lesser extent in reproductive and various endocrine organs.

Therefore, polynucleotides and polypeptides of the invention are useful as
 25 reagents for differential identification of the tissue(s) or cell type(s) present in a
 biological sample and for diagnosis of diseases and conditions which include, but are
 not limited to, cancer, cardiovascular and immune defects. Similarly, polypeptides and
 antibodies directed to these polypeptides are useful in providing immunological probes
 for differential identification of the tissue(s) or cell type(s). For a number of disorders
 30 of the above tissues or cells, particularly of the immune, cardiovascular, and
 reproductive systems, expression of this gene at significantly higher or lower levels
 may be routinely detected in certain tissues or cell types (e.g. endothelial, reproductive,
 cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine,
 synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual
 35 having such a disorder, relative to the standard gene expression level, i.e., the
 expression level in healthy tissue or bodily fluid from an individual not having the
 disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:214 as residues: Ser-44 to Ala-50.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of cancer, cardiovascular and reproductive disorders.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:89 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2072 of SEQ ID NO:89, b is an integer of 15 to 2086, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:89, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 80

This gene is expressed primarily in human tongue and TNF-induced epithelium.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, mucosal, oral, and inflammatory conditions. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of mucosal and epidermal tissues, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. tongue, epithelial, cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:215 as residues: Ser-39 to Leu-48, Ala-65 to Pro-75, Pro-81 to Cys-87.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the study and treatment of disorders of the oral and intestinal mucosa, inflammation and other epithelial disorders.

Many polynucleotide sequences, such as EST sequences, are publicly available
5 and accessible through sequence databases. Some of these sequences are related to SEQ
ID NO:90 and may have been publicly available prior to conception of the present
invention. Preferably, such related polynucleotides are specifically excluded from the
scope of the present invention. To list every related sequence is cumbersome.
Accordingly, preferably excluded from the present invention are one or more
10 polynucleotides comprising a nucleotide sequence described by the general formula of
a-b, where a is any integer between 1 to 877 of SEQ ID NO:90, b is an integer of 15 to
891, where both a and b correspond to the positions of nucleotide residues shown in
SEQ ID NO:90, and where b is greater than or equal to a + 14.

15

FEATURES OF PROTEIN ENCODED BY GENE NO: 81

This gene is expressed primarily in activated neutrophils.

Therefore, polynucleotides and polypeptides of the invention are useful as
20 reagents for differential identification of the tissue(s) or cell type(s) present in a
biological sample and for diagnosis of diseases and conditions which include, but are
not limited to, immune, autoimmune, and inflammatory conditions. Similarly,
polypeptides and antibodies directed to these polypeptides are useful in providing
immunological probes for differential identification of the tissue(s) or cell type(s). For a
25 number of disorders of the above tissues or cells, particularly of the immune system,
expression of this gene at significantly higher or lower levels may be routinely detected
in certain tissues or cell types (e.g. immune, cancerous and wounded tissues) or bodily
fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another
tissue or cell sample taken from an individual having such a disorder, relative to the
30 standard gene expression level, i.e., the expression level in healthy tissue or bodily
fluid from an individual not having the disorder.

The tissue distribution indicates that polynucleotides and polypeptides
corresponding to this gene are useful for the study, diagnosis and treatment of immune,
autoimmune, and inflammatory disorders. Furthermore, this gene product may be
35 involved in the regulation of cytokine production, antigen presentation, or other
processes that may also suggest a usefulness in the treatment of cancer (e.g. by
boosting immune responses). Since the gene is expressed in cells of lymphoid origin,

the gene or protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Expression of this gene product in neutrophils strongly indicates a role for this protein in immune function and immune surveillance.

- 5 Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:91 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome.
- 10 Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1960 of SEQ ID NO:91, b is an integer of 15 to 1974, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:91, and where b is greater than or equal to a + 14.

15

FEATURES OF PROTEIN ENCODED BY GENE NO: 82

- 20 This gene is expressed primarily in primary dendritic cells, and to a lesser extent in neutrophils, monocytes, and osteoblasts.

- Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune and hematopoietic conditions. Similarly, polypeptides and
- 25 antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune and hematopoietic systems, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. immune, cancerous and wounded tissues) or bodily
- 30 fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

- Preferred epitopes include those comprising a sequence shown in SEQ ID
- 35 NO:217 as residues: Gly-47 to Arg-53.

 The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the study and treatment of immune,

inflammatory and hematopoietic disorders. Furthermore, this gene product may be involved in the regulation of cytokine production, antigen presentation, or other processes that may also suggest a usefulness in the treatment of cancer (e.g. by boosting immune responses). Since the gene is expressed in cells of lymphoid origin, the gene or protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues. Expression of this gene product in neutrophils and primary dendritic cells also strongly indicates a role for this protein in immune function and immune surveillance.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:92 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1409 of SEQ ID NO:92, b is an integer of 15 to 1423, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:92, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 83

One embodiment of this gene comprises polypeptides of the following amino acid sequence:

MPKRKVTFGVGDEEDEDEIIVPKKKLVDPVAGSGGPGSRFKGKHS LDSDEEE
 DDDGGSSKYDILASEDVEGQEAATLPSEGGVRITPFNLQEEMEEGHFDADGN
 YFLNRDAQIRDSWLDNIDWVKIRERPPGQRQASDSEEDSLGQTSMSAQALLEG
 LLELLLPRETVAGALRRLGARGGGKGRKGPQPSSPQRLDRLSGLADQMVAR
 GNLGVYQETRERLAMRLKGLGCQTLGPHNPTPPPSLDMFAEELAELEELETPTPT
 QRGAEESRGDGLVDVMWEYKWENTGDAELYGPFTSAQMQTWVSEGYFPDGV
 YCRKLDPPGGQFYNSKRIDFDLYT (SEQ ID NO:373), TFQGVGDEEDEDEIIVP
 KKKLVDP (SEQ ID NO:374), PGSRFKGKHS LDSDEEEDDDGGSSKY (SEQ ID
 NO:375), EAATLPSEGGVRITPFNLQEEMEEG (SEQ ID NO:376), FLNRDAQIRDS
 WLDNIDWVKIRERPPGQR (SEQ ID NO:377), SLGQTSMSAQALLEG LLELLL
 PRETV (SEQ ID NO:378), RGGGKGRKGPQPSSPQRLDRLSGLADQ (SEQ ID
 NO:379), QETRERLAMRLKGLGCQTLGPHNP (SEQ ID NO:380), DMFAEELAELEE
 LETPTPTQRGAEESRGD (SEQ ID NO:381), and/or ELYGPFTSAQMQTW

VSEGYFPDGVYCRKLD (SEQ ID NO:382). An additional embodiment is the polynucleotides encoding these polypeptides.

This gene is expressed primarily in fetal lung, stromal cells and lymphoma cells.

Therefore, polynucleotides and polypeptides of the invention are useful as
5 reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, haemopoietic and respiratory disorders and cancer. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a
10 number of disorders of the above tissues or cells, particularly of the haemopoietic and respiratory systems, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. lung, cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder,
15 relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:218 as residues: Met-1 to Trp-15, Thr-52 to Met-58.

The tissue distribution indicates that polynucleotides and polypeptides
20 corresponding to this gene are useful for the treatment and diagnosis of diseases of the haemopoietic and respiratory systems. Protein, as well as, antibodies directed against the protein may show utility as a tissue-specific marker and/or immunotherapy target for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available
25 and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:93 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more
30 polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1351 of SEQ ID NO:93, b is an integer of 15 to 1365, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:93, and where b is greater than or equal to a + 14.

35

FEATURES OF PROTEIN ENCODED BY GENE NO: 84

In specific embodiments, polypeptides of the invention comprise the following amino acid sequence: PHSSRVSFQLQSLSF (SEQ ID NO:383), RGQPRPCVSGVCLS PHSRFECCSFYLQGLPALRCSRTPPGCHFFRVFPSCPFSSSRSPSCFT HICPV
 5 VRIQFSRALWVSTCLVLAITPGKWLLPEDRALSLMLLASLQCCPPPGAWWMQ VLTHKGRQAGLG PGVSSRPL (SEQ ID NO:384, S NIKSLPPTNSLSLLRA QTGTDCAVSPGLAGPCHQRGLEDTPGPRPACLPCLCVSTCIHQAPKGGGQHWRE
 EA SSIRDRAISSGRSHFPGVMAKTKHVDTHNARENWIRTTGQMWVKHEG EREEEKGHEGKTLKK (SEQ ID NO:385), VCLSPHSRFECCSFYLQGLPALRC
 10 (SEQ ID NO:386), QFSRALWVSTCLVLAITPGKWLLPEDR (SEQ ID NO:387), SLSLLRAQTGTDCAVSPGLAGPCHQRG (SEQ ID NO:388), and/or SGRSHFPG VMAKTKHVDTHNARENWIRT (SEQ ID NO:389). Polynucleotides encoding these polypeptides are also encompassed by the invention. When tested against U937 cell
 15 lines, supernatants removed from cells containing this gene activated the GAS (gamma activating sequence) promoter element. Thus, it is likely that this gene activates myeloid cells, including their progenitors, through the Jak-STAT signal transduction pathway. GAS is a promoter element found upstream of many genes which are involved in the Jak-STAT pathway. The Jak-STAT pathway is a large, signal transduction pathway involved in the differentiation and proliferation of cells. Therefore, activation of the Jak-
 20 STAT pathway, reflected by the binding of the GAS element, can be used to indicate proteins involved in the proliferation and differentiation of cells.

This gene is expressed primarily in T-cells and lungs.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a
 25 biological sample and for diagnosis of diseases and conditions which include, but are not limited to, respiratory and immune diseases. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune and respiratory systems,
 30 expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. pulmonary, immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, pulmonary surfactant or sputum, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene
 35 expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:219 as residues: His-38 to Ala-43.

5 The tissue distribution in T-cells and lung tissue, combined with the detected GAS biological activity indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of disorders of the respiratory and immune systems. This gene product may be involved in the regulation of cytokine production, antigen presentation, or other processes that may also suggest a usefulness in the treatment of cancer (e.g. by boosting immune responses). Since the gene is expressed in cells of lymphoid origin, the natural gene product may be involved in
10 immune functions. Therefore it may be also used as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as
15 host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's disease, scleroderma and tissues. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in
20 the differentiation and/or proliferation of various cell types. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues. The protein may show utility in modulating the immune response to various pulmonary disorders or conditions, particularly in emphysema, or ARDS.

25 Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:94 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome.
30 Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 742 of SEQ ID NO:94, b is an integer of 15 to 756, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:94, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 85

In specific embodiments, polypeptides of the invention comprise the following amino acid sequence: ARVEVQQQGPGAKVDAGEGQ (SEQ ID NO:390), WVVL
5 SQLQA QGVAGMMCSYPEGQKKGKEATRSHRWVPRSLPGMGSLAAPH
NPWLAPLALLEIPXPVLCEWKRKLIAL EEVSECRPGVGGGGGFLSXCRR
GHLSFLSGAPYPLFPISPLX (SEQ ID NO:391), ELRHGGPRQVKDSFLDYM
GYPDEDRA GPPSRWFPRERFLSPPTV VPLCVELRLGFESGMGWGVPGSSHS
EGGPEARWPLIAPMYTVTQWFQRPNSGRGPQPPQXRGEIGKRGY GAPER
10 KLRWPLLXWERXPPPPPTPGRHSETSSSAISFLFHSQRTGWGISSANGASQGL
LWGAARXLPIP GRDLGTHLWDLVASFPFFCPSPG (SEQ ID NO:392), PEGQKK
GKEATRSHRWVPRSLPGM (SEQ ID NO:393), LRLGFESGMGWGVPGSSHSE
GPEAR (SEQ ID NO:394), and/or HSQRTGWGISSANGASQGLLWGA (SEQ ID
NO:395). Polynucleotides encoding these polypeptides are also encompassed by the
15 invention.

This gene is expressed primarily in eosinophils, dendritic cells, Jurkat cells and tonsils.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a
20 biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune, or hematopoietic disorders, particularly inflammatory, autoimmune, allergy, and hypersensitivity conditions. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders
25 of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g. immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the
30 standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in a variety of immune and hematopoietic-specific cells and tissues indicates that polynucleotides and polypeptides corresponding to this gene are useful for modifying the response of the immune system in autoimmune diseases
35 and inflammatory conditions. Moreover, polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of hematopoietic related disorders such as anemia, pancytopenia, leukopenia, thrombocytopenia or

leukemia since stromal cells are important in the production of cells of hematopoietic lineages. The uses include bone marrow cell ex vivo culture, bone marrow transplantation, bone marrow reconstitution, radiotherapy or chemotherapy of neoplasia. The gene product may also be involved in lymphopoiesis, therefore, it can be used in immune disorders such as infection, inflammation, allergy, immunodeficiency etc. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. It may also have a very wide range of biological activities. Typical of these are cytokine, cell proliferation/differentiation modulating activity or induction of other cytokines; immunostimulating/immunosuppressant activities (e.g. for treating human immunodeficiency virus infection, cancer, autoimmune diseases and allergy); regulation of hematopoiesis (e.g. for treating anaemia or as adjunct to chemotherapy); stimulation or growth of bone, cartilage, tendons, ligaments and/or nerves (e.g. for treating wounds, stimulation of follicle stimulating hormone (for control of fertility); chemotactic and chemokinetic activities (e.g. for treating infections, tumors); hemostatic or thrombolytic activity (e.g. for treating haemophilia, cardiac infarction etc.); anti-inflammatory activity (e.g. for treating septic shock, Crohn's disease); as antimicrobials; for treating psoriasis or other hyperproliferative diseases; for regulation of metabolism, and behaviour. Also contemplated is the use of the corresponding nucleic acid in gene therapy procedures. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:95 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 924 of SEQ ID NO:95, b is an integer of 15 to 938, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:95, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 86

This gene is expressed primarily in cells from fibrosarcoma tumors.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, muscle, or endothelial disorders, particularly fibrosarcomas and fibroids. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the skeleto-muscular system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. skeleto-muscular, cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in fibrosarcoma tissue indicates that polynucleotides and polypeptides corresponding to this gene are useful for the detection, treatment, and/or prevention of various muscle disorders, in particular fibrosarcomas. In addition, the expression of this gene product in synovium would suggest a role in the detection and treatment of disorders and conditions affecting the skeletal system, in particular osteoporosis as well as disorders afflicting connective tissues (e.g. arthritis, trauma, tendonitis, chondromalacia and inflammation). The gene or protein product of this gene may also show utility in modulating the immune response to proliferative tissues. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:96 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 914 of SEQ ID NO:96, b is an integer of 15 to 928, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:96, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 87

5 This gene is expressed primarily in helper T-Cells, cerebellum, and to a lesser extent, in mesangial cells, fetal lung, fetal liver, cortex, and adipose tissue.

 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are
10 not limited to, immune, or neural disorders, particularly, for modulation of immune responses to viral or bacterial infections, or neurodeficiencies. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of
15 this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g. renal, developmental, pulmonary, hepatic, neural, metabolic, immune, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, bile, amniotic fluid, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene
20 expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

 The tissue distribution in helper T-cells indicates that polynucleotides and polypeptides corresponding to this gene are useful for modifying the immune response to foreign agents such as bacteria or virus. In addition, this gene product may be
25 involved in the regulation of cytokine production, antigen presentation, or other processes that may also suggest a usefulness in the treatment of cancer (e.g. by boosting immune responses). Since the gene is expressed in cells of lymphoid origin, the natural gene product may be involved in immune functions. Therefore it may be also used as an agent for immunological disorders including arthritis, asthma,
30 immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue
35 injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's disease, scleroderma and tissues. In addition, this gene product may have commercial utility in the expansion of stem cells and committed

progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Moreover, based upon the expression within the cerebellum and cortex, polynucleotides and polypeptides corresponding to this gene are useful for the detection/treatment of neurodegenerative disease states, behavioural disorders, or inflammatory conditions such as Alzheimers Disease, Parkinsons Disease, Huntingtons Disease, Tourette Syndrome, meningitis, encephalitis, demyelinating diseases, peripheral neuropathies, neoplasia, trauma, congenital malformations, spinal cord injuries, ischemia and infarction, aneurysms, hemorrhages, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and preception. In addition, elevated expression of this gene product in regions of the brain indicates that it plays a role in normal neural function. Potentially, this gene product is involved in synapse formation, neurotransmission, learning, cognition, homeostasis, or neuronal differentiation or survival. Moreover, the gene or gene product may also play a role in the treatment and/or detection of developmental disorders associated with the developing embryo, sexually-linked disorders, or disorders of the cardiovascular system. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:97 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1701 of SEQ ID NO:97, b is an integer of 15 to 1715, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:97, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 88

When tested against U937 cell lines, supernatants removed from cells containing this gene activated the GAS (gamma activating sequence) promoter element. Thus, it is likely that this gene activates promyelocytic cells through the JAK-STAT signal transduction pathway. GAS is a promoter element found upstream of many

genes which are involved in the Jak-STAT pathway. The Jak-STAT pathway is a large, signal transduction pathway involved in the differentiation and proliferation of cells.

Therefore, activation of the Jak-STAT pathway, reflected by the binding of the GAS element, can be used to indicate proteins involved in the proliferation and differentiation

5 of cells. In specific embodiments, polypeptides of the invention comprise the following amino acid sequence:

FIMKLLYQLLMLTTSSSYSLITHLCYSIFLCSFYFHFPCNVSLFVLISEEFTYD
(SEQ ID NO:396), LMLTTSSSYSLITHLCYSIFL (SEQ ID NO:397), LCSFYFH
FPCNVSLFVLISEE (SEQ ID NO:398), MRKNIFAILDKMLTCLIINELFRNQYKET
10 NITREVKIKGTEENGIAQMSYKAI (SEQ ID NO:399), DKMLTCLIINELFRNQ
YKETN (SEQ ID NO:400), and/or NITREVKIKGTEENGIAQMSY (SEQ ID
NO:401). Polynucleotides encoding these polypeptides are also encompassed by the
invention.

This gene is expressed primarily in fetal lung.

15 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, pulmonary and developmental disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes
20 for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the developmental and pulmonary systems, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g. Pulmonary, developmental, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, pulmonary surfactant or sputum,
25 amniotic fluid, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution of this gene only in fetal lung, combined with the
30 detected GAS biological activity indicates that it plays a key role in development of the pulmonary system. This would suggest that misregulation of the expression of this protein product in the adult could lead to lymphoma or sarcoma formation, particularly in the lung. It may also be involved in the predisposition to certain pulmonary defects such as pulmonary edema and embolism, bronchitis and cystic fibrosis. Moreover, the
35 protein product of this gene may be beneficial in the treatment of underdeveloped lung tissue, as exists in premature infants, both through the use of antibodies directed against the protein, through a gene therapy-based regime, or through the action of the protein

itself, either directly or indirectly. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:98 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 664 of SEQ ID NO:98, b is an integer of 15 to 678, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:98, and where b is greater than or equal to a + 14.

15

FEATURES OF PROTEIN ENCODED BY GENE NO: 89

When tested against U937 cell lines, supernatants removed from cells containing this gene activated the GAS (gamma activating sequence) promoter element. Thus, it is likely that this gene activates myeloid cells, including their progenitors, through the Jak-STAT signal transduction pathway. GAS is a promoter element found upstream of many genes which are involved in the Jak-STAT pathway. The Jak-STAT pathway is a large, signal transduction pathway involved in the differentiation and proliferation of cells. Therefore, activation of the Jak-STAT pathway, reflected by the binding of the GAS element, can be used to indicate proteins involved in the proliferation and differentiation of cells. In specific embodiments, polypeptides of the invention comprise the following amino acid sequence: GISERKP (SEQ ID NO:402). Polynucleotides encoding these polypeptides are also encompassed by the invention.

This gene is expressed primarily in brain.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, neural or immune disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the central nervous system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues

or cell types (e.g. neural, immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:224 as residues: Ile-40 to Trp-50.

The tissue distribution in brain combined with the detected GAS biological activity indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of central nervous system disorders. Moreover, polynucleotides and polypeptides corresponding to this gene are useful for the detection/treatment of neurodegenerative disease states, behavioural disorders, or inflammatory conditions such as Alzheimers Disease, Parkinsons Disease, Huntingtons Disease, Tourette Syndrome, meningitis, encephalitis, demyelinating diseases, peripheral neuropathies, neoplasia, trauma, congenital malformations, spinal cord injuries, ischemia and infarction, aneurysms, hemorrhages, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and perception. In addition, elevated expression of this gene product in regions of the brain indicates that it plays a role in normal neural function. Potentially, this gene product is involved in synapse formation, neurotransmission, learning, cognition, homeostasis, or neuronal differentiation or survival. Moreover, the gene or gene product may also play a role in the treatment and/or detection of developmental disorders associated with the developing embryo, sexually-linked disorders, or disorders of the cardiovascular system. Furthermore, the protein may show utility in modulating the immune response to various neurodegenerative conditions. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:99 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome.

Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1527 of SEQ ID NO:99, b is an integer of 15

to 1541, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:99, and where b is greater than or equal to a + 14.

5 FEATURES OF PROTEIN ENCODED BY GENE NO: 90

In specific embodiments, polypeptides of the invention comprise the following amino acid sequence: QSPAVSYTVTSQVPWGLGLLAGEKR (SEQ ID NO:403), LP SHPLRPLTFS SAMCMHLPPPLCRRALSAPFATQHRPWSVAAACLPRIHQN
 10 PLDAEYPSGCCRMSFLPAACSNIYSQECH YTLMSHSEASTLQXAQLL (SEQ ID NO:404), MLLQAAGRKL MRQQPDGYSASRGFWWMRGRQAAATLHGRCWVA KGADSAAL RQRGGGRCMHIAD EKVRLSGCDGS (SEQ ID NO:405), LCRRA ALSAPFATQHRPWSVAAACL (SEQ ID NO:406), RGFWWMRGRQAAATLHGR CWVAKG (SEQ ID NO:407), and/or QRGGGRCMHIAD EKVRLSGCDG (SEQ ID
 15 NO:408). Polynucleotides encoding these polypeptides are also encompassed by the invention.

This gene is expressed primarily in neutrophils.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a
 20 biological sample and for diagnosis of diseases and conditions which include, but are not limited to, inflammatory and immune conditions. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene
 25 at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily
 30 fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:225 as residues: Pro-34 to His-39, Pro-44 to His-54.

The tissue distribution in neutrophils indicates that polynucleotides and polypeptides corresponding to this gene are useful for the study, diagnosis, and
 35 treatment of inflammatory, general immune, and infectious diseases. Moreover, the expression of this gene indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem

cells. This gene product may be involved in the regulation of cytokine production, antigen presentation, or other processes that may also suggest a usefulness in the treatment of cancer (e.g. by boosting immune responses). Since the gene is expressed in cells of lymphoid origin, the natural gene product may be involved in immune functions. Therefore it may be also used as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's disease, scleroderma and tissues. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:100 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 867 of SEQ ID NO:100, b is an integer of 15 to 881, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:100, and where b is greater than or equal to a + 14.

30 FEATURES OF PROTEIN ENCODED BY GENE NO: 91

When tested against Jurkat cell lines, supernatants removed from cells containing this gene activated the GAS (gamma activating sequence) promoter element. Thus, it is likely that this gene activates myeloid cells through the JAK-STAT signal transduction pathway. GAS is a promoter element found upstream of many genes which are involved in the Jak-STAT pathway. The Jak-STAT pathway is a large, signal transduction pathway involved in the differentiation and proliferation of cells.

Therefore, activation of the Jak-STAT pathway, reflected by the binding of the GAS element, can be used to indicate proteins involved in the proliferation and differentiation of cells. In addition, contact of cells with supernatant expressing the product of this gene has been shown to increase the permeability of the plasma membrane of stromal cells to calcium. Thus, it is likely that the product of this gene is involved in a signal transduction pathway that is initiated when the product binds a receptor on the surface of the plasma membrane of both stromal, in addition to other cell-lines or tissue cell types. Thus, polynucleotides and polypeptides have uses which include, but are not limited to, activating stromal cells. Binding of a ligand to a receptor is known to alter intracellular levels of small molecules, such as calcium, potassium and sodium, as well as alter pH and membrane potential. Alterations in small molecule concentration can be measured to identify supernatants which bind to receptors of a particular cell. In specific embodiments, polypeptides of the invention comprise the following amino acid sequence:

15 THPSHSIVIQSTVSLCLTASSRRKKSDCLSLCQVSCSQRPGSHKTNVAWGFLM
SRVHFSVRWVSGGRI TGAICKESSLPCKEIQKGKACYFCHHPAQQSTPFSHI
(SEQ ID NO:409, VIQSTVSLCLTASSRRKKSDCLSLCQV (SEQ ID NO:410),
and/or ICKESSLPCKEIQKGKACYFCHHPAQQ (SEQ ID NO:411). Polynucleotides
encoding these polypeptides are also encompassed by the invention.

20 This gene is expressed primarily in neutrophils, and to a lesser extent, in cord blood.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune or developmental disorders, particularly inflammatory conditions. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune and hemopoietic systems, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. immune, developmental, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, amniotic fluid, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

35 Preferred epitopes include those comprising a sequence shown in SEQ ID NO:226 as residues: Glu-32 to Arg-37.

The tissue distribution in neutrophils, combined with the detected GAS and calcium flux biological activities, indicates that polynucleotides and polypeptides corresponding to this gene are useful for the study and treatment of inflammatory, infectious, and hemopoietic disorders. Similarly, expression within cord blood indicates that this protein may play a role in the regulation of cellular division, and may show utility in the diagnosis and treatment of cancer and other proliferative disorders, particularly of the developing hematopoietic system. Similarly, developmental tissues rely on decisions involving cell differentiation and/or apoptosis in pattern formation. Thus, this protein may also be involved in apoptosis or tissue differentiation and could again be useful in cancer therapy. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:101 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 933 of SEQ ID NO:101, b is an integer of 15 to 947, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:101, and where b is greater than or equal to a + 14.

25 FEATURES OF PROTEIN ENCODED BY GENE NO: 92

The gene encoding the disclosed cDNA is thought to reside on chromosome 15. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 15.

30 This gene is expressed primarily in macrophages, T cells, dendritic cells, testes and pancreas tumors.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune disorders including testis and pancreas tumors. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a

number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. immune, metabolic, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, bile, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:227 as residues: Gln-85 to Lys-91, Pro-106 to Ser-117, Pro-124 to Ala-130, Trp-154 to Trp-160.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of immune disorders such as testes and pancreas tumors. Furthermore, polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and/or treatment of hematopoietic disorders. This gene product is primarily expressed in hematopoietic cells and tissues, suggesting that it plays a role in the survival, proliferation, and/or differentiation of hematopoietic lineages. Expression of this gene product in T cells and primary dendritic cells also strongly indicates a role for this protein in immune function and immune surveillance. Since the gene is expressed in cells of lymphoid origin, the natural gene product may be involved in immune functions. Therefore it may be also used as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's disease, scleroderma and tissues. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:102 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome.

Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1355 of SEQ ID NO:102, b is an integer of 15 to 1369, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:102, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 93

10 This gene is expressed primarily in brain tissue from a patient suffering from manic depression.

 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, neural disorders, particularly manic depression. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the brain, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. brain, cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

25 The tissue distribution in brain tissue indicates that polynucleotides and polypeptides corresponding to this gene are useful for diagnosis of manic depression and other disorders of the CNS. Furthermore, polynucleotides and polypeptides corresponding to this gene are useful for the detection/treatment of neurodegenerative disease states and behavioural disorders such as Alzheimers Disease, Parkinsons Disease, Huntingtons Disease, Tourette Syndrome, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and perception. In addition, elevated expression of this gene product in regions of the brain indicates that it plays a role in normal neural function.

35 Potentially, this gene product is involved in synapse formation, neurotransmission, learning, cognition, homeostasis, or neuronal differentiation or survival. Moreover, the gene or gene product may also play a role in the treatment and/or detection of

developmental disorders associated with the developing embryo, sexually-linked disorders, or disorders of the cardiovascular system. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

- 5 Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:103 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome.
- 10 Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1217 of SEQ ID NO:103, b is an integer of 15 to 1231, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:103, and where b is greater than or equal to a + 14.

15

FEATURES OF PROTEIN ENCODED BY GENE NO: 94

This gene is expressed primarily in anergic T-cells.

- 20 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune or hematopoietic disorders, particularly autoimmune disorders such as lupus. Similarly, polypeptides and antibodies directed to these polypeptides are
- 25 useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine,
- 30 synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

- The tissue distribution in T-cells indicates that polynucleotides and polypeptides
- 35 corresponding to this gene are useful for the diagnosis and treatment of a variety of immune system disorders. Moreover, the protein product of this gene may play a role in the regulation of the proliferation; survival; differentiation; and/or activation of

potentially all hematopoietic cell lineages, including blood stem cells. This gene product may be involved in the regulation of cytokine production, antigen presentation, or other processes that may also suggest a usefulness in the treatment of cancer (e.g. by boosting immune responses). Expression of this gene product in T cells also strongly indicates a role for this protein in immune function and immune surveillance. Since the gene is expressed in cells of lymphoid origin, the natural gene product may be involved in immune functions. Therefore it may be also used as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's disease, scleroderma and tissues. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:104 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1228 of SEQ ID NO:104, b is an integer of 15 to 1242, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:104, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 95

This gene is expressed primarily in the spinal cord.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are

not limited to, neural disorders, particularly CNS, PNS, and a variety of congenital malformations of the spinal column and injuries of the spinal cord. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s) present in a biological sample. For a number of disorders of the above tissues or cells, particularly of the central nervous system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. CNS, cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:230 as residues: Ser-44 to His-52.

The tissue distribution in spinal cord tissue indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and/or treatment of disorders of the brain and nervous system. Such involvement may impact many processes, such as learning and cognition. It may also be useful in the treatment of such neurodegenerative disorders as schizophrenia; ALS; or Alzheimer's. Protein, as well as, antibodies directed against the protein may show utility as a tissue-specific marker and/or immunotherapy target for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:105 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1137 of SEQ ID NO:105, b is an integer of 15 to 1151, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:105, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 96

This gene is expressed primarily in smooth muscle.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, muscular, vascular, or cardiopulmonary disorders, particularly a variety of diseases that include wasting and muscle mass loss including amyotrophic lateral sclerosis, embolism, atherosclerosis, stroke, and aneurysm. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the neuromuscular system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. muscle, cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:231 as residues: Leu-37 to Trp-44.

The tissue distribution in smooth muscle indicates that polynucleotides and polypeptides corresponding to this gene are useful for the detection, treatment, and/or prevention of various muscle disorders, such as muscular dystrophy, cardiomyopathy, fibroids, myomas, vascular disorders, and rhabdomyosarcomas. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:106 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1614 of SEQ ID NO:106, b is an integer of 15 to 1628, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:106, and where b is greater than or equal to a + 14.

35

FEATURES OF PROTEIN ENCODED BY GENE NO: 97

This gene is expressed primarily in the brain.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, disorders affecting the brain and central nervous system, such as Alzheimer's disease. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the brain and central nervous system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. brain, cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in brain tissue indicates that polynucleotides and polypeptides corresponding to this gene are useful for the detection/treatment of neurodegenerative disease states and behavioural disorders such as Alzheimers Disease, Parkinsons Disease, Huntingtons Disease, Tourette Syndrome, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, panic disorder, learning disabilities, ALS, psychoses, autism, and altered bahaviors, including disorders in feeding, sleep patterns, balance, and perception. In addition, the gene or gene product may also play a role in the treatment and/or detection of developmental disorders associated with the developing embryo. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:107 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1451 of SEQ ID NO:107, b is an integer of 15 to 1465, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:107, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 98

5 In specific embodiments, polypeptides of the invention comprise the following amino acid sequence: SLQVLRTLGSKCGDFLRSRFCKDVLPLAGSLVT QAPISARAGPVYSHTLAFKLQLAVLQGLGPLCERLDLGEGLNKNVADACLIYLS VKQPVKLQEAARSVFL HLMKVDPDSTWFLNLNELYCPVQFTPPHPSLHPVQLX GASGQQNPXHDQRAPAAQGAAVTLLPHHRGHRSL PYCQPEAGLTPPRP (SEQ
10 ID NO:412), GADGNVSDFDNEEEEQSVPPKVDENDTRPDVEPPLPLQIQIAM DVMERCIHLLSDKNLQIRLKVLDVLDL CVVVLQSHKNQLLPLAHQAWPSL VHRLTRDAPLAVLRAFKFYVPWEASVVTFFAAGSAKMSCQSWLAP (SEQ ID NO:413), TLGSKCGDFLRSRFCKDVLPLAGSL (SEQ ID NO:414), PVYSHTL AFKLQLAVLQGLGPLCERLDLG (SEQ ID NO:415), SVPPKVDENDTRPDV
15 EPPLPLQIQIAM (SEQ ID NO:416), and/or WPSLVHRLTRDAPLAVLRAFK FYVPW (SEQ ID NO:417). Polynucleotides encoding these polypeptides are also encompassed by the invention.

This gene is expressed primarily in kidney cortex, hemangiopericytoma, fetal spleen, infant brain, and to a lesser extent, in pancreas, lymph node, fetal liver, ovarian
20 tumor, T-cells and other tissues.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, renal, immune, neural, or developmental disorders, particularly tumors. Similarly,
25 polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the reproductive system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g. renal, immune, neural, developmental,
30 reproductive, ovarian, hepatic, metabolic, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, bile, amniotic fluid, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

35 Preferred epitopes include those comprising a sequence shown in SEQ ID NO:233 as residues: Pro-24 to Pro-37.

The tissue distribution in proliferating tissues and cells, combined with its distribution in developing tissues indicates that polynucleotides and polypeptides corresponding to this gene are useful for diagnosing and treating tumors. Thus this protein may also be involved in apoptosis or tissue differentiation and could again be useful in cancer therapy. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:108 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1251 of SEQ ID NO:108, b is an integer of 15 to 1265, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:108, and where b is greater than or equal to a + 14.

20 FEATURES OF PROTEIN ENCODED BY GENE NO: 99

In specific embodiments, polypeptides of the invention comprise the following amino acid sequence: SLGISTFGIMVFSVYFGGIMISIPYSGISFGNKKELNID SCYNMVNLKNIMFSERSQT (SEQ ID NO:418), HASGNNDPLWFLTYL (SEQ ID NO:419), MVFSVYFGGIMISIPYSGISF (SEQ ID NO:420), and/or FGNKKELNID SCYNMVNLKN (SEQ ID NO:421). Polynucleotides encoding these polypeptides are also encompassed by the invention.

This gene is expressed primarily in T-cells, spleen, and pancreas.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune or endocrine disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g. Immune, hematopoietic, endocrine, pancreatic, cancerous and wounded

tissues) or bodily fluids (e.g. lymph, serum, bile, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

- 5 Preferred epitopes include those comprising a sequence shown in SEQ ID NO:234 as residues: Thr-24 to Arg-29.

The tissue distribution of this gene predominantly in cell types or tissues associated with the immune system indicates that the gene could be important for the treatment or detection of immune or hematopoietic disorders including, but not limited to, arthritis, asthma, immunodeficiency diseases and leukemia. Moreover, the expression within pancreatic tissues indicates that the protein product of this gene may be useful in the treatment or prevention of a variety of metabolic disorders, such as diabetes. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

- 10 to, arthritis, asthma, immunodeficiency diseases and leukemia. Moreover, the expression within pancreatic tissues indicates that the protein product of this gene may be useful in the treatment or prevention of a variety of metabolic disorders, such as diabetes. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.
- 15 Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:109 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome.
- 20 Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 992 of SEQ ID NO:109, b is an integer of 15 to 1006, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:109, and where b is greater than or equal to a + 14.

25

FEATURES OF PROTEIN ENCODED BY GENE NO: 100

- The gene encoding the disclosed cDNA is believed to reside on the X chromosome. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for the X chromosome.

This gene is expressed primarily in urinary bladder carcinoma HSC172 cells, and to a lesser extent, in human adult heart, lung, osteoclastoma, and liver.

- Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, urogenital, or renal disorders, particularly urinary bladder carcinoma and

other cancers. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the bladder, expression of this gene at significantly higher or lower levels may be
5 routinely detected in certain tissues or cell types (e.g. renal, cardiopulmonary, hepatic, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, bile, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the
10 disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:235 as residues: Gly-18 to Lys-23, Pro-31 to Gly-38.

The tissue distribution in urinary bladder carcinoma indicates that polynucleotides and polypeptides corresponding to this gene are useful for diagnosis
15 and therapeutic targeting of urinary bladder carcinoma, osteoclastoma, and other cancers. Additionally, the tissue distribution in heart, lung and osteocarcinoma indicates an indication for the use of this gene and gene product in diagnosis and treatment of disorders in the heart and lung. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above
20 listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:110 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the
25 scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1244 of SEQ ID NO:110, b is an integer of 15 to 1258, where both a and b correspond to the positions of nucleotide residues shown
30 in SEQ ID NO:110, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 101

35 In specific embodiments, polypeptides of the invention comprise the following amino acid sequence:

MNSFSVIASIVLLPFPGLSVSACLPSHSHQCKTFILLFLPSSEKTLXXXPP

SHSSTLGGQGGQIMRSGDRXHXG (SEQ ID NO:422), VVFFXXFFEMESH
 SVAQAGVQWRNLGSLQAL PPGFMPFSCLSLPGSWDYRRPPPPSPANLXCIF
 SRDGGHHVSQXGLDLLTS (SEQ ID NO:423), IVVLLPFPGLSVSACLPS
 HSHQCKTFIL (SEQ ID NO:424), and/or PGFMPFSCLSLPGSWDYRRPPPPSPAN
 5 (SEQ ID NO:425). Polynucleotides encoding these polypeptides are also encompassed
 by the invention.

This gene is expressed primarily in adipose tissue.

Therefore, polynucleotides and polypeptides of the invention are useful as
 reagents for differential identification of the tissue(s) or cell type(s) present in a
 10 biological sample and for diagnosis of diseases and conditions which include, but are
 not limited to, obesity and other metabolic disorders. Similarly, polypeptides and
 antibodies directed to these polypeptides are useful in providing immunological probes
 for differential identification of the tissue(s) or cell type(s). For a number of disorders
 of the above tissues or cells, particularly of the endocrine system, expression of this
 15 gene at significantly higher or lower levels may be routinely detected in certain tissues
 or cell types (e.g. adipose, metabolic, neural, and cancerous and wounded tissues) or
 bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or
 another tissue or cell sample taken from an individual having such a disorder, relative to
 the standard gene expression level, i.e., the expression level in healthy tissue or bodily
 20 fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID
 NO:236 as residues: Arg-28 to Asn-33.

The tissue distribution in adipose tissue indicates that polynucleotides and
 polypeptides corresponding to this gene are useful for the treatment of obesity and other
 25 metabolic and endocrine conditions or disorders. Furthermore, the protein product of
 this gene may show utility in ameliorating conditions which occur secondary to aberrant
 fatty-acid metabolism (e.g. aberrant myelin sheath development), either directly or
 indirectly. Protein, as well as, antibodies directed against the protein may show utility
 as a tumor marker and/or immunotherapy targets for the above listed tissues.

30 Many polynucleotide sequences, such as EST sequences, are publicly available
 and accessible through sequence databases. Some of these sequences are related to SEQ
 ID NO:111 and may have been publicly available prior to conception of the present
 invention. Preferably, such related polynucleotides are specifically excluded from the
 scope of the present invention. To list every related sequence is cumbersome.
 35 Accordingly, preferably excluded from the present invention are one or more
 polynucleotides comprising a nucleotide sequence described by the general formula of
 a-b, where a is any integer between 1 to 1439 of SEQ ID NO:111, b is an integer of 15

to 1453, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:111, and where b is greater than or equal to a + 14.

5 FEATURES OF PROTEIN ENCODED BY GENE NO: 102

In specific embodiments, polypeptides of the invention comprise the following amino acid sequence: YRFKNPKCRLFSVPCR (SEQ ID NO:426), TQNRELLAWK
 PKGTDDICTSHNTTHIQKMPGE ANSCCPRGA KSYHIDCWPPALFPRCVA YLFL
 10 NKPATLRKKYYCKPYHTQLHPA WHREKSAFWIFETVSQS KQSLTSLVYS
 VNELLVLSNLAQWALG (SEQ ID NO:427), AWKPKGTDDICTSHNTTHIQKMP
 (SEQ ID NO:428), CPRGA KSYHIDCWPPALFPRCVA YL (SEQ ID NO:429), SYHI
 DCWPPALFPRCVA YLFLNKPAT (SEQ ID NO:430), and/or RKKYYCKPY
 HTQLHPA WHREKSAFWIFET (SEQ ID NO:431). Polynucleotides encoding these
 15 polypeptides are also encompassed by the invention.

This gene is expressed primarily in dendritic cells and activated monocytes.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are
 20 not limited to, immune or hematopoietic disorders, particularly inflammation, immune defects, multiple myeloma, or immunodeficiencies. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at
 25 significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily
 30 fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:237 as residues: Thr-27 to Arg-33.

The tissue distribution in dendritic cells and monocytes indicates that polynucleotides and polypeptides corresponding to this gene are useful for the
 35 diagnosis and treatment of inflammatory and immune disorders such as cancers, particularly of dendritic cells and monocytes, but also of hematopoietic progenitors. Similarly, polynucleotides and polypeptides corresponding to this gene are useful for

the treatment and diagnosis of hematopoietic related disorders such as anemia, pancytopenia, leukopenia, thrombocytopenia or leukemia since stromal cells are important in the production of cells of hematopoietic lineages. The uses include bone marrow cell ex vivo culture, bone marrow transplantation, bone marrow reconstitution, radiotherapy or chemotherapy of neoplasia. The gene product may also be involved in lymphopoiesis, therefore, it can be used in immune disorders such as infection, inflammation, allergy, immunodeficiency, etc. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types.

Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:112 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1538 of SEQ ID NO:112, b is an integer of 15 to 1552, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:112, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 103

25

When tested against K562 leukemia cell lines, supernatants removed from cells containing this gene activated the ISRE assay. Thus, it is likely that this gene activates leukemia cells through the Jak-STAT signal transduction pathway. The interferon-sensitive response element is a promoter element found upstream of many genes which are involved in the Jak-STAT pathway. The Jak-STAT pathway is a large, signal transduction pathway involved in the differentiation and proliferation of cells. Therefore, activation of the Jak-STAT pathway, reflected by the binding of the ISRE element, can be used to indicate proteins involved in the proliferation and differentiation of cells. The gene encoding the disclosed cDNA is thought to reside on chromosome 5.

Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 5.

This gene is expressed primarily in placenta, adipose tissue and fibroblasts.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, disorders of the skin, developing organs and metabolic disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the epidermal system metabolic system and embryogenesis, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. epidermal, cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of disorders of the epidermal system, metabolic system and embryogenesis. Furthermore, the tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and/or treatment of disorders of the placenta. Specific expression within the placenta indicates that this gene product may play a role in the proper establishment and maintenance of placental function. Alternately, this gene product may be produced by the placenta and then transported to the embryo, where it may play a crucial role in the development and/or survival of the developing embryo or fetus. Expression of this gene product in a vascular-rich tissue such as the placenta also indicates that this gene product may be produced more generally in endothelial cells or within the circulation. In such instances, it may play more generalized roles in vascular function, such as in angiogenesis. It may also be produced in the vasculature and have effects on other cells within the circulation, such as hematopoietic cells. It may serve to promote the proliferation, survival, activation, and/or differentiation of hematopoietic cells, as well as other cells throughout the body.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:113 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more

polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1475 of SEQ ID NO:113, b is an integer of 15 to 1489, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:113, and where b is greater than or equal to a + 14.

5

FEATURES OF PROTEIN ENCODED BY GENE NO: 104

In specific embodiments, polypeptides of the invention comprise the following amino acid sequence: ICLDSCSQVSVTSLWSFLRVHSLVQTLW (SEQ ID NO:432). Polynucleotides encoding these polypeptides are also encompassed by the invention.

This gene is expressed primarily in neutrophils.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, diseases of the immune system, including neutropenia, cancer, inflammatory diseases and allergies. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g. Immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:239 as residues: Ala-35 to Asp-44.

The tissue distribution in neutrophils indicates that polynucleotides and polypeptides corresponding to this gene are useful for treatment/diagnosis of diseases of the immune system since expression is primarily in neutrophils, and may be useful as a growth factor for the differentiation or proliferation of neutrophils for the treatment of neutropenia following chemotherapy or may be useful in the treatment of immune dysfunction or anti-inflammatory by inhibiting infiltration of neutrophils to the site of injury or distress. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:114 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 593 of SEQ ID NO:114, b is an integer of 15 to 607, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:114, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 105

This gene is expressed primarily in stromal cells.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune or hematopoietic disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in stromal cells indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and/or treatment of immune disorders. This gene product may be involved in the regulation of cytokine production, antigen presentation, or other processes that may also suggest a usefulness in the treatment of cancer (e.g. by boosting immune responses). Since the gene is expressed in cells of lymphoid origin, the natural gene product may be involved in immune functions. Therefore it may be also used as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous disease, inflammatory bowel disease,

sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus
5 erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's disease, scleroderma and tissues. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or
10 immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:115 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the
15 scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1484 of SEQ ID NO:115, b is an integer of 15 to 1498, where both a and b correspond to the positions of nucleotide residues shown
20 in SEQ ID NO:115, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 106

25 In specific embodiments, polypeptides of the invention comprise the following amino acid sequence: HYCC DFGTSL LGFYVPFHYYVHVMVNIILTTIDFYHYKFC CSQNANKHCFKHFQIMTTVPYLNINKENLRFKNIF K (SEQ ID NO:433), TSL LGFYVPFHYYVHVMVNIIL TTIDFY (SEQ ID NO:434), and/or FQIMTTVPYLN INKENLRFKNI (SEQ ID NO:435). Polynucleotides encoding these polypeptides are
30 also encompassed by the invention. The gene encoding the disclosed cDNA is believed to reside on chromosome 5. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 5.

This gene is expressed primarily in spleen, breast, placenta, ovarian cancer, and to a lesser extent, in B-cell lymphoma, pancreas tumor, osteoclastoma, thyroid, bone
35 marrow, fetal liver, and stromal cells.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a

biological sample and for diagnosis of diseases and conditions which include, but are not limited to, diseases characterized by immune cell activation and proliferation, particularly of the reproductive system. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential
5 identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g. Immune, reproductive, metabolic, skeletal, endocrine, hepatic, placental, ovarian, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, bile,
10 amniotic fluid, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID
15 NO:241 as residues: Ser-21 to Ser-27.

The tissue distribution in spleen and reproductive tissues indicates that the product of this gene is useful for modifying or detecting the proliferation or activation of cells in the hematopoietic system. The secreted protein can also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or
20 receptors, to identify agents that modulate their interactions and as nutritional supplements. It may also have a very wide range of biological activities. Typical of these are cytokine, cell proliferation/differentiation modulating activity or induction of other cytokines; immunostimulating/immunosuppressant activities (e.g. for treating human immunodeficiency virus infection, cancer, autoimmune diseases and allergy);
25 regulation of hematopoiesis (e.g. for treating anaemia or as adjunct to chemotherapy); stimulation or growth of bone, cartilage, tendons, ligaments and/or nerves (e.g. for treating wounds, stimulation of follicle stimulating hormone (for control of fertility); chemotactic and chemokinetic activities (e.g. for treating infections, tumors); hemostatic or thrombolytic activity (e.g. for treating haemophilia, cardiac infarction etc.); anti-
30 inflammatory activity (e.g. for treating septic shock, Crohn's disease); as antimicrobials; for treating psoriasis or other hyperproliferative diseases; for regulation of metabolism, and behaviour. Also contemplated is the use of the corresponding nucleic acid in gene therapy procedures. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the
35 above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ

ID NO:116 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome.

- Accordingly, preferably excluded from the present invention are one or more
 5 polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1783 of SEQ ID NO:116, b is an integer of 15 to 1797, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:116, and where b is greater than or equal to a + 14.

10

FEATURES OF PROTEIN ENCODED BY GENE NO: 107

In specific embodiments, polypeptides of the invention comprise the following amino acid sequence:

- 15 ISESMSLVRS LQFYRGKNRAERTVISSSSHSCHLIDLEFQPRSDGEVSISFLEKGV
 ELRWGMGLEDLIGLGLGVSTRRSTVRRKEPTKAGMHTACSEEMEPENREN
 (SEQ ID NO:436), DGSRSVAQARVQWHHRGSLPPLPPRFKQFPLRHLRVGGITG
 ACRHTQIIFVVLVQMGMFHHVG QAGLELLTSGDPPALASQSAGITGVSHSTRPKL
 LSWLPSDNLLGMALYSIQWALLANS LYFQVPSPLSML CAFLPLWVPSA (SEQ
 20 ID NO:437), RGKNRAERTVISSSSHSCHLIDLEFQP (SEQ ID NO:438), LGLGVST
 RRSTVRRKEPTKAGMHTACSEEMEP (SEQ ID NO:439), GDPPALASQSAGI
 TGVSHSTRPKL (SEQ ID NO:440), and/or ALYSIQWALLANS LYFQVPSPLSML
 (SEQ ID NO:441). Polynucleotides encoding these polypeptides are also encompassed
 by the invention.

- 25 This gene is expressed primarily in bone marrow.

- Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune or hematopoietic disorders, particularly bone marrow related
 30 diseases such as multiple myeloma, immunodeficiencies, and hematopoietic disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the bone marrow, expression of this gene at significantly higher or lower levels may be routinely
 35 detected in certain tissues and cell types (e.g. Immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a

disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:242 as residues: Gln-46 to Asn-56.

5 The tissue distribution in bone marrow indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of central nervous system disorders and hemopoietic system developmental disorders. Moreover, polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of hematopoietic related disorders such as anemia,
10 pancytopenia, leukopenia, thrombocytopenia or leukemia since stromal cells are important in the production of cells of hematopoietic lineages. The uses include bone marrow cell ex vivo culture, bone marrow transplantation, bone marrow reconstitution, radiotherapy or chemotherapy of neoplasia. The gene product may also be involved in lymphopoiesis, therefore, it can be used in immune disorders such as infection,
15 inflammation, allergy, immunodeficiency etc. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

20 Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:117 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome.
25 Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 938 of SEQ ID NO:117, b is an integer of 15 to 952, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:117, and where b is greater than or equal to a + 14.

30

FEATURES OF PROTEIN ENCODED BY GENE NO: 108

35 In specific embodiments, polypeptides of the invention comprise the following amino acid sequence: DRILLFYSRDGGTTSTKGPNPACCLFLLKKFYWNTA (SEQ ID NO:442), and/or DGQTTSKGPNPACCLFLLKKF (SEQ ID NO:443). Polynucleotides encoding these polypeptides are also encompassed by the invention.

This gene is expressed primarily in early stage human brain.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, neural disorders, particularly developmental disorders of the brain. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the early stage human brain, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g. neural, developmental, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, amniotic fluid, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:243 as residues: Asn-16 to Gln-21.

The tissue distribution in early stage brain indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of brain development disorders. In addition, polynucleotides and polypeptides corresponding to this gene are useful for the detection/treatment of neurodegenerative disease states, behavioural disorders, or inflammatory conditions such as Alzheimers Disease, Parkinsons Disease, Huntingtons Disease, Tourette Syndrome, meningitis, encephalitis, demyelinating diseases, peripheral neuropathies, neoplasia, trauma, congenital malformations, spinal cord injuries, ischemia and infarction, aneurysms, hemorrhages, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and preception. In addition, elevated expression of this gene product in regions of the brain indicates that it plays a role in normal neural function. Potentially, this gene product is involved in synapse formation, neurotransmission, learning, cognition, homeostasis, or neuronal differentiation or survival. Moreover, the gene or gene product may also play a role in the treatment and/or detection of developmental disorders associated with the developing embryo, sexually-linked disorders, or disorders of the cardiovascular system. Moreover, the expression within embryonic tissue indicates that this protein may play a role in the regulation of cellular division, and may show utility in the diagnosis and treatment of cancer and other proliferative disorders. Similarly,

developmental tissues rely on decisions involving cell differentiation and/or apoptosis in pattern formation. Thus this protein may also be involved in apoptosis or tissue differentiation and could again be useful in cancer therapy. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:118 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1171 of SEQ ID NO:118, b is an integer of 15 to 1185, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:118, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 109

In specific embodiments, polypeptides of the invention comprise the following amino acid sequence: DPRVRRITLDLGITLYFLYIFLSL (SEQ ID NO:444), PALGECCLDAFLFLLGKQLKKSGEKPLLGGSLMEYAILSALAAAMNEPKTCSTTA LKKYV LENHPGTNSNYQMHLKKTLQKCEKNGWMEQISGKGFSGTFQL CFPYYPSPGVLFPKKEPDDSRDEDEDE DESSEEDSEDEEPPPKRRLQKKTPAKS PGKAASVKQRGSKPAPKVSAQAQRGKARPLPKKAPPAKTPAKK TRPSSTV IKKPSGGSSKKPAT SARKEVKLPGKGKSTMKKSFRVKK (SEQ ID NO:445), DFEFHDTLFSYKIYFFTLKDFFMVDLPLPGNFTSFLALVAGFF EEPPLGFLM TVDEGLVFLAGVLALGGAFLGKGLAFPRWAAETLGAGLDPLCFTDAAFPGDLA GVFFCNLL LGGGSSSSSESSSSDDSSSSSSSSLESSGSFFGNRTPGLG (SEQ ID NO:446), CLDAFLFLLGKQLKKSGEKPLLGGSLME (SEQ ID NO:447), YQMHLK KTLQKCEKNGWMEQISGKGFSGT (SEQ ID NO:448), KTPAKSPGKAAS VKQRGSKPAPKVSAQA (SEQ ID NO:449), SSKPAT SARKEVKLPGKGKSTM KKSFR (SEQ ID NO:450), VDEGLVFLAGVLALGGAFLGKGL (SEQ ID NO:451), and/or GLDPLCFTDAAFPGDLAGVFFCNLL (SEQ ID NO:452). Polynucleotides encoding these polypeptides are also encompassed by the invention.

This gene is expressed primarily in bone marrow stromal cells, and to a lesser extent, in human osteoblasts and T cells (helper I).

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, connective tissues, haemopoietic, or immune disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the skeletal and immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g. hematopoietic, immune, skeletal, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:244 as residues: Glu-18 to Cys-38.

The tissue distribution in bone marrow stromal cells and T-cells suggest that polynucleotides and polypeptides corresponding to this gene are useful for diagnosis and treatment of defects of stromal development, and immune system disorders. Moreover, polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of hematopoietic related disorders such as anemia, pancytopenia, leukopenia, thrombocytopenia or leukemia since stromal cells are important in the production of cells of hematopoietic lineages. The uses include bone marrow cell ex vivo culture, bone marrow transplantation, bone marrow reconstitution, radiotherapy or chemotherapy of neoplasia. The gene product may also be involved in lymphopoiesis, therefore, it can be used in immune disorders such as infection, inflammation, allergy, immunodeficiency etc. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Moreover, the expression of this gene product in osteoblasts would suggest a role in the detection and treatment of disorders and conditions affecting the skeletal system, in particular osteoporosis, bone cancer, as well as, disorders afflicting connective tissues (e.g. arthritis, trauma, tendonitis, chondromalacia and inflammation), such as in the diagnosis or treatment of various autoimmune disorders such as rheumatoid arthritis, lupus, scleroderma, and dermatomyositis as well as dwarfism, spinal deformation, and specific joint abnormalities as well as chondrodysplasias (ie. spondyloepiphyseal dysplasia congenita, familial osteoarthritis, Atelosteogenesis type II, metaphyseal

chondrodysplasia type Schmid). Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:119 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1084 of SEQ ID NO:119, b is an integer of 15 to 1098, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:119, and where b is greater than or equal to a + 14.

15

FEATURES OF PROTEIN ENCODED BY GENE NO: 110

This gene is expressed primarily in rhabdomyosarcoma, CD34 positive cells, breast lymph nodes, neutrophils and endothelial cells.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune or hematopoietic disorders, developmental, proliferative, and vascular disorders, particularly fibroids or atherosclerosis. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune or hematopoietic systems, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. immune, hematopoietic, developmental, vascular, endothelial, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, amniotic fluid, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in neutrophils and lymph nodes indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and intervention of disorders in immune or hematopoietic systems. Similarly,

the secreted protein can also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, and as nutritional supplements. It may also have a very wide range of biological activities. Typical of these are cytokine, cell proliferation/differentiation modulating activity or induction of other cytokines; immunostimulating/immunosuppressant activities (e.g. for treating human immunodeficiency virus infection, cancer, autoimmune diseases and allergy); regulation of hematopoiesis (e.g. for treating anaemia or as adjunct to chemotherapy); stimulation or growth of bone, cartilage, tendons, ligaments and/or nerves (e.g. for treating wounds, stimulation of follicle stimulating hormone (for control of fertility); chemotactic and chemokinetic activities (e.g. for treating infections, tumors); hemostatic or thrombolytic activity (e.g. for treating haemophilia, cardiac infarction etc.); anti-inflammatory activity (e.g. for treating septic shock, Crohn's disease); as antimicrobials; for treating psoriasis or other hyperproliferative diseases; for regulation of metabolism, and behaviour. Also contemplated is the use of the corresponding nucleic acid in gene therapy procedures. The protein may also show utility in the treatment or prevention of a variety of vascular disorders, particularly embolism, thrombosis, aneurysms, stroke, or atherosclerosis. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:120 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 791 of SEQ ID NO:120, b is an integer of 15 to 805, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:120, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 111

In specific embodiments, polypeptides of the invention comprise the following amino acid sequence: TMLFYLSQPDWQLDFFRVSFNG PVFFIIIFNDRAGFRM QALVSQAACRRSRYKLSVVY (SEQ ID NO:453), and/or DRAGFRMQALVS

QAACRRSRYKL (SEQ ID NO:454). Polynucleotides encoding these polypeptides are also encompassed by the invention. The gene encoding the disclosed cDNA is believed to reside on chromosome 1. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 1.

5 This gene is expressed primarily in human cerebellum, and to a lesser extent, in colon carcinoma cells, activated T-cells, fetal spleen, and placenta.

 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are
10 not limited to, immune, hematopoietic, or neural disorders, particularly neurodegenerative disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune or central nervous systems, expression of this gene at
15 significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. neural, immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily
20 fluid from an individual not having the disorder.

 The tissue distribution in human cerebellum indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of diseases in the central nervous system and immune disorders. Moreover, polynucleotides and polypeptides corresponding to this gene are useful for the
25 detection/treatment of neurodegenerative disease states, behavioural disorders, or inflammatory conditions such as Alzheimers Disease, Parkinsons Disease, Huntingtons Disease, Tourette Syndrome, meningitis, encephalitis, demyelinating diseases, peripheral neuropathies, neoplasia, trauma, congenital malformations, spinal cord injuries, ischemia and infarction, aneurysms, hemorrhages, schizophrenia, mania,
30 dementia, paranoia, obsessive compulsive disorder, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and preception. In addition, elevated expression of this gene product in regions of the brain indicates that it plays a role in normal neural function. Potentially, this gene product is involved in synapse formation,
35 neurotransmission, learning, cognition, homeostasis, or neuronal differentiation or survival. Moreover, the gene or gene product may also play a role in the treatment and/or detection of developmental disorders associated with the developing embryo,

sexually-linked disorders, or disorders of the cardiovascular system. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:121 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1584 of SEQ ID NO:121, b is an integer of 15 to 1598, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:121, and where b is greater than or equal to a + 14.

15

FEATURES OF PROTEIN ENCODED BY GENE NO: 112

The gene encoding the disclosed cDNA is thought to reside on chromosome 8. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 8.

This gene is expressed primarily in testes, fetal brain, and fetal liver.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, brain and liver diseases, reproductive disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the liver and brain expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g. immune, reproductive, hepatic, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, bile, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in brain and liver tissues indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment of neural, hepatic, or metabolic diseases. Furthermore, the tissue distribution indicates that

polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and/or treatment of disorders of the brain and nervous system. Such involvement may impact many processes, such as learning and cognition. It may also be useful in the treatment of such neurodegenerative disorders as schizophrenia; ALS; or Alzheimer's. The tissue distribution further indicates that polynucleotides and polypeptides corresponding to this gene are useful for the detection and treatment of liver disorders and cancers (e.g. hepatoblastoma, jaundice, hepatitis, liver metabolic diseases and conditions that are attributable to the differentiation of hepatocyte progenitor cells). Additionally, the tissue distribution indicates that the protein product of this gene is useful for the treatment and diagnosis of conditions concerning proper testicular function (e.g. endocrine function, sperm maturation), as well as cancer. Therefore, this gene product is useful in the treatment of male infertility and/or impotence. This gene product is also useful in assays designed to identify binding agents as such agents (antagonists) are useful as male contraceptive agents. Similarly, the protein is believed to be useful in the treatment and/or diagnosis of testicular cancer. The testes are also a site of active gene expression of transcripts that may be expressed, particularly at low levels, in other tissues of the body. Therefore, this gene product may be expressed in other specific tissues or organs where it may play related functional roles in other processes, such as hematopoiesis, inflammation, bone formation, and kidney function, to name a few possible target indications. Protein, as well as, antibodies directed against the protein may show utility as a tissue-specific marker and/or immunotherapy target for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:122 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1006 of SEQ ID NO:122, b is an integer of 15 to 1020, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:122, and where b is greater than or equal to a + 14.

35 FEATURES OF PROTEIN ENCODED BY GENE NO: 113

neurotransmission, learning, cognition, homeostasis, or neuronal differentiation or survival. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available
5 and accessible through sequence databases. Some of these sequences are related to SEQ
ID NO:123 and may have been publicly available prior to conception of the present
invention. Preferably, such related polynucleotides are specifically excluded from the
scope of the present invention. To list every related sequence is cumbersome.
Accordingly, preferably excluded from the present invention are one or more
10 polynucleotides comprising a nucleotide sequence described by the general formula of
a-b, where a is any integer between 1 to 1364 of SEQ ID NO:123, b is an integer of 15
to 1378, where both a and b correspond to the positions of nucleotide residues shown
in SEQ ID NO:123, and where b is greater than or equal to a + 14.

15

FEATURES OF PROTEIN ENCODED BY GENE NO: 114

This gene is expressed primarily in neutrophils.

Therefore, polynucleotides and polypeptides of the invention are useful as
20 reagents for differential identification of the tissue(s) or cell type(s) present in a
biological sample and for diagnosis of diseases and conditions which include, but are
not limited to, immune or hematopoietic disorders, particularly inflammatory conditions
or immunodeficiencies. Similarly, polypeptides and antibodies directed to these
polypeptides are useful in providing immunological probes for differential identification
25 of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells,
particularly of the immune system, expression of this gene at significantly higher or
lower levels may be routinely detected in certain tissues or cell types (e.g. immune, and
cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine,
synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual
30 having such a disorder, relative to the standard gene expression level, i.e., the
expression level in healthy tissue or bodily fluid from an individual not having the
disorder.

The tissue distribution in neutrophils indicates that polynucleotides and
polypeptides corresponding to this gene are useful for the diagnosis and treatment of a
35 malfunctioning immune system response to foreign antigens. Furthermore, this gene
product may be involved in the regulation of cytokine production, antigen presentation,
or other processes that may also suggest a usefulness in the treatment of cancer (e.g. by

boosting immune responses). Since the gene is expressed in cells of lymphoid origin, the gene or protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Expression of this gene product in neutrophils also strongly indicates a role for this protein in immune function and immune surveillance.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:124 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome.

Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1132 of SEQ ID NO:124, b is an integer of 15 to 1146, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:124, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 115

In specific embodiments, polypeptides of the invention comprise the following amino acid sequence: LAAGILNSSLPALYHSVEEISQ (SEQ ID NO:455), XYRMNTKFLESYKMSTTLSRRHQNVSLCKDKMTPAGTDTKIAFLE (SEQ ID NO:456), SYKMSTTLSRRHQNVSLCKDM (SEQ ID NO:457), ICIESLMLHYIALVFEMAFMFPLVYHEMGSDSIRFHLQCQVDSCLPSMMRFFFSFPFL (SEQ ID NO:458), YIALVFEMAFMFPLVYHEMG (SEQ ID NO:459), and/or SDSIRFHLQCQVDSCLPSMMRF (SEQ ID NO:460). Polynucleotides encoding these polypeptides are also encompassed by the invention.

This gene is expressed primarily in melanocytes, merkel cells, synovial cells, ulcerative colitis, and to a lesser extent, in fetal spleen, bone marrow, jurkat cells, adrenal gland tumor rejected kidney from a failed transplantation.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, integumentary, skeletal, or gastrointestinal disorders, particularly tumors, including melanoma, lymphoma, and adrenal gland tumors. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a

number of disorders of the above tissues or cells, particularly of the integumentary system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g. Integumentary, skeletal, gastrointestinal, immune, hematopoietic, renal, endocrine, and cancerous and wounded tissues) or
5 bodily fluids (e.g. lymph, serum, amniotic fluid, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in melanocytes indicates that polynucleotides and
10 polypeptides corresponding to this gene are useful for detecting and treating tumors particularly those involving melanocytes, lymphocytes and the adrenal gland. The secreted protein can also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions and as nutritional supplements. It may also have a very wide range of
15 biological activities. Typical of these are cytokine, cell proliferation/differentiation modulating activity or induction of other cytokines; immunostimulating/immunosuppressant activities (e.g. for treating human immunodeficiency virus infection, cancer, autoimmune diseases and allergy); regulation of hematopoiesis (e.g. for treating anaemia or as adjunct to chemotherapy); stimulation
20 or growth of bone, cartilage, tendons, ligaments and/or nerves (e.g. for treating wounds, stimulation of follicle stimulating hormone (for control of fertility); chemotactic and chemokinetic activities (e.g. for treating infections, tumors); hemostatic or thrombolytic activity (e.g. for treating haemophilia, cardiac infarction etc.); anti-inflammatory activity (e.g. for treating septic shock, Crohn's disease); as
25 antimicrobials; for treating psoriasis or other hyperproliferative diseases; for regulation of metabolism, and behaviour. Also contemplated is the use of the corresponding nucleic acid in gene therapy procedures. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

30 Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:125 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome.
35 Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1661 of SEQ ID NO:125, b is an integer of 15

to 1675, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:125, and where b is greater than or equal to a + 14.

5 FEATURES OF PROTEIN ENCODED BY GENE NO: 116

When tested against fibroblast cell lines, supernatants removed from cells containing this gene activated the EGR1 (early growth response gene 1) promoter element. Thus, it is likely that this gene activates fibroblast cells through the EGR1
 10 signal transduction pathway. EGR1 is a separate signal transduction pathway from Jak-STAT, genes containing the EGR1 promoter are induced in various tissues and cell types upon activation, leading the cells to undergo differentiation and proliferation. In specific embodiments, polypeptides of the invention comprise the following amino acid sequence:

15 GGVSVDGSLREETDVGEGRPRGGQSEGARVTRRPSPDSNASAFDLDLDFS
 PFCIWCYRLETPAEVVF SPAPLRLSGPGLAPVVFVSTLPSLQPSSFCGWD
 LPARPRGLSGFR (SEQ ID NO:461), FTNKSCSKMSSTHLYKGSDVLCYARS
 SESMSLSCGDVANAGR LTPRLHLARSASQGPPTLPRVPPRGSRPPTA GESPA
 PRTXSLENHKNIDHLSSNSHGKFRIYGQNDIKI (SEQ ID NO:462), QDVIYTFVQ
 20 RFRRPMLCTILRKYEPVVRGRRKRWQA HPSSAFGKKRLPRPPHPAQGAPQRE
 QASHSWREPGPQNTFFPRKP (SEQ ID NO:463), REETDVGEGRPRGGQSEGA
 RV (SEQ ID NO:464), GPGLAPVVFVSTLPSLQPSSFCGWDLP (SEQ ID NO:465),
 MSSTHLYKGSDVLCYARSSMSL (SEQ ID NO:466), SQGPPTLPRVPPRG
 SRPPTAGESPAPRT (SEQ ID NO:467), RFRRPMLCTILRKYEPVVRGRRKRW
 25 (SEQ ID NO:468), and/or RLPRPPHPAQGAPQREQASHSWRE (SEQ ID NO:469).
 Polynucleotides encoding these polypeptides are also encompassed by the invention.

This gene is expressed primarily in hematopoietic cells, endothelial cells, and in spleen.

Therefore, polynucleotides and polypeptides of the invention are useful as
 30 reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, hematopoietic, integumentary, and immune disorders, particularly multiple myeloma, immunodeficiencies, leukemias, and vascular conditions. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing
 35 immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the hematopoietic, immune, and vascular systems, expression of this gene at significantly higher or lower

levels may be routinely detected in certain tissues and cell types (e.g. Immune, hematopoietic, integumentary, endothelial, and cancerous and wounded tissues) or bodily fluids (e.g.lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in spleen and hematopoietic cells, combined with the detected EGR1 biological activity indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and/or detection of vascular, immune and/or hematopoietic disorders including arthritis, ischemia, auto-immune diseases, host-graft rejection, AIDS, leukemia and microbial infection. Moreover, polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of hematopoietic related disorders such as anemia, pancytopenia, leukopenia, thrombocytopenia or leukemia since stromal cells are important in the production of cells of hematopoietic lineages. The uses include bone marrow cell ex vivo culture, bone marrow transplantation, bone marrow reconstitution, radiotherapy or chemotherapy of neoplasia. The gene product may also be involved in lymphopoiesis, therefore, it can be used in immune disorders such as infection, inflammation, allergy, immunodeficiency etc. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Furthermore, a utility for treating or preventing vascular or integumentary disorders may be anticipated for this gene based upon its expression within endothelial tissues in addition to its EGR1 activity. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:126 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1050 of SEQ ID NO:126, b is an integer of 15 to 1064, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:126, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 117

In specific embodiments, polypeptides of the invention comprise the following amino acid sequence:

5 RGMGRWL VSSGA AFPIPLNGFCESREFFPDSGS VLLHWRPNXVLIEIKVFGS
RSQSLISSK NLKTS LTFIYGKVEEV LNN (SEQ ID NO:470), LKLSSADSQA
IMNIFSADCM PRLHIALQTEMIPNRAPQGGAAANLWHEAQYRRLPFSR APEX
TDAHQASAQRGAAQLPREQ (SEQ ID NO:471, PIPLNGFCESREFFPDSGS
10 VLLHWRPNX (SEQ ID NO:472), and/or NIFSADCM PRLHIALQTEMIP NRA
PQGGA (SEQ ID NO:473). Polynucleotides encoding these polypeptides are also encompassed by the invention.

This gene is expressed primarily in neutrophils.

Therefore, polynucleotides and polypeptides of the invention are useful as

15 reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, diseases of the immune system, including neutropenia, cancer, inflammatory diseases and allergies. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential

20 identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g. Immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another

25 tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in neutrophils indicates that polynucleotides and polypeptides corresponding to this gene are useful for treatment/diagnosis of diseases

30 of the immune system since expression is primarily in neutrophils, and may be useful as a growth factor for the differentiation or proliferation of neutrophils for the treatment of neutropenia following chemotherapy or may be useful in the treatment of immune dysfunction or anti-inflammatory by inhibiting infiltration of neutrophils to the site of injury or distress. Protein, as well as, antibodies directed against the protein may show

35 utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ

ID NO:127 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome.

- Accordingly, preferably excluded from the present invention are one or more
- 5 polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1593 of SEQ ID NO:127, b is an integer of 15 to 1607, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:127, and where b is greater than or equal to a + 14.

10

FEATURES OF PROTEIN ENCODED BY GENE NO: 118

- Contact of cells with supernatant expressing the product of this gene has been shown to increase the permeability of the plasma membrane of renal mesangial cells to
- 15 calcium. Thus it is likely that the product of this gene is involved in a signal transduction pathway that is initiated when the product binds a receptor on the surface of the plasma membrane of both mesangial cells and other cell types, in addition to other cell-lines or tissue cell types. Thus, polynucleotides and polypeptides have uses which include, but are not limited to, activating mesangial cells. Binding of a ligand to a
- 20 receptor is known to alter intracellular levels of small molecules, such as calcium, potassium and sodium, as well as alter pH and membrane potential. Alterations in small molecule concentration can be measured to identify supernatants which bind to receptors of a particular cell. In addition, when tested against fibroblast cell lines, supernatants removed from cells containing this gene activated the EGR1 (early growth
- 25 response gene 1) promoter element. Thus, it is likely that this gene activates fibroblast cells through the EGR1 signal transduction pathway. EGR1 is a separate signal transduction pathway from Jak-STAT, genes containing the EGR1 promoter are induced in various tissues and cell types upon activation, leading the cells to undergo differentiation and proliferation. In specific embodiments, polypeptides of the invention
- 30 comprise the following amino acid sequence:
TFRLVSAHLKTRKLINPEAAERRWRDWSRQGWLSVK (SEQ ID NO:474),
and/or KTRKLINPEAAERRWRDWSR (SEQ ID NO:475). Polynucleotides encoding these polypeptides are also encompassed by the invention.

- This gene is expressed primarily in bone marrow cell lines, and to a lesser
- 35 extent, in human endometrial stromal cells, human adult small intestine and human pancreas tumor.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, haemopoietic and gastrointestinal tract disorders and stomatosis, in addition to endothelial, mucosal, or epithelial cell disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune and digestive systems, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g. haemopoietic, immune, reproductive, gastrointestinal, endocrine, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:253 as residues: Gly-25 to Arg-31, Ile-47 to Glu-57, Glu-120 to Arg-138.

The tissue distribution in bone marrow cells, combined with the detected calcium flux and EGR1 biological activity indicates that polynucleotides and polypeptides corresponding to this gene are useful for immune and gastrointestinal tract disorders, and stomatosis, particularly tumors and proliferative disorders. More specifically, polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of hematopoietic related disorders such as anemia, pancytopenia, leukopenia, thrombocytopenia or leukemia since stromal cells are important in the production of cells of hematopoietic lineages. The uses include bone marrow cell ex vivo culture, bone marrow transplantation, bone marrow reconstitution, radiotherapy or chemotherapy of neoplasia. The gene product may also be involved in lymphopoiesis, therefore, it can be used in immune disorders such as infection, inflammation, allergy, immunodeficiency etc. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:128 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the

scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1023 of SEQ ID NO:128, b is an integer of 15 to 1037, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:128, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 119

10

In specific embodiments, polypeptides of the invention comprise the following amino acid sequence: WNYTVNNLYLFSFSIVSMKFMHVLSINIF FGRARWLT PVIPALLEAEAGGSLGQEFKTSLGKDGETPSLLKIQKLAGHGGRRL (SEQ ID NO:476, DQPGKHGETLSLLKMQKLTWCGGMPFVIP SYSRSPRPENRLNL GDRGCTELLHSSLGNRVRLSKKKEVYMMELYSK (SEQ ID NO:477), VIPALLE AEAGGSLGQEFKTSLGKDGET (SEQ ID NO:478), and/or NRLNLGDRGCT ELLHSSLGNRVRLSKKKE (SEQ ID NO:479). Polynucleotides encoding these polypeptides are also encompassed by the invention.

This gene is expressed primarily in human fetal brain.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, neurological, developmental, and immunological disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the central nervous and immune systems, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g. neural, developmental, immune, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, amniotic fluid, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in fetal brain indicates that polynucleotides and polypeptides corresponding to this gene are useful for diagnosis and treatment of disorders relating to CNS and immune system. In addition, polynucleotides and polypeptides corresponding to this gene are useful for the detection/treatment of

neurodegenerative disease states, behavioural disorders, or inflammatory conditions such as Alzheimers Disease, Parkinsons Disease, Huntingtons Disease, Tourette Syndrome, meningitis, encephalitis, demyelinating diseases, peripheral neuropathies, neoplasia, trauma, congenital malformations, spinal cord injuries, ischemia and infarction, aneurysms, hemorrhages, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, panic disorder, learning disabilities, ALS, psychoses, autism, and altered bahaviors, including disorders in feeding, sleep patterns, balance, and preception. In addition, elevated expression of this gene product in regions of the brain indicates that it plays a role in normal neural function. Potentially, this gene product is involved in synapse formation, neurotransmission, learning, cognition, homeostasis, or neuronal differentiation or survival. Moreover, the gene or gene product may also play a role in the treatment and/or detection of developmental disorders associated with the developing embryo, sexually-linked disorders, or disorders of the cardiovascular system. Furthermore, expression within fetal tissue indicates that this protein may play a role in the regulation of cellular division, and may show utility in the diagnosis and treatment of cancer and other proliferative disorders. Similarly, developmental tissues rely on decisions involving cell differentiation and/or apoptosis in pattern formation. Thus this protein may also be involved in apoptosis or tissue differentiation and could again be useful in cancer therapy. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:129 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1132 of SEQ ID NO:129, b is an integer of 15 to 1146, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:129, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 120

In specific embodiments, polypeptides of the invention comprise the following amino acid sequence: HASEHLAALPVNVKIGK (SEQ ID NO:480). Polynucleotides

encoding these polypeptides are also encompassed by the invention. The gene encoding the disclosed cDNA is believed to reside on chromosome 5. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 5.

5 This gene is expressed primarily in T cells/helper I.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune, or haemopoitic disorders. Similarly, polypeptides and
10 antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g. Immune, haemopoitic disorders, and cancerous and wounded tissues)
15 or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID
20 NO:255 as residues: Ile-31 to Glu-36, Leu-59 to Glu-73, Ser-109 to Ser-121, Ser-175 to Gln-182, Lys-258 to Lys-264.

The tissue distribution in T-cells indicates that polynucleotides and polypeptides corresponding to this gene are useful for diagnosis and treatment of immune disorders. Moreover, expression of this gene product indicates a role in regulating the
25 proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. This gene product may be involved in the regulation of cytokine production, antigen presentation, or other processes that may also suggest a usefulness in the treatment of cancer (e.g. by boosting immune responses). Since the gene is expressed in cells of lymphoid origin, the natural gene product may be involved
30 in immune functions. Therefore it may be also used as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as
35 host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's disease,

scleroderma and tissues. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:130 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1158 of SEQ ID NO:130, b is an integer of 15 to 1172, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:130, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 121

In specific embodiments, polypeptides of the invention comprise the following amino acid sequence:
LVCILLVHWIPPLGAWGLSLMLFLILEQRCGKGKWRNALLSVSFSVPQLQMOK
VS LDSTPLNVNHDKMDIWKLTPKL (SEQ ID NO:481), IMIKWIFGNLLL SCD
LGCISTSGLPQYQGLRLLNFEYSLGFMLRSLWSRSAIQCFPS (SEQ ID NO:482),
LLLSCDLGCISTSGLPQYQGL (SEQ ID NO:483), and/or LRLLNFEYSLGFML
LRSLWSRS (SEQ ID NO:484). Polynucleotides encoding these polypeptides are also encompassed by the invention. The gene encoding the disclosed cDNA is believed to reside on chromosome 11. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 11.

This gene is expressed primarily in human gall bladder.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, metabolic, or gastrointestinal disorders, particularly those relating to the gall bladder. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of

the gastrointestinal tract system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g. Metabolic, gastrointestinal, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, bile, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:256 as residues: Ser-18 to Gly-26.

The tissue distribution in gall bladder tissue indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of gall bladder disorders, or related metabolic conditions, such as gall stones. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:131 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome.

Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 649 of SEQ ID NO:131, b is an integer of 15 to 663, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:131, and where b is greater than or equal to a + 14.

25

FEATURES OF PROTEIN ENCODED BY GENE NO: 122

In specific embodiments, polypeptides of the invention comprise the following amino acid sequence: ASPHL FIEKWGRAFILRKLLLVPVISKRIINIMAHQVKPPI FCAMIMCNLFCSGYEHLFTLMRFFSFEQIFDEV VFH (SEQ ID NO:485), and/or KLLLVPVISKRIINIMAH QVK PPIF (SEQ ID NO:486). Polynucleotides encoding these polypeptides are also encompassed by the invention. The gene encoding the disclosed cDNA is believed to reside on chromosome 4. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 4.

This gene is expressed primarily in glioblastoma.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, neural disorders, particularly glioblastoma multiform. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the central nervous system (CNS), expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g. neural, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:257 as residues: Ser-40 to Gly-45, Leu-73 to Arg-80.

The tissue distribution in glioblastoma indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of neural cell disorders. Moreover, polynucleotides and polypeptides corresponding to this gene are useful for the detection/treatment of neurodegenerative disease states, behavioural disorders, or inflammatory conditions such as Alzheimers Disease, Parkinsons Disease, Huntingtons Disease, Tourette Syndrome, meningitis, encephalitis, demyelinating diseases, peripheral neuropathies, neoplasia, trauma, congenital malformations, spinal cord injuries, ischemia and infarction, aneurysms, hemorrhages, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and perception. In addition, elevated expression of this gene product in regions of the brain indicates that it plays a role in normal neural function. Potentially, this gene product is involved in synapse formation, neurotransmission, learning, cognition, homeostasis, or neuronal differentiation or survival. Moreover, the gene or gene product may also play a role in the treatment and/or detection of developmental disorders associated with the developing embryo, sexually-linked disorders, or disorders of the cardiovascular system. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:132 and may have been publicly available prior to conception of the present

invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome.

Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of
 5 a-b, where a is any integer between 1 to 762 of SEQ ID NO:132, b is an integer of 15 to 776, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:132, and where b is greater than or equal to a + 14.

10 FEATURES OF PROTEIN ENCODED BY GENE NO: 123

When tested against U937 and Jurket cell lines, supernatants removed from cells containing this gene activated the GAS (gamma activating sequence) promoter element. Thus, it is likely that this gene activates myeloid cells, including their
 15 progenitors, through the Jak-STAT signal transduction pathway. GAS is a promoter element found upstream of many genes which are involved in the Jak-STAT pathway. The Jak-STAT pathway is a large, signal transduction pathway involved in the differentiation and proliferation of cells. Therefore, activation of the Jak-STAT pathway, reflected by the binding of the GAS element, can be used to indicate proteins
 20 involved in the proliferation and differentiation of cells. In specific embodiments, polypeptides of the invention comprise the following amino acid sequence:
 FAVIRFESIIHEFDPWFNRYRSTHHLASHGFIYFLNWFDERAWYPLGRIVGGTVY
 PGLMITAGLIHWILNT LNITVHIRDVCVFLAPTFSGLTSISTFLLTRELWN QGA
 GLLAACFIAIVPGYISRSVAGSFDNEGIAIFA LQFTYYLWVKSVKTGSVFW
 25 TMCCCLSYFYMVSAWGGYVFIIINLIPLHVFVLLLMQRYSKRVYIAYSTFYI VGL
 ILSMQIPFVGFPQIRTSEHMAAAGVFALLQAYAFLLQYLRDRLTKQEFQTLFFLG
 SLAAGAVFLSVI YLTYTGFIAPWWSGRFYSLWDTGYAKIHIPIIASVSEHQ PTT
 WVSFFFDLHILVCTFPAGLWFCIKNINDE RXFGKXGF (SEQ ID NO:487), EFD
 PWFNRYRSTHHLASHGFIYFLNWFD (SEQ ID NO:488), TRELWNQGAGLL
 30 AACFIAIVPGY (SEQ ID NO:489), TYYLWVKSVKTGSVFWTMCCCL (SEQ ID
 NO:490), GVFALLQAYAFLLQYLRDRLTKQEFQ (SEQ ID NO:491), and/or YSLWD
 TGYAKIHIPIIASVSEHQPTTW (SEQ ID NO:492). Polynucleotides encoding these polypeptides are also encompassed by the invention.

This gene is expressed primarily in human colon carcinoma (HCC) cell line, and
 35 to a lesser extent, in human eosinophils.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a

biological sample and for diagnosis of diseases and conditions which include, but are not limited to, gastrointestinal or immune disorders, particularly colon carcinoma and leukemia. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the excretory and immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g. gastrointestinal, immune, hematopoietic, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO:258 as residues: Glu-49 to Ser-54.

The tissue distribution in human colon carcinoma cell lines, combined with the detected GAS biological activity, indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of colon cancer and immune disorders. In addition, expression within cellular sources marked by proliferating cells indicates that this protein may play a role in the regulation of cellular division, and may show utility in the diagnosis and treatment of cancer and other proliferative disorders. Similarly, developmental tissues rely on decisions involving cell differentiation and/or apoptosis in pattern formation. Thus, this protein may also be involved in apoptosis or tissue differentiation and could again be useful in cancer therapy, particularly in modulating the immune response to cancer-specific antigens. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:133 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1529 of SEQ ID NO:133, b is an integer of 15 to 1543, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:133, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 124

- 5 This gene shares homology with elongation factor 1-Alpha (*giardia intestinalis*), which promotes the GTP-dependent binding of aminoacyl tRNA to the A-site of ribosomes during protein biosynthesis. One embodiment of this gene comprises polypeptides of the following amino acid sequence:
- MGHMLYLLGNINKRTMHKYXQESKKAGKASFAY AWVLDETGEERER
 10 GVTMDVGMTKFETTTKVIT LMDAPGHKDFIPNMITGAAQADVAVLVVDASR
 GEFEAGFETGGQTREHGLLVRS LGVTQLAVAVN KMDQVNWQQERF QEIT
 GKLGHFLKQAGFKESDV GFIPTSGLSGENLITRSQSSELTKWYKGLCLLEQ
 IDSFKPPQRSIDKPFRLCVSDVFKDQSGFCITG KIEAGYIQTGDRLL AMP
 NETCTVKGITLHDEPV DWAAAGDHVSLTLVGMDIINVCIFCGPKVP
 15 IKACTRFRARILIFNIEIPITKGFPVLLHYQTVSE PAVIKRLISVLNKSTG
 EVTKKKPKFLTKGQNAL VELQTRPIALELYKDFKELGRFMLRYGGSTIAA
 GVVTEIKE (SEQ ID NO:493), LYLLGNINKRTMHKYXQESKK (SEQ ID
 NO:494), LDETGEERERGVTMDVGMTKFET (SEQ ID NO:495), GHKDFIPNMIT
 GAAQADVAVLV (SEQ ID NO:496), GFETGGQTREHGLLVRS LGVTQL (SEQ ID
 20 NO:497), WQERFQEITGKLGHFLKQAGFK (SEQ ID NO:498), TSGLSGENLI
 TRSQSSELTKWY (SEQ ID NO:499), PQRSIDKPFRLCVSDVFKDQSG (SEQ ID
 NO:500), LISVLNKSTGEVTKKKPKFLTK (SEQ ID NO:501), and/or QRPIALELY
 KDFKELGRFMLRYGGS (SEQ ID NO:502). An additional embodiment is the
 polynucleotides encoding these polypeptides. The gene encoding the disclosed cDNA is
 25 thought to reside on chromosome 6. Accordingly, polynucleotides related to this
 invention are useful as a marker in linkage analysis for chromosome 6.

This gene is expressed primarily in colon tissue from a patient having ulcerative colitis, brain tissue, lung tissue, testes and endometrial tumor.

- Therefore, polynucleotides and polypeptides of the invention are useful as
 30 reagents for differential identification of the tissue(s) or cell type(s) present in a
 biological sample and for diagnosis of diseases and conditions which include, but are
 not limited to, ulcerative colitis, and testes and endometrial tumors. Similarly,
 polypeptides and antibodies directed to these polypeptides are useful in providing
 immunological probes for differential identification of the tissue(s) or cell type(s). For a
 35 number of disorders of the above tissues or cells, particularly of the immune system
 and reproductive system, expression of this gene at significantly higher or lower levels
 may be routinely detected in certain tissues or cell types (e.g. reproductive, immune,

cancerous and wounded tissues) or bodily fluids (e.g., serum, seminal fluid, pulmonary surfactant or sputum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in ulcerative colitis, testes and endometrial tumors indicates that polynucleotides and polypeptides corresponding to this gene are useful for diagnosing or treatment of a variety of reproductive or gastrointestinal disorders. Furthermore, the tissue distribution indicates that the protein product of this gene is useful for the treatment and diagnosis of conditions concerning proper testicular function (e.g. endocrine function, sperm maturation), as well as cancer. Therefore, this gene product is useful in the treatment of male infertility and/or impotence. This gene product is also useful in assays designed to identify binding agents as such agents (antagonists) are useful as male contraceptive agents. Similarly, the protein is believed to be useful in the treatment and/or diagnosis of testicular cancer. The testes are also a site of active gene expression of transcripts that may be expressed, particularly at low levels, in other tissues of the body. Therefore, this gene product may be expressed in other specific tissues or organs where it may play related functional roles in other processes, such as hematopoiesis, inflammation, bone formation, and kidney function, to name a few possible target indications.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:134 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of $a-b$, where a is any integer between 1 to 2143 of SEQ ID NO:134, b is an integer of 15 to 2157, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:134, and where b is greater than or equal to $a + 14$.

FEATURES OF PROTEIN ENCODED BY GENE NO: 125

This gene is expressed primarily in skin.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a

biological sample and for diagnosis of diseases and conditions which include, but are not limited to, integumentary diseases. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the integumentary system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g. integumentary, melanocyte, and cancerous and wounded tissues) or bodily fluids (e.g. lymph, serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in skin indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and/or treatment of diseases relating to integumentary conditions. Specifically, polynucleotides and polypeptides corresponding to this gene are useful for the treatment, diagnosis, and/or prevention of various skin disorders including congenital disorders (i.e. nevi, moles, freckles, Mongolian spots, hemangiomas, port-wine syndrome), integumentary tumors (i.e. keratoses, Bowen's disease, basal cell carcinoma, squamous cell carcinoma, malignant melanoma, Paget's disease, mycosis fungoides, and Kaposi's sarcoma), injuries and inflammation of the skin (i.e. wounds, rashes, prickly heat disorder, psoriasis, dermatitis), atherosclerosis, urticaria, eczema, photosensitivity, autoimmune disorders (i.e. lupus erythematosus, vitiligo, dermatomyositis, morphea, scleroderma, pemphigoid, and pemphigus), keloids, striae, erythema, petechiae, purpura, and xanthelasma. In addition, such disorders may predispose increased susceptibility to viral and bacterial infections of the skin (i.e. cold sores, warts, chickenpox, molluscum contagiosum, herpes zoster, boils, cellulitis, erysipelas, impetigo, tinea, athlete's foot, and ringworm). Moreover, the protein product of this gene may also be useful for the treatment or diagnosis of various connective tissue disorders such as arthritis, trauma, tendonitis, chondromalacia and inflammation, autoimmune disorders such as rheumatoid arthritis, lupus, scleroderma, and dermatomyositis as well as dwarfism, spinal deformation, and specific joint abnormalities as well as chondrodysplasias (i.e. spondyloepiphyseal dysplasia congenita, familial osteoarthritis, Atelosteogenesis type II, metaphyseal chondrodysplasia type Schmid). Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ

ID NO:135 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence is cumbersome.

- Accordingly, preferably excluded from the present invention are one or more
- 5 polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 406 of SEQ ID NO:135, b is an integer of 15 to 420, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:135, and where b is greater than or equal to a + 14.

Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
1	HUSIG64	209423 10/30/97	pSport1	11	1010	1	1010	9	9	136	1	21	22	334
2	HATCI78	209368 10/16/97	Uni-ZAP XR	12	1559	1	1559	283	283	137	1	20	21	42
3	HSIDR70	209368 10/16/97	Uni-ZAP XR	13	1589	1	1589	110	110	138	1	17	18	86
4	HFADD53	209368 10/16/97	Uni-ZAP XR	14	1255	1	1255	183	183	139	1	22	23	121
5	HPMGT51	209423 10/30/97	Uni-ZAP XR	15	1191	1	1191	152	152	140	1	29	30	275
6	HFVAB79	209368 10/16/97	Uni-ZAP XR	16	1186	1	1186	139	139	141	1	15	16	194
7	HDTBP51	209407 10/23/97	pCMVSPORT 2.0	17	1182	1	1182	93	93	142	1	25	26	182

Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of 5' NT Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
8	HLHFR19	209407 10/23/97	Uni-ZAP XR	18	1171	1	1171	24	24	143	1	30	31	121
9	HMEET96	209407 10/23/97	Lambda ZAP II	19	1337	73	1200	121	121	144	1	30	31	266
10	HTXCV12	209423 10/30/97	Uni-ZAP XR	20	1162	1	1162	183	183	145	1	27	28	91
11	HCEFB70	209423 10/30/97	Uni-ZAP XR	21	1837	1	1837	223	223	146	1	24	25	108
12	HDTAV25	209423 10/30/97	pCMVSPORT 2.0	22	1054	1	1054	100	100	147	1	38	39	87
13	HSATA21	209368 10/16/97	Uni-ZAP XR	23	1066	1	1060	49	49	148	1	25	26	73
14	HKIXI03	209368 10/16/97	pBluescript	24	928	1	928	61	61	149	1	24	25	71
15	HDTDC56	209407 10/23/97	pCMVSPORT 2.0	25	966	1	966	210	210	150	1	24	25	151

Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
16	HLTBF35	209407 10/23/97	Uni-ZAP XR	26	1146	1	1132	136	136	151	1	16	17	60
17	HEPAB80	209423 10/30/97	Uni-ZAP XR	27	802	1	802	67	67	152	1	28	29	122
18	HFOXBI3	209423 10/30/97	pSport1	28	1169	1	1169	36	36	153	1	21	22	54
19	HTOAK16	209368 10/16/97	Uni-ZAP XR	29	1466	1	1466	87	87	154	1	18	19	110
20	HBXDC63	209368 10/16/97	ZAP Express	30	1226	1	1226	165	165	155	1	30	31	47
21	HASAU43	209407 10/23/97	Uni-ZAP XR	31	1094	1	1094	33	33	156	1	17	18	81
22	HAGEA31	209423 10/30/97	Uni-ZAP XR	32	1037	1	1037	151	151	157	1	25	26	155
23	HEQAF19	209423 10/30/97	pCMVSPORT 3.0	33	1376	1	1376	84	84	158	1	23	24	294

Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
24	HTXHB33	209368 10/16/97	Uni-ZAP XR	34	1220	1	1220	243	243	159	1	17	18	59
25	HMWFT65	209368 10/16/97	Uni-Zap XR	35	1346	1	1346	72	72	160	1	28	29	121
26	HNGAZ68	209368 10/16/97	Uni-ZAP XR	36	1026	1	1026	238	238	161	1	18	19	72
27	HTWFH07	209407 10/23/97	pSport1	37	832	1	832	14	14	162	1	25	26	122
28	HMQDF12	209407 10/23/97	Uni-ZAP XR	38	706	1	627	63	63	163	1	27	28	142
29	HFABH95	209407 10/23/97	Uni-ZAP XR	39	1347	1	1347	199	199	164	1	21	22	116
30	HNGDD48	209423 10/30/97	Uni-ZAP XR	40	1467	1	1467	85	85	165	1	20	21	58
31	HPMBY46	209423 10/30/97	Uni-ZAP XR	41	914	1	914	63	63	166	1	21	22	125

Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
32	HRKPA09	209423 10/30/97	pBluescript	42	1131	1	1131	101	101	167	1	33	34	86
33	HAGAQ26	209368 10/16/97	Uni-ZAP XR	43	1333	157	1333	251	251	168	1	20	21	62
34	HCWFL55	209368 10/16/97	ZAP Express	44	1004	1	1004	40	40	169	1	19	20	47
35	HKA AE44	209368 10/16/97	pCMV Sport 2.0	45	1494	1	1494	113	113	170	1	39	40	136
36	HNGEU90	209407 10/23/97	Uni-ZAP XR	46	1166	1	1166	17	17	171	1	20	21	88
37	HCFCC07	209407 10/23/97	pSport1	47	1536	1	1536	94	94	172	1	47	48	57
38	HLWBI63	209407 10/23/97	pCMV Sport 3.0	48	1038	1	1038	149	149	173	1	30	31	63
39	HDUAC77	209423 10/30/97	pSport1	49	1176	1	1176	193	193	174	1	19	20	60

Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	5' NT 3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
40	HFOYV27	209423 10/30/97	pSport1	50	731	1	731	171	171	175	1	18	19	103
41	HGBHI35	209423 10/30/97	Uni-ZAP XR	51	1437	71	1276	87	87	176	1	16	17	292
42	HRDEU27	209423 10/30/97	Uni-ZAP XR	52	1369	1	1369	285	285	177	1	18	19	45
43	HNGJE50	209368 10/16/97	Uni-ZAP XR	53	1037	1	1037	77	77	178	1	36	37	46
44	HNH DU48	209368 10/16/97	Uni-ZAP XR	54	1373	1	1373	99	99	179	1	20	21	54
45	HFXJU68	209423 10/30/97	Lambda ZAP II	55	1347	1	1347	148	148	180	1	25	26	66
46	HMMAH60	209368 10/16/97	pSport1	56	822	1	822	142	142	181	1	15	16	50
47	HNGFR31	209407 10/23/97	Uni-ZAP XR	57	536	1	536	108	108	182	1	23	24	90

Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
48	HFPDB26	209423 10/30/97	Uni-ZAP XR	58	1262	50	1192	65	65	183	1	29	30	54
49	HFRAW86	209423 10/30/97	Uni-ZAP XR	59	1269	1	1269	162	162	184	1	16	17	63
50	HTEDX90	209368 10/16/97	Uni-ZAP XR	60	1829	1	1829	63	63	185	1	17	18	112
51	HTXGG45	209407 10/23/97	Uni-ZAP XR	61	1112	1	1112	52	52	186	1	19	20	59
52	HTXJI95	209407 10/23/97	Uni-ZAP XR	62	1674	1	1674	164	164	187	1	23	24	63
53	HLYBD32	209407 10/23/97	pSport1	63	1045	35	1045	98	98	188	1	23	24	70
54	HOUDK26	209423 10/30/97	Uni-ZAP XR	64	1051	1	1051	218	218	189	1	24	25	62
55	HROAJ03	209423 10/30/97	Uni-ZAP XR	65	1182	1	1182	19	19	190	1	20	21	192

Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
56	HTXAJ12	209423 10/30/97	Uni-ZAP XR	66	675	1	675	91	91	191	1	18	19	111
57	HKAEL80	209423 10/30/97	pCMVSPORT 2.0	67	1105	1	1105	398	398	192	1	17	18	79
58	HNHFL04	209423 10/30/97	Uni-ZAP XR	68	1279	1	1279	162	162	193	1	16	17	87
59	HPCAM01	209368 10/16/97	Uni-ZAP XR	69	1638	1	1638	311	311	194	1	24	25	41
60	HJACA79	209368 10/16/97	pBluescript SK-	70	887	1	887	84	84	195	1	28	29	68
61	HMADK33	209368 10/16/97	Uni-ZAP XR	71	864	1	864	161	161	196	1	24	25	152
62	HMSFI26	209368 10/16/97	Uni-ZAP XR	72	1217	1	1217	120	120	197	1	34	35	62
63	HMSJR08	209368 10/16/97	Uni-ZAP XR	73	1717	1	1717	165	165	198	1	28	29	63

Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	5' NT 3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
64	HMWIO93	209368 10/16/97	Uni-Zap XR	74	1276	1	1276	72	72	199	1	18	19	42
65	HNGAK47	209368 10/16/97	Uni-ZAP XR	75	1144	1	1144	89	89	200	1	23	24	40
66	HNGAL31	209368 10/16/97	Uni-ZAP XR	76	918	1	918	34	34	201	1	20	21	43
67	HNGIZ06	209368 10/16/97	Uni-ZAP XR	77	1065	1	1065	108	108	202	1	16	17	41
68	HNHBI75	209368 10/16/97	Uni-ZAP XR	78	1126	1	1126	12	12	203	1	15	16	41
69	HOFNT24	209368 10/16/97	pCMVSPORT 2.0	79	984	1	984	63	63	204	1	22	23	112
70	HSAXI95	209368 10/16/97	Uni-ZAP XR	80	1247	1	1247	147	147	205	1	19	20	44
71	HCMTB45	209368 10/16/97	Uni-ZAP XR	81	946	1	946	209	209	206	1	27	28	70

Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
72	HE9CP41	209368 10/16/97	Uni-ZAP XR	82	1392	1	1392	132	132	207	1	21	22	41
73	HHENV10	209368 10/16/97	pCMVSPORT 3.0	83	1155	1	1155	143	143	208	1	27	28	50
74	HSKDD72	209407 10/23/97	Uni-ZAP XR	84	1373	1	1373	94	94	209	1	23	24	64
75	HAGDO20	209407 10/23/97	Uni-ZAP XR	85	1258	184	1258	218	218	210	1	20	21	76
76	HCFBH15	209407 10/23/97	pSPORT	86	1318	1	1318	156	156	211	1	22	23	44
77	HSYBX48	209423 10/30/97	pCMVSPORT 3.0	87	978	38	961	246	246	212	1	34	35	65
78	HATDQ62	209423 10/30/97	Uni-ZAP XR	88	1863	323	1863	412	412	213	1	25	26	61
79	HMEIE13	209423 10/30/97	Lambda ZAP II	89	2086	1	1131	147	147	214	1	26	27	55

Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
80	HNAAF65	209423 10/30/97	pSport1	90	891	1	891	140	140	215	1	21	22	212
81	HNFHY30	209423 10/30/97	Uni-ZAP XR	91	1974	1	1974	134	134	216	1	30	31	40
82	HNFIR81	209423 10/30/97	pBluescript	92	1423	1	1423	19	19	217	1	20	21	59
83	HNTBI57	209423 10/30/97	pCMVSPORT 3.0	93	1365	134	1365	210	210	218	1	26	27	58
84	HSAYRI3	209423 10/30/97	Uni-ZAP XR	94	756	1	756	171	171	219	1	19	20	45
85	HTOHV49	209407 10/23/97	Uni-ZAP XR	95	938	1	729	62	62	220	1	19	20	61
86	HSFAG37	209368 10/16/97	Uni-ZAP XR	96	928	1	928	264	264	221	1	18	19	51
87	HTXBU52	209407 10/23/97	Uni-ZAP XR	97	1715	557	1715	574	574	222	1	34	35	50

Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
88	HLHFP18	209407 10/23/97	Uni-ZAP XR	98	678	1	678	25	25	223	1	24	25	46
89	HFXBW09	209423 10/30/97	Lambda ZAP II	99	1541	1	1541	159	159	224	1	29	30	51
90	HNGEM62	209423 10/30/97	Uni-ZAP XR	100	881	1	881	78	78	225	1	21	22	65
91	HNGJF92	209423 10/30/97	Uni-ZAP XR	101	947	1	947	40	40	226	1	31	32	46
92	HMEED18	209368 10/16/97	Lambda ZAP II	102	1369	28	1369	34	34	227	1	34	35	221
93	HMIAM45	209368 10/16/97	Uni-ZAP XR	103	1231	1	1231	68	68	228	1	37	38	48
94	HSAVK10	209368 10/16/97	Uni-ZAP XR	104	1242	1	1242	131	131	229	1	32	33	40
95	HSDHC81	209368 10/16/97	Uni-ZAP XR	105	1151	1	1151	184	184	230	1	22	23	52

Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
96	HSLCT04	209368 10/16/97	Uni-ZAP XR	106	1628	1	1628	159	159	231	1	36	37	49
97	HMDAB56	209368 10/16/97	Uni-ZAP XR	107	1465	1	1465	273	273	232	1	33	34	44
98	HUDBZ89	209407 10/23/97	ZAP Express	108	1265	1	1265	197	197	233	1	17	18	54
99	HL YCT47	209407 10/23/97	pSport1	109	1006	1	1006	47	47	234	1	22	23	68
100	HOSDJ25	209423 10/30/97	Uni-ZAP XR	110	1258	1	1258	146	146	235	1	18	19	40
101	HADAO89	209423 10/30/97	pSport1	111	1453	1	1453	244	244	236	1	22	23	44
102	HMSG B14	209423 10/30/97	Uni-ZAP XR	112	1552	1	1552	138	138	237	1	18	19	77
103	HPMGD01	209423 10/30/97	Uni-ZAP XR	113	1489	140	1489	157	157	238	1	36	37	52

Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
104	HNHFU32	209407 10/23/97	Uni-ZAP XR	114	607	1	607	175	175	239	1	30	31	52
105	HMIAL40	209368 10/16/97	Uni-ZAP XR	115	1498	1	1498	235	235	240	1	19	20	42
106	HAMFY69	209407 10/23/97	pCMVSPORT 3.0	116	1797	314	1797	359	359	241	1	17	18	48
107	HBMCT17	209407 10/23/97	pBluescript	117	952	1	952	160	160	242	1	25	26	74
108	HEBFI91	209407 10/23/97	Uni-ZAP XR	118	1185	1	1185	132	132	243	1	20	21	43
109	HHEAH86	209407 10/23/97	pCMVSPORT 3.0	119	1098	1	1098	75	75	244	1	16	17	64
110	HRDFD27	209423 10/30/97	Uni-ZAP XR	120	805	1	805	82	82	245	1	36	37	83
111	HTPCS72	209423 10/30/97	Uni-ZAP XR	121	1598	306	1598	530	530	246	1	29	30	71

Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
112	HFFAL36	209368 10/16/97	Lambda ZAP II	122	1020	1	1020	68	68	247	1	35	36	56
113	HFXBT12	209368 10/16/97	Lambda ZAP II	123	1378	1	1378	79	79	248	1	18	19	66
114	HNGJF70	209368 10/16/97	Uni-ZAP XR	124	1146	1	1146	94	94	249	1	16	17	45
115	HATEE46	209407 10/23/97	Uni-ZAP XR	125	1675	136	863	241	241	250	1	21	22	53
116	HJMBN89	209407 10/23/97	pCMVSPORT 3.0	126	1064	306	1064	348	348	251	1	13	14	56
117	HNHEK61	209407 10/23/97	Uni-ZAP XR	127	1607	1	1607	45	45	252	1	24	25	41
118	HEQAO65	209407 10/23/97	pCMVSPORT 3.0	128	1037	5	1037	152	152	253	1	27	28	160
119	HFCDV54	209407 10/23/97	Uni-ZAP XR	129	1146	1	1146	27	27	254	1	29	30	50

Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
120	HHEAD14	209407 10/23/97	pCMVSPORT 3.0	130	1172	1	1172	53	53	255	1	18	19	278
121	HGBHE57	209407 10/23/97	Uni-ZAP XR	131	663	1	663	14	14	256	1	19	20	68
122	HGLAF75	209407 10/23/97	Uni-ZAP XR	132	776	1	776	231	231	257	1	28	29	121
123	HHEMQ28	209407 10/23/97	pCMVSPORT 3.0	133	1543	286	1543	442	442	258	1	31	32	58
124	HMWEC56	209368 10/16/97	Uni-Zap XR	134	2157	1013	2146	1067	1067	259	1	17	18	67
125	HERAR44	209407 10/23/97	Uni-ZAP XR	135	420	1	420	60	60	260	1	40	41	45

Table 1 summarizes the information corresponding to each "Gene No." described above. The nucleotide sequence identified as "NT SEQ ID NO:X" was assembled from partially homologous ("overlapping") sequences obtained from the "cDNA clone ID" identified in Table 1 and, in some cases, from additional related DNA clones. The overlapping sequences were assembled into a single contiguous sequence of high redundancy (usually three to five overlapping sequences at each nucleotide position), resulting in a final sequence identified as SEQ ID NO:X.

The cDNA Clone ID was deposited on the date and given the corresponding deposit number listed in "ATCC Deposit No:Z and Date." Some of the deposits contain multiple different clones corresponding to the same gene. "Vector" refers to the type of vector contained in the cDNA Clone ID.

"Total NT Seq." refers to the total number of nucleotides in the contig identified by "Gene No." The deposited clone may contain all or most of these sequences, reflected by the nucleotide position indicated as "5' NT of Clone Seq." and the "3' NT of Clone Seq." of SEQ ID NO:X. The nucleotide position of SEQ ID NO:X of the putative start codon (methionine) is identified as "5' NT of Start Codon." Similarly, the nucleotide position of SEQ ID NO:X of the predicted signal sequence is identified as "5' NT of First AA of Signal Pep."

The translated amino acid sequence, beginning with the methionine, is identified as "AA SEQ ID NO:Y," although other reading frames can also be easily translated using known molecular biology techniques. The polypeptides produced by these alternative open reading frames are specifically contemplated by the present invention.

The first and last amino acid position of SEQ ID NO:Y of the predicted signal peptide is identified as "First AA of Sig Pep" and "Last AA of Sig Pep." The predicted first amino acid position of SEQ ID NO:Y of the secreted portion is identified as "Predicted First AA of Secreted Portion." Finally, the amino acid position of SEQ ID NO:Y of the last amino acid in the open reading frame is identified as "Last AA of ORF."

SEQ ID NO:X and the translated SEQ ID NO:Y are sufficiently accurate and otherwise suitable for a variety of uses well known in the art and described further below. For instance, SEQ ID NO:X is useful for designing nucleic acid hybridization probes that will detect nucleic acid sequences contained in SEQ ID NO:X or the cDNA contained in the deposited clone. These probes will also hybridize to nucleic acid molecules in biological samples, thereby enabling a variety of forensic and diagnostic methods of the invention. Similarly, polypeptides identified from SEQ ID NO:Y may be used to generate antibodies which bind specifically to the secreted proteins encoded by the cDNA clones identified in Table 1.

Nevertheless, DNA sequences generated by sequencing reactions can contain sequencing errors. The errors exist as misidentified nucleotides, or as insertions or deletions of nucleotides in the generated DNA sequence. The erroneously inserted or deleted nucleotides cause frame shifts in the reading frames of the predicted amino acid sequence. In these cases, the predicted amino acid sequence diverges from the actual amino acid sequence, even though the generated DNA sequence may be greater than 99.9% identical to the actual DNA sequence (for example, one base insertion or deletion in an open reading frame of over 1000 bases).

Accordingly, for those applications requiring precision in the nucleotide sequence or the amino acid sequence, the present invention provides not only the generated nucleotide sequence identified as SEQ ID NO:X and the predicted translated amino acid sequence identified as SEQ ID NO:Y, but also a sample of plasmid DNA containing a human cDNA of the invention deposited with the ATCC, as set forth in Table 1. The nucleotide sequence of each deposited clone can readily be determined by sequencing the deposited clone in accordance with known methods. The predicted amino acid sequence can then be verified from such deposits. Moreover, the amino acid sequence of the protein encoded by a particular clone can also be directly determined by peptide sequencing or by expressing the protein in a suitable host cell containing the deposited human cDNA, collecting the protein, and determining its sequence.

The present invention also relates to the genes corresponding to SEQ ID NO:X, SEQ ID NO:Y, or the deposited clone. The corresponding gene can be isolated in accordance with known methods using the sequence information disclosed herein. Such methods include preparing probes or primers from the disclosed sequence and identifying or amplifying the corresponding gene from appropriate sources of genomic material.

Also provided in the present invention are species homologs. Species homologs may be isolated and identified by making suitable probes or primers from the sequences provided herein and screening a suitable nucleic acid source for the desired homologue.

The polypeptides of the invention can be prepared in any suitable manner. Such polypeptides include isolated naturally occurring polypeptides, recombinantly produced polypeptides, synthetically produced polypeptides, or polypeptides produced by a combination of these methods. Means for preparing such polypeptides are well understood in the art.

The polypeptides may be in the form of the secreted protein, including the mature form, or may be a part of a larger protein, such as a fusion protein (see below).

It is often advantageous to include an additional amino acid sequence which contains secretory or leader sequences, pro-sequences, sequences which aid in purification, such as multiple histidine residues, or an additional sequence for stability during recombinant production.

5 The polypeptides of the present invention are preferably provided in an isolated form, and preferably are substantially purified. A recombinantly produced version of a polypeptide, including the secreted polypeptide, can be substantially purified by the one-step method described in Smith and Johnson, *Gene* 67:31-40 (1988).

10 Polypeptides of the invention also can be purified from natural or recombinant sources using antibodies of the invention raised against the secreted protein in methods which are well known in the art.

Signal Sequences

15 Methods for predicting whether a protein has a signal sequence, as well as the cleavage point for that sequence, are available. For instance, the method of McGeoch, *Virus Res.* 3:271-286 (1985), uses the information from a short N-terminal charged region and a subsequent uncharged region of the complete (uncleaved) protein. The method of von Heinje, *Nucleic Acids Res.* 14:4683-4690 (1986) uses the information from the residues surrounding the cleavage site, typically residues -13 to +2, where +1
20 indicates the amino terminus of the secreted protein. The accuracy of predicting the cleavage points of known mammalian secretory proteins for each of these methods is in the range of 75-80%. (von Heinje, *supra*.) However, the two methods do not always produce the same predicted cleavage point(s) for a given protein.

25 In the present case, the deduced amino acid sequence of the secreted polypeptide was analyzed by a computer program called SignalP (Henrik Nielsen et al., *Protein Engineering* 10:1-6 (1997)), which predicts the cellular location of a protein based on the amino acid sequence. As part of this computational prediction of localization, the methods of McGeoch and von Heinje are incorporated. The analysis of the amino acid sequences of the secreted proteins described herein by this program provided the results
30 shown in Table 1.

 As one of ordinary skill would appreciate, however, cleavage sites sometimes vary from organism to organism and cannot be predicted with absolute certainty. Accordingly, the present invention provides secreted polypeptides having a sequence shown in SEQ ID NO:Y which have an N-terminus beginning within 5 residues (i.e., +
35 or - 5 residues) of the predicted cleavage point. Similarly, it is also recognized that in some cases, cleavage of the signal sequence from a secreted protein is not entirely

uniform, resulting in more than one secreted species. These polypeptides, and the polynucleotides encoding such polypeptides, are contemplated by the present invention.

Moreover, the signal sequence identified by the above analysis may not necessarily predict the naturally occurring signal sequence. For example, the naturally occurring signal sequence may be further upstream from the predicted signal sequence. However, it is likely that the predicted signal sequence will be capable of directing the secreted protein to the ER. These polypeptides, and the polynucleotides encoding such polypeptides, are contemplated by the present invention.

10 Polynucleotide and Polypeptide Variants

"Variant" refers to a polynucleotide or polypeptide differing from the polynucleotide or polypeptide of the present invention, but retaining essential properties thereof. Generally, variants are overall closely similar, and, in many regions, identical to the polynucleotide or polypeptide of the present invention.

By a polynucleotide having a nucleotide sequence at least, for example, 95% "identical" to a reference nucleotide sequence of the present invention, it is intended that the nucleotide sequence of the polynucleotide is identical to the reference sequence except that the polynucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence encoding the polypeptide. In other words, to obtain a polynucleotide having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. The query sequence may be an entire sequence shown in Table 1, the ORF (open reading frame), or any fragment specified as described herein.

As a practical matter, whether any particular nucleic acid molecule or polypeptide is at least 90%, 95%, 96%, 97%, 98% or 99% identical to a nucleotide sequence of the present invention can be determined conventionally using known computer programs. A preferred method for determining the best overall match between a query sequence (a sequence of the present invention) and a subject sequence, also referred to as a global sequence alignment, can be determined using the FASTDB computer program based on the algorithm of Brutlag et al. (Comp. App. Biosci. (1990) 6:237-245). In a sequence alignment the query and subject sequences are both DNA sequences. An RNA sequence can be compared by converting U's to T's. The result of said global sequence alignment is in percent identity. Preferred parameters used in a FASTDB alignment of DNA sequences to calculate percent identity are: Matrix=Unitary, k-tuple=4, Mismatch Penalty=1, Joining Penalty=30, Randomization

Group Length=0, Cutoff Score=1, Gap Penalty=5, Gap Size Penalty 0.05, Window Size=500 or the length of the subject nucleotide sequence, whichever is shorter.

If the subject sequence is shorter than the query sequence because of 5' or 3' deletions, not because of internal deletions, a manual correction must be made to the results. This is because the FASTDB program does not account for 5' and 3' truncations of the subject sequence when calculating percent identity. For subject sequences truncated at the 5' or 3' ends, relative to the query sequence, the percent identity is corrected by calculating the number of bases of the query sequence that are 5' and 3' of the subject sequence, which are not matched/aligned, as a percent of the total bases of the query sequence. Whether a nucleotide is matched/aligned is determined by results of the FASTDB sequence alignment. This percentage is then subtracted from the percent identity, calculated by the above FASTDB program using the specified parameters, to arrive at a final percent identity score. This corrected score is what is used for the purposes of the present invention. Only bases outside the 5' and 3' bases of the subject sequence, as displayed by the FASTDB alignment, which are not matched/aligned with the query sequence, are calculated for the purposes of manually adjusting the percent identity score.

For example, a 90 base subject sequence is aligned to a 100 base query sequence to determine percent identity. The deletions occur at the 5' end of the subject sequence and therefore, the FASTDB alignment does not show a matched/alignment of the first 10 bases at 5' end. The 10 unpaired bases represent 10% of the sequence (number of bases at the 5' and 3' ends not matched/total number of bases in the query sequence) so 10% is subtracted from the percent identity score calculated by the FASTDB program. If the remaining 90 bases were perfectly matched the final percent identity would be 90%. In another example, a 90 base subject sequence is compared with a 100 base query sequence. This time the deletions are internal deletions so that there are no bases on the 5' or 3' of the subject sequence which are not matched/aligned with the query. In this case the percent identity calculated by FASTDB is not manually corrected. Once again, only bases 5' and 3' of the subject sequence which are not matched/aligned with the query sequence are manually corrected for. No other manual corrections are to be made for the purposes of the present invention.

By a polypeptide having an amino acid sequence at least, for example, 95% "identical" to a query amino acid sequence of the present invention, it is intended that the amino acid sequence of the subject polypeptide is identical to the query sequence except that the subject polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the query amino acid sequence. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a query

amino acid sequence, up to 5% of the amino acid residues in the subject sequence may be inserted, deleted, (indels) or substituted with another amino acid. These alterations of the reference sequence may occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions,
5 interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

As a practical matter, whether any particular polypeptide is at least 90%, 95%, 96%, 97%, 98% or 99% identical to, for instance, the amino acid sequences shown in Table 1 or to the amino acid sequence encoded by deposited DNA clone can be
10 determined conventionally using known computer programs. A preferred method for determining the best overall match between a query sequence (a sequence of the present invention) and a subject sequence, also referred to as a global sequence alignment, can be determined using the FASTDB computer program based on the algorithm of Brutlag et al. (Comp. App. Biosci. (1990) 6:237-245). In a sequence alignment the query and
15 subject sequences are either both nucleotide sequences or both amino acid sequences. The result of said global sequence alignment is in percent identity. Preferred parameters used in a FASTDB amino acid alignment are: Matrix=PAM 0, k-tuple=2, Mismatch Penalty=1, Joining Penalty=20, Randomization Group Length=0, Cutoff Score=1, Window Size=sequence length, Gap Penalty=5, Gap Size Penalty=0.05, Window
20 Size=500 or the length of the subject amino acid sequence, whichever is shorter.

If the subject sequence is shorter than the query sequence due to N- or C-terminal deletions, not because of internal deletions, a manual correction must be made to the results. This is because the FASTDB program does not account for N- and C-terminal truncations of the subject sequence when calculating global percent identity.
25 For subject sequences truncated at the N- and C-termini, relative to the the query sequence, the percent identity is corrected by calculating the number of residues of the query sequence that are N- and C-terminal of the subject sequence, which are not matched/aligned with a corresponding subject residue, as a percent of the total bases of the query sequence. Whether a residue is matched/aligned is determined by results of
30 the FASTDB sequence alignment. This percentage is then subtracted from the percent identity, calculated by the above FASTDB program using the specified parameters, to arrive at a final percent identity score. This final percent identity score is what is used for the purposes of the present invention. Only residues to the N- and C-termini of the subject sequence, which are not matched/aligned with the query sequence, are
35 considered for the purposes of manually adjusting the percent identity score. That is, only query residue positions outside the farthest N- and C-terminal residues of the subject sequence.

For example, a 90 amino acid residue subject sequence is aligned with a 100 residue query sequence to determine percent identity. The deletion occurs at the N-terminus of the subject sequence and therefore, the FASTDB alignment does not show a matching/alignment of the first 10 residues at the N-terminus. The 10 unpaired
5 residues represent 10% of the sequence (number of residues at the N- and C- termini not matched/total number of residues in the query sequence) so 10% is subtracted from the percent identity score calculated by the FASTDB program. If the remaining 90 residues were perfectly matched the final percent identity would be 90%. In another example, a 90 residue subject sequence is compared with a 100 residue query sequence.
10 This time the deletions are internal deletions so there are no residues at the N- or C-termini of the subject sequence which are not matched/aligned with the query. In this case the percent identity calculated by FASTDB is not manually corrected. Once again, only residue positions outside the N- and C-terminal ends of the subject sequence, as displayed in the FASTDB alignment, which are not matched/aligned with the query
15 sequence are manually corrected for. No other manual corrections are to made for the purposes of the present invention.

The variants may contain alterations in the coding regions, non-coding regions, or both. Especially preferred are polynucleotide variants containing alterations which produce silent substitutions, additions, or deletions, but do not alter the properties or
20 activities of the encoded polypeptide. Nucleotide variants produced by silent substitutions due to the degeneracy of the genetic code are preferred. Moreover, variants in which 5-10, 1-5, or 1-2 amino acids are substituted, deleted, or added in any combination are also preferred. Polynucleotide variants can be produced for a variety of reasons, e.g., to optimize codon expression for a particular host (change codons in
25 the human mRNA to those preferred by a bacterial host such as *E. coli*).

Naturally occurring variants are called "allelic variants," and refer to one of several alternate forms of a gene occupying a given locus on a chromosome of an organism. (Genes II, Lewin, B., ed., John Wiley & Sons, New York (1985).) These allelic variants can vary at either the polynucleotide and/or polypeptide level.
30 Alternatively, non-naturally occurring variants may be produced by mutagenesis techniques or by direct synthesis.

Using known methods of protein engineering and recombinant DNA technology, variants may be generated to improve or alter the characteristics of the polypeptides of the present invention. For instance, one or more amino acids can be
35 deleted from the N-terminus or C-terminus of the secreted protein without substantial loss of biological function. The authors of Ron et al., *J. Biol. Chem.* 268: 2984-2988 (1993), reported variant KGF proteins having heparin binding activity even after

deleting 3, 8, or 27 amino-terminal amino acid residues. Similarly, Interferon gamma exhibited up to ten times higher activity after deleting 8-10 amino acid residues from the carboxy terminus of this protein. (Dobeli et al., J. Biotechnology 7:199-216 (1988).)

Moreover, ample evidence demonstrates that variants often retain a biological activity similar to that of the naturally occurring protein. For example, Gayle and coworkers (J. Biol. Chem 268:22105-22111 (1993)) conducted extensive mutational analysis of human cytokine IL-1a. They used random mutagenesis to generate over 3,500 individual IL-1a mutants that averaged 2.5 amino acid changes per variant over the entire length of the molecule. Multiple mutations were examined at every possible amino acid position. The investigators found that "[m]ost of the molecule could be altered with little effect on either [binding or biological activity]." (See, Abstract.) In fact, only 23 unique amino acid sequences, out of more than 3,500 nucleotide sequences examined, produced a protein that significantly differed in activity from wild-type.

Furthermore, even if deleting one or more amino acids from the N-terminus or C-terminus of a polypeptide results in modification or loss of one or more biological functions, other biological activities may still be retained. For example, the ability of a deletion variant to induce and/or to bind antibodies which recognize the secreted form will likely be retained when less than the majority of the residues of the secreted form are removed from the N-terminus or C-terminus. Whether a particular polypeptide lacking N- or C-terminal residues of a protein retains such immunogenic activities can readily be determined by routine methods described herein and otherwise known in the art.

Thus, the invention further includes polypeptide variants which show substantial biological activity. Such variants include deletions, insertions, inversions, repeats, and substitutions selected according to general rules known in the art so as have little effect on activity. For example, guidance concerning how to make phenotypically silent amino acid substitutions is provided in Bowie, J. U. et al., Science 247:1306-1310 (1990), wherein the authors indicate that there are two main strategies for studying the tolerance of an amino acid sequence to change.

The first strategy exploits the tolerance of amino acid substitutions by natural selection during the process of evolution. By comparing amino acid sequences in different species, conserved amino acids can be identified. These conserved amino acids are likely important for protein function. In contrast, the amino acid positions where substitutions have been tolerated by natural selection indicates that these positions are not critical for protein function. Thus, positions tolerating amino acid substitution could be modified while still maintaining biological activity of the protein.

The second strategy uses genetic engineering to introduce amino acid changes at specific positions of a cloned gene to identify regions critical for protein function. For example, site directed mutagenesis or alanine-scanning mutagenesis (introduction of single alanine mutations at every residue in the molecule) can be used. (Cunningham and Wells, Science 244:1081-1085 (1989).) The resulting mutant molecules can then be tested for biological activity.

As the authors state, these two strategies have revealed that proteins are surprisingly tolerant of amino acid substitutions. The authors further indicate which amino acid changes are likely to be permissive at certain amino acid positions in the protein. For example, most buried (within the tertiary structure of the protein) amino acid residues require nonpolar side chains, whereas few features of surface side chains are generally conserved. Moreover, tolerated conservative amino acid substitutions involve replacement of the aliphatic or hydrophobic amino acids Ala, Val, Leu and Ile; replacement of the hydroxyl residues Ser and Thr; replacement of the acidic residues Asp and Glu; replacement of the amide residues Asn and Gln, replacement of the basic residues Lys, Arg, and His; replacement of the aromatic residues Phe, Tyr, and Trp, and replacement of the small-sized amino acids Ala, Ser, Thr, Met, and Gly.

Besides conservative amino acid substitution, variants of the present invention include (i) substitutions with one or more of the non-conserved amino acid residues, where the substituted amino acid residues may or may not be one encoded by the genetic code, or (ii) substitution with one or more of amino acid residues having a substituent group, or (iii) fusion of the mature polypeptide with another compound, such as a compound to increase the stability and/or solubility of the polypeptide (for example, polyethylene glycol), or (iv) fusion of the polypeptide with additional amino acids, such as an IgG Fc fusion region peptide, or leader or secretory sequence, or a sequence facilitating purification. Such variant polypeptides are deemed to be within the scope of those skilled in the art from the teachings herein.

For example, polypeptide variants containing amino acid substitutions of charged amino acids with other charged or neutral amino acids may produce proteins with improved characteristics, such as less aggregation. Aggregation of pharmaceutical formulations both reduces activity and increases clearance due to the aggregate's immunogenic activity. (Pinckard et al., Clin. Exp. Immunol. 2:331-340 (1967); Robbins et al., Diabetes 36: 838-845 (1987); Cleland et al., Crit. Rev. Therapeutic Drug Carrier Systems 10:307-377 (1993).)

Polynucleotide and Polypeptide Fragments

In the present invention, a "polynucleotide fragment" refers to a short polynucleotide having a nucleic acid sequence contained in the deposited clone or shown in SEQ ID NO:X. The short nucleotide fragments are preferably at least about 15 nt, and more preferably at least about 20 nt, still more preferably at least about 30 nt, and even more preferably, at least about 40 nt in length. A fragment "at least 20 nt in length," for example, is intended to include 20 or more contiguous bases from the cDNA sequence contained in the deposited clone or the nucleotide sequence shown in SEQ ID NO:X. These nucleotide fragments are useful as diagnostic probes and primers as discussed herein. Of course, larger fragments (e.g., 50, 150, 500, 600, 2000 nucleotides) are preferred.

Moreover, representative examples of polynucleotide fragments of the invention, include, for example, fragments having a sequence from about nucleotide number 1-50, 51-100, 101-150, 151-200, 201-250, 251-300, 301-350, 351-400, 401-450, 451-500, 501-550, 551-600, 651-700, 701-750, 751-800, 800-850, 851-900, 901-950, 951-1000, 1001-1050, 1051-1100, 1101-1150, 1151-1200, 1201-1250, 1251-1300, 1301-1350, 1351-1400, 1401-1450, 1451-1500, 1501-1550, 1551-1600, 1601-1650, 1651-1700, 1701-1750, 1751-1800, 1801-1850, 1851-1900, 1901-1950, 1951-2000, or 2001 to the end of SEQ ID NO:X or the cDNA contained in the deposited clone. In this context "about" includes the particularly recited ranges, larger or smaller by several (5, 4, 3, 2, or 1) nucleotides, at either terminus or at both termini. Preferably, these fragments encode a polypeptide which has biological activity. More preferably, these polynucleotides can be used as probes or primers as discussed herein.

In the present invention, a "polypeptide fragment" refers to a short amino acid sequence contained in SEQ ID NO:Y or encoded by the cDNA contained in the deposited clone. Protein fragments may be "free-standing," or comprised within a larger polypeptide of which the fragment forms a part or region, most preferably as a single continuous region. Representative examples of polypeptide fragments of the invention, include, for example, fragments from about amino acid number 1-20, 21-40, 41-60, 61-80, 81-100, 102-120, 121-140, 141-160, or 161 to the end of the coding region. Moreover, polypeptide fragments can be about 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, or 150 amino acids in length. In this context "about" includes the particularly recited ranges, larger or smaller by several (5, 4, 3, 2, or 1) amino acids, at either extreme or at both extremes.

Preferred polypeptide fragments include the secreted protein as well as the mature form. Further preferred polypeptide fragments include the secreted protein or the mature form having a continuous series of deleted residues from the amino or the carboxy terminus, or both. For example, any number of amino acids, ranging from 1-

60, can be deleted from the amino terminus of either the secreted polypeptide or the mature form. Similarly, any number of amino acids, ranging from 1-30, can be deleted from the carboxy terminus of the secreted protein or mature form. Furthermore, any combination of the above amino and carboxy terminus deletions are preferred.

5 Similarly, polynucleotide fragments encoding these polypeptide fragments are also preferred.

Also preferred are polypeptide and polynucleotide fragments characterized by structural or functional domains, such as fragments that comprise alpha-helix and alpha-helix forming regions, beta-sheet and beta-sheet-forming regions, turn and turn-
10 forming regions, coil and coil-forming regions, hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions, substrate binding region, and high antigenic index regions. Polypeptide fragments of SEQ ID NO:Y falling within conserved domains are specifically contemplated by the present invention. Moreover, polynucleotide
15 fragments encoding these domains are also contemplated.

Other preferred fragments are biologically active fragments. Biologically active fragments are those exhibiting activity similar, but not necessarily identical, to an activity of the polypeptide of the present invention. The biological activity of the fragments may include an improved desired activity, or a decreased undesirable activity.
20

Epitopes & Antibodies

In the present invention, "epitopes" refer to polypeptide fragments having antigenic or immunogenic activity in an animal, especially in a human. A preferred embodiment of the present invention relates to a polypeptide fragment comprising an
25 epitope, as well as the polynucleotide encoding this fragment. A region of a protein molecule to which an antibody can bind is defined as an "antigenic epitope." In contrast, an "immunogenic epitope" is defined as a part of a protein that elicits an antibody response. (See, for instance, Geysen et al., Proc. Natl. Acad. Sci. USA 81:3998- 4002 (1983).)

30 Fragments which function as epitopes may be produced by any conventional means. (See, e.g., Houghten, R. A., Proc. Natl. Acad. Sci. USA 82:5131-5135 (1985) further described in U.S. Patent No. 4,631,211.)

In the present invention, antigenic epitopes preferably contain a sequence of at least seven, more preferably at least nine, and most preferably between about 15 to
35 about 30 amino acids. Antigenic epitopes are useful to raise antibodies, including monoclonal antibodies, that specifically bind the epitope. (See, for instance, Wilson et al., Cell 37:767-778 (1984); Sutcliffe, J. G. et al., Science 219:660-666 (1983).)

Similarly, immunogenic epitopes can be used to induce antibodies according to methods well known in the art. (See, for instance, Sutcliffe et al., supra; Wilson et al., supra; Chow, M. et al., Proc. Natl. Acad. Sci. USA 82:910-914; and Bittle, F. J. et al., J. Gen. Virol. 66:2347-2354 (1985).) A preferred immunogenic epitope includes the secreted protein. The immunogenic epitopes may be presented together with a carrier protein, such as an albumin, to an animal system (such as rabbit or mouse) or, if it is long enough (at least about 25 amino acids), without a carrier. However, immunogenic epitopes comprising as few as 8 to 10 amino acids have been shown to be sufficient to raise antibodies capable of binding to, at the very least, linear epitopes in a denatured polypeptide (e.g., in Western blotting.)

As used herein, the term "antibody" (Ab) or "monoclonal antibody" (Mab) is meant to include intact molecules as well as antibody fragments (such as, for example, Fab and F(ab')₂ fragments) which are capable of specifically binding to protein. Fab and F(ab')₂ fragments lack the Fc fragment of intact antibody, clear more rapidly from the circulation, and may have less non-specific tissue binding than an intact antibody. (Wahl et al., J. Nucl. Med. 24:316-325 (1983).) Thus, these fragments are preferred, as well as the products of a FAB or other immunoglobulin expression library. Moreover, antibodies of the present invention include chimeric, single chain, and humanized antibodies.

Fusion Proteins

Any polypeptide of the present invention can be used to generate fusion proteins. For example, the polypeptide of the present invention, when fused to a second protein, can be used as an antigenic tag. Antibodies raised against the polypeptide of the present invention can be used to indirectly detect the second protein by binding to the polypeptide. Moreover, because secreted proteins target cellular locations based on trafficking signals, the polypeptides of the present invention can be used as targeting molecules once fused to other proteins.

Examples of domains that can be fused to polypeptides of the present invention include not only heterologous signal sequences, but also other heterologous functional regions. The fusion does not necessarily need to be direct, but may occur through linker sequences.

Moreover, fusion proteins may also be engineered to improve characteristics of the polypeptide of the present invention. For instance, a region of additional amino acids, particularly charged amino acids, may be added to the N-terminus of the polypeptide to improve stability and persistence during purification from the host cell or subsequent handling and storage. Also, peptide moieties may be added to the

polypeptide to facilitate purification. Such regions may be removed prior to final preparation of the polypeptide. The addition of peptide moieties to facilitate handling of polypeptides are familiar and routine techniques in the art.

Moreover, polypeptides of the present invention, including fragments, and
5 specifically epitopes, can be combined with parts of the constant domain of
immunoglobulins (IgG), resulting in chimeric polypeptides. These fusion proteins
facilitate purification and show an increased half-life in vivo. One reported example
describes chimeric proteins consisting of the first two domains of the human CD4-
polypeptide and various domains of the constant regions of the heavy or light chains of
10 mammalian immunoglobulins. (EP A 394,827; Traunecker et al., *Nature* 331:84-86
(1988).) Fusion proteins having disulfide-linked dimeric structures (due to the IgG)
can also be more efficient in binding and neutralizing other molecules, than the
monomeric secreted protein or protein fragment alone. (Fountoulakis et al., *J.*
Biochem. 270:3958-3964 (1995).)

15 Similarly, EP-A-O 464 533 (Canadian counterpart 2045869) discloses fusion
proteins comprising various portions of constant region of immunoglobulin molecules
together with another human protein or part thereof. In many cases, the Fc part in a
fusion protein is beneficial in therapy and diagnosis, and thus can result in, for
example, improved pharmacokinetic properties. (EP-A 0232 262.) Alternatively,
20 deleting the Fc part after the fusion protein has been expressed, detected, and purified,
would be desired. For example, the Fc portion may hinder therapy and diagnosis if the
fusion protein is used as an antigen for immunizations. In drug discovery, for
example, human proteins, such as hIL-5, have been fused with Fc portions for the
purpose of high-throughput screening assays to identify antagonists of hIL-5. (See, D.
25 Bennett et al., *J. Molecular Recognition* 8:52-58 (1995); K. Johanson et al., *J. Biol.*
Chem. 270:9459-9471 (1995).)

Moreover, the polypeptides of the present invention can be fused to marker
sequences, such as a peptide which facilitates purification of the fused polypeptide. In
preferred embodiments, the marker amino acid sequence is a hexa-histidine peptide,
30 such as the tag provided in a pQE vector (QIAGEN, Inc., 9259 Eton Avenue,
Chatsworth, CA, 91311), among others, many of which are commercially available.
As described in Gentz et al., *Proc. Natl. Acad. Sci. USA* 86:821-824 (1989), for
instance, hexa-histidine provides for convenient purification of the fusion protein.
Another peptide tag useful for purification, the "HA" tag, corresponds to an epitope
35 derived from the influenza hemagglutinin protein. (Wilson et al., *Cell* 37:767 (1984).)

Thus, any of these above fusions can be engineered using the polynucleotides
or the polypeptides of the present invention.

Vectors, Host Cells, and Protein Production

The present invention also relates to vectors containing the polynucleotide of the present invention, host cells, and the production of polypeptides by recombinant techniques. The vector may be, for example, a phage, plasmid, viral, or retroviral vector. Retroviral vectors may be replication competent or replication defective. In the latter case, viral propagation generally will occur only in complementing host cells.

The polynucleotides may be joined to a vector containing a selectable marker for propagation in a host. Generally, a plasmid vector is introduced in a precipitate, such as a calcium phosphate precipitate, or in a complex with a charged lipid. If the vector is a virus, it may be packaged in vitro using an appropriate packaging cell line and then transduced into host cells.

The polynucleotide insert should be operatively linked to an appropriate promoter, such as the phage lambda PL promoter, the E. coli lac, trp, phoA and tac promoters, the SV40 early and late promoters and promoters of retroviral LTRs, to name a few. Other suitable promoters will be known to the skilled artisan. The expression constructs will further contain sites for transcription initiation, termination, and, in the transcribed region, a ribosome binding site for translation. The coding portion of the transcripts expressed by the constructs will preferably include a translation initiating codon at the beginning and a termination codon (UAA, UGA or UAG) appropriately positioned at the end of the polypeptide to be translated.

As indicated, the expression vectors will preferably include at least one selectable marker. Such markers include dihydrofolate reductase, G418 or neomycin resistance for eukaryotic cell culture and tetracycline, kanamycin or ampicillin resistance genes for culturing in E. coli and other bacteria. Representative examples of appropriate hosts include, but are not limited to, bacterial cells, such as E. coli, Streptomyces and Salmonella typhimurium cells; fungal cells, such as yeast cells; insect cells such as Drosophila S2 and Spodoptera Sf9 cells; animal cells such as CHO, COS, 293, and Bowes melanoma cells; and plant cells. Appropriate culture mediums and conditions for the above-described host cells are known in the art.

Among vectors preferred for use in bacteria include pQE70, pQE60 and pQE-9, available from QIAGEN, Inc.; pBluescript vectors, Phagescript vectors, pNH8A, pNH16a, pNH18A, pNH46A, available from Stratagene Cloning Systems, Inc.; and ptrc99a, pKK223-3, pKK233-3, pDR540, pRIT5 available from Pharmacia Biotech, Inc. Among preferred eukaryotic vectors are pWLNEO, pSV2CAT, pOG44, pXT1 and pSG available from Stratagene; and pSVK3, pBPV, pMSG and pSVL available from Pharmacia. Other suitable vectors will be readily apparent to the skilled artisan.

Introduction of the construct into the host cell can be effected by calcium phosphate transfection, DEAE-dextran mediated transfection, cationic lipid-mediated transfection, electroporation, transduction, infection, or other methods. Such methods are described in many standard laboratory manuals, such as Davis et al., Basic Methods
5 In Molecular Biology (1986). It is specifically contemplated that the polypeptides of the present invention may in fact be expressed by a host cell lacking a recombinant vector.

A polypeptide of this invention can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography,
10 phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography ("HPLC") is employed for purification.

Polypeptides of the present invention, and preferably the secreted form, can also
15 be recovered from: products purified from natural sources, including bodily fluids, tissues and cells, whether directly isolated or cultured; products of chemical synthetic procedures; and products produced by recombinant techniques from a prokaryotic or eukaryotic host, including, for example, bacterial, yeast, higher plant, insect, and mammalian cells. Depending upon the host employed in a recombinant production
20 procedure, the polypeptides of the present invention may be glycosylated or may be non-glycosylated. In addition, polypeptides of the invention may also include an initial modified methionine residue, in some cases as a result of host-mediated processes. Thus, it is well known in the art that the N-terminal methionine encoded by the translation initiation codon generally is removed with high efficiency from any protein
25 after translation in all eukaryotic cells. While the N-terminal methionine on most proteins also is efficiently removed in most prokaryotes, for some proteins, this prokaryotic removal process is inefficient, depending on the nature of the amino acid to which the N-terminal methionine is covalently linked.

30 Uses of the Polynucleotides

Each of the polynucleotides identified herein can be used in numerous ways as reagents. The following description should be considered exemplary and utilizes known techniques.

The polynucleotides of the present invention are useful for chromosome
35 identification. There exists an ongoing need to identify new chromosome markers, since few chromosome marking reagents, based on actual sequence data (repeat

polymorphisms), are presently available. Each polynucleotide of the present invention can be used as a chromosome marker.

Briefly, sequences can be mapped to chromosomes by preparing PCR primers (preferably 15-25 bp) from the sequences shown in SEQ ID NO:X. Primers can be
5 selected using computer analysis so that primers do not span more than one predicted exon in the genomic DNA. These primers are then used for PCR screening of somatic cell hybrids containing individual human chromosomes. Only those hybrids containing the human gene corresponding to the SEQ ID NO:X will yield an amplified fragment.

Similarly, somatic hybrids provide a rapid method of PCR mapping the
10 polynucleotides to particular chromosomes. Three or more clones can be assigned per day using a single thermal cycler. Moreover, sublocalization of the polynucleotides can be achieved with panels of specific chromosome fragments. Other gene mapping strategies that can be used include in situ hybridization, prescreening with labeled flow-sorted chromosomes, and preselection by hybridization to construct chromosome
15 specific-cDNA libraries.

Precise chromosomal location of the polynucleotides can also be achieved using fluorescence in situ hybridization (FISH) of a metaphase chromosomal spread. This technique uses polynucleotides as short as 500 or 600 bases; however, polynucleotides 2,000-4,000 bp are preferred. For a review of this technique, see Verma et al.,
20 "Human Chromosomes: a Manual of Basic Techniques," Pergamon Press, New York (1988).

For chromosome mapping, the polynucleotides can be used individually (to mark a single chromosome or a single site on that chromosome) or in panels (for marking multiple sites and/or multiple chromosomes). Preferred polynucleotides
25 correspond to the noncoding regions of the cDNAs because the coding sequences are more likely conserved within gene families, thus increasing the chance of cross hybridization during chromosomal mapping.

Once a polynucleotide has been mapped to a precise chromosomal location, the physical position of the polynucleotide can be used in linkage analysis. Linkage
30 analysis establishes coinheritance between a chromosomal location and presentation of a particular disease. (Disease mapping data are found, for example, in V. McKusick, Mendelian Inheritance in Man (available on line through Johns Hopkins University Welch Medical Library) .) Assuming 1 megabase mapping resolution and one gene per 20 kb, a cDNA precisely localized to a chromosomal region associated with the disease
35 could be one of 50-500 potential causative genes.

Thus, once coinheritance is established, differences in the polynucleotide and the corresponding gene between affected and unaffected individuals can be examined.

First, visible structural alterations in the chromosomes, such as deletions or translocations, are examined in chromosome spreads or by PCR. If no structural alterations exist, the presence of point mutations are ascertained. Mutations observed in some or all affected individuals, but not in normal individuals, indicates that the mutation may cause the disease. However, complete sequencing of the polypeptide and the corresponding gene from several normal individuals is required to distinguish the mutation from a polymorphism. If a new polymorphism is identified, this polymorphic polypeptide can be used for further linkage analysis.

Furthermore, increased or decreased expression of the gene in affected individuals as compared to unaffected individuals can be assessed using polynucleotides of the present invention. Any of these alterations (altered expression, chromosomal rearrangement, or mutation) can be used as a diagnostic or prognostic marker.

In addition to the foregoing, a polynucleotide can be used to control gene expression through triple helix formation or antisense DNA or RNA. Both methods rely on binding of the polynucleotide to DNA or RNA. For these techniques, preferred polynucleotides are usually 20 to 40 bases in length and complementary to either the region of the gene involved in transcription (triple helix - see Lee et al., Nucl. Acids Res. 6:3073 (1979); Cooney et al., Science 241:456 (1988); and Dervan et al., Science 251:1360 (1991)) or to the mRNA itself (antisense - Okano, J. Neurochem. 56:560 (1991); Oligodeoxy-nucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988).) Triple helix formation optimally results in a shut-off of RNA transcription from DNA, while antisense RNA hybridization blocks translation of an mRNA molecule into polypeptide. Both techniques are effective in model systems, and the information disclosed herein can be used to design antisense or triple helix polynucleotides in an effort to treat disease.

Polynucleotides of the present invention are also useful in gene therapy. One goal of gene therapy is to insert a normal gene into an organism having a defective gene, in an effort to correct the genetic defect. The polynucleotides disclosed in the present invention offer a means of targeting such genetic defects in a highly accurate manner. Another goal is to insert a new gene that was not present in the host genome, thereby producing a new trait in the host cell.

The polynucleotides are also useful for identifying individuals from minute biological samples. The United States military, for example, is considering the use of restriction fragment length polymorphism (RFLP) for identification of its personnel. In this technique, an individual's genomic DNA is digested with one or more restriction enzymes, and probed on a Southern blot to yield unique bands for identifying

personnel. This method does not suffer from the current limitations of "Dog Tags" which can be lost, switched, or stolen, making positive identification difficult. The polynucleotides of the present invention can be used as additional DNA markers for RFLP.

5 The polynucleotides of the present invention can also be used as an alternative to RFLP, by determining the actual base-by-base DNA sequence of selected portions of an individual's genome. These sequences can be used to prepare PCR primers for amplifying and isolating such selected DNA, which can then be sequenced. Using this
10 of DNA sequences. Once an unique ID database is established for an individual, positive identification of that individual, living or dead, can be made from extremely small tissue samples.

 Forensic biology also benefits from using DNA-based identification techniques as disclosed herein. DNA sequences taken from very small biological samples such as
15 tissues, e.g., hair or skin, or body fluids, e.g., blood, saliva, semen, etc., can be amplified using PCR. In one prior art technique, gene sequences amplified from polymorphic loci, such as DQa class II HLA gene, are used in forensic biology to identify individuals. (Erlich, H., PCR Technology, Freeman and Co. (1992).) Once these specific polymorphic loci are amplified, they are digested with one or more
20 restriction enzymes, yielding an identifying set of bands on a Southern blot probed with DNA corresponding to the DQa class II HLA gene. Similarly, polynucleotides of the present invention can be used as polymorphic markers for forensic purposes.

 There is also a need for reagents capable of identifying the source of a particular tissue. Such need arises, for example, in forensics when presented with tissue of
25 unknown origin. Appropriate reagents can comprise, for example, DNA probes or primers specific to particular tissue prepared from the sequences of the present invention. Panels of such reagents can identify tissue by species and/or by organ type. In a similar fashion, these reagents can be used to screen tissue cultures for contamination.

30 In the very least, the polynucleotides of the present invention can be used as molecular weight markers on Southern gels, as diagnostic probes for the presence of a specific mRNA in a particular cell type, as a probe to "subtract-out" known sequences in the process of discovering novel polynucleotides, for selecting and making oligomers for attachment to a "gene chip" or other support, to raise anti-DNA antibodies using
35 DNA immunization techniques, and as an antigen to elicit an immune response.

Uses of the Polypeptides

Each of the polypeptides identified herein can be used in numerous ways. The following description should be considered exemplary and utilizes known techniques.

A polypeptide of the present invention can be used to assay protein levels in a biological sample using antibody-based techniques. For example, protein expression in tissues can be studied with classical immunohistological methods. (Jalkanen, M., et al., J. Cell. Biol. 101:976-985 (1985); Jalkanen, M., et al., J. Cell. Biol. 105:3087-3096 (1987).) Other antibody-based methods useful for detecting protein gene expression include immunoassays, such as the enzyme linked immunosorbent assay (ELISA) and the radioimmunoassay (RIA). Suitable antibody assay labels are known in the art and include enzyme labels, such as, glucose oxidase, and radioisotopes, such as iodine (^{125}I , ^{121}I), carbon (^{14}C), sulfur (^{35}S), tritium (^3H), indium (^{112}In), and technetium ($^{99\text{m}}\text{Tc}$), and fluorescent labels, such as fluorescein and rhodamine, and biotin.

In addition to assaying secreted protein levels in a biological sample, proteins can also be detected in vivo by imaging. Antibody labels or markers for in vivo imaging of protein include those detectable by X-radiography, NMR or ESR. For X-radiography, suitable labels include radioisotopes such as barium or cesium, which emit detectable radiation but are not overtly harmful to the subject. Suitable markers for NMR and ESR include those with a detectable characteristic spin, such as deuterium, which may be incorporated into the antibody by labeling of nutrients for the relevant hybridoma.

A protein-specific antibody or antibody fragment which has been labeled with an appropriate detectable imaging moiety, such as a radioisotope (for example, ^{131}I , ^{112}In , $^{99\text{m}}\text{Tc}$), a radio-opaque substance, or a material detectable by nuclear magnetic resonance, is introduced (for example, parenterally, subcutaneously, or intraperitoneally) into the mammal. It will be understood in the art that the size of the subject and the imaging system used will determine the quantity of imaging moiety needed to produce diagnostic images. In the case of a radioisotope moiety, for a human subject, the quantity of radioactivity injected will normally range from about 5 to 20 millicuries of $^{99\text{m}}\text{Tc}$. The labeled antibody or antibody fragment will then preferentially accumulate at the location of cells which contain the specific protein. In vivo tumor imaging is described in S.W. Burchiel et al., "Immunopharmacokinetics of Radiolabeled Antibodies and Their Fragments." (Chapter 13 in Tumor Imaging: The Radiochemical Detection of Cancer, S.W. Burchiel and B. A. Rhodes, eds., Masson Publishing Inc. (1982).)

Thus, the invention provides a diagnostic method of a disorder, which involves (a) assaying the expression of a polypeptide of the present invention in cells or body fluid of an individual; (b) comparing the level of gene expression with a standard gene expression level, whereby an increase or decrease in the assayed polypeptide gene expression level compared to the standard expression level is indicative of a disorder.

Moreover, polypeptides of the present invention can be used to treat disease. For example, patients can be administered a polypeptide of the present invention in an effort to replace absent or decreased levels of the polypeptide (e.g., insulin), to supplement absent or decreased levels of a different polypeptide (e.g., hemoglobin S for hemoglobin B), to inhibit the activity of a polypeptide (e.g., an oncogene), to activate the activity of a polypeptide (e.g., by binding to a receptor), to reduce the activity of a membrane bound receptor by competing with it for free ligand (e.g., soluble TNF receptors used in reducing inflammation), or to bring about a desired response (e.g., blood vessel growth).

Similarly, antibodies directed to a polypeptide of the present invention can also be used to treat disease. For example, administration of an antibody directed to a polypeptide of the present invention can bind and reduce overproduction of the polypeptide. Similarly, administration of an antibody can activate the polypeptide, such as by binding to a polypeptide bound to a membrane (receptor).

At the very least, the polypeptides of the present invention can be used as molecular weight markers on SDS-PAGE gels or on molecular sieve gel filtration columns using methods well known to those of skill in the art. Polypeptides can also be used to raise antibodies, which in turn are used to measure protein expression from a recombinant cell, as a way of assessing transformation of the host cell. Moreover, the polypeptides of the present invention can be used to test the following biological activities.

Biological Activities

The polynucleotides and polypeptides of the present invention can be used in assays to test for one or more biological activities. If these polynucleotides and polypeptides do exhibit activity in a particular assay, it is likely that these molecules may be involved in the diseases associated with the biological activity. Thus, the polynucleotides and polypeptides could be used to treat the associated disease.

Immune Activity

A polypeptide or polynucleotide of the present invention may be useful in treating deficiencies or disorders of the immune system, by activating or inhibiting the

proliferation, differentiation, or mobilization (chemotaxis) of immune cells. Immune cells develop through a process called hematopoiesis, producing myeloid (platelets, red blood cells, neutrophils, and macrophages) and lymphoid (B and T lymphocytes) cells from pluripotent stem cells. The etiology of these immune deficiencies or disorders
5 may be genetic, somatic, such as cancer or some autoimmune disorders, acquired (e.g., by chemotherapy or toxins), or infectious. Moreover, a polynucleotide or polypeptide of the present invention can be used as a marker or detector of a particular immune system disease or disorder.

A polynucleotide or polypeptide of the present invention may be useful in
10 treating or detecting deficiencies or disorders of hematopoietic cells. A polypeptide or polynucleotide of the present invention could be used to increase differentiation and proliferation of hematopoietic cells, including the pluripotent stem cells, in an effort to treat those disorders associated with a decrease in certain (or many) types hematopoietic cells. Examples of immunologic deficiency syndromes include, but are not limited to:
15 blood protein disorders (e.g. agammaglobulinemia, dysgammaglobulinemia), ataxia telangiectasia, common variable immunodeficiency, Digeorge Syndrome, HIV infection, HTLV-BLV infection, leukocyte adhesion deficiency syndrome, lymphopenia, phagocyte bactericidal dysfunction, severe combined immunodeficiency (SCIDs), Wiskott-Aldrich Disorder, anemia, thrombocytopenia, or hemoglobinuria.

Moreover, a polypeptide or polynucleotide of the present invention could also
20 be used to modulate hemostatic (the stopping of bleeding) or thrombolytic activity (clot formation). For example, by increasing hemostatic or thrombolytic activity, a polynucleotide or polypeptide of the present invention could be used to treat blood coagulation disorders (e.g., afibrinogenemia, factor deficiencies), blood platelet
25 disorders (e.g. thrombocytopenia), or wounds resulting from trauma, surgery, or other causes. Alternatively, a polynucleotide or polypeptide of the present invention that can decrease hemostatic or thrombolytic activity could be used to inhibit or dissolve clotting. These molecules could be important in the treatment of heart attacks (infarction), strokes, or scarring.

A polynucleotide or polypeptide of the present invention may also be useful in
30 treating or detecting autoimmune disorders. Many autoimmune disorders result from inappropriate recognition of self as foreign material by immune cells. This inappropriate recognition results in an immune response leading to the destruction of the host tissue. Therefore, the administration of a polypeptide or polynucleotide of the
35 present invention that inhibits an immune response, particularly the proliferation, differentiation, or chemotaxis of T-cells, may be an effective therapy in preventing autoimmune disorders.

Examples of autoimmune disorders that can be treated or detected by the present invention include, but are not limited to: Addison's Disease, hemolytic anemia, antiphospholipid syndrome, rheumatoid arthritis, dermatitis, allergic encephalomyelitis, glomerulonephritis, Goodpasture's Syndrome, Graves' Disease, Multiple Sclerosis, Myasthenia Gravis, Neuritis, Ophthalmia, Bullous Pemphigoid, Pemphigus, Polyendocrinopathies, Purpura, Reiter's Disease, Stiff-Man Syndrome, Autoimmune Thyroiditis, Systemic Lupus Erythematosus, Autoimmune Pulmonary Inflammation, Guillain-Barre Syndrome, insulin dependent diabetes mellitus, and autoimmune inflammatory eye disease.

Similarly, allergic reactions and conditions, such as asthma (particularly allergic asthma) or other respiratory problems, may also be treated by a polypeptide or polynucleotide of the present invention. Moreover, these molecules can be used to treat anaphylaxis, hypersensitivity to an antigenic molecule, or blood group incompatibility.

A polynucleotide or polypeptide of the present invention may also be used to treat and/or prevent organ rejection or graft-versus-host disease (GVHD). Organ rejection occurs by host immune cell destruction of the transplanted tissue through an immune response. Similarly, an immune response is also involved in GVHD, but, in this case, the foreign transplanted immune cells destroy the host tissues. The administration of a polypeptide or polynucleotide of the present invention that inhibits an immune response, particularly the proliferation, differentiation, or chemotaxis of T-cells, may be an effective therapy in preventing organ rejection or GVHD.

Similarly, a polypeptide or polynucleotide of the present invention may also be used to modulate inflammation. For example, the polypeptide or polynucleotide may inhibit the proliferation and differentiation of cells involved in an inflammatory response. These molecules can be used to treat inflammatory conditions, both chronic and acute conditions, including inflammation associated with infection (e.g., septic shock, sepsis, or systemic inflammatory response syndrome (SIRS)), ischemia-reperfusion injury, endotoxin lethality, arthritis, complement-mediated hyperacute rejection, nephritis, cytokine or chemokine induced lung injury, inflammatory bowel disease, Crohn's disease, or resulting from over production of cytokines (e.g., TNF or IL-1.)

Hyperproliferative Disorders

A polypeptide or polynucleotide can be used to treat or detect hyperproliferative disorders, including neoplasms. A polypeptide or polynucleotide of the present invention may inhibit the proliferation of the disorder through direct or indirect

interactions. Alternatively, a polypeptide or polynucleotide of the present invention may proliferate other cells which can inhibit the hyperproliferative disorder.

For example, by increasing an immune response, particularly increasing antigenic qualities of the hyperproliferative disorder or by proliferating, differentiating, or mobilizing T-cells, hyperproliferative disorders can be treated. This immune response may be increased by either enhancing an existing immune response, or by initiating a new immune response. Alternatively, decreasing an immune response may also be a method of treating hyperproliferative disorders, such as a chemotherapeutic agent.

Examples of hyperproliferative disorders that can be treated or detected by a polynucleotide or polypeptide of the present invention include, but are not limited to neoplasms located in the: abdomen, bone, breast, digestive system, liver, pancreas, peritoneum, endocrine glands (adrenal, parathyroid, pituitary, testicles, ovary, thymus, thyroid), eye, head and neck, nervous (central and peripheral), lymphatic system, pelvic, skin, soft tissue, spleen, thoracic, and urogenital.

Similarly, other hyperproliferative disorders can also be treated or detected by a polynucleotide or polypeptide of the present invention. Examples of such hyperproliferative disorders include, but are not limited to: hypergammaglobulinemia, lymphoproliferative disorders, paraproteinemias, purpura, sarcoidosis, Sezary Syndrome, Waldenstrom's Macroglobulinemia, Gaucher's Disease, histiocytosis, and any other hyperproliferative disease, besides neoplasia, located in an organ system listed above.

Infectious Disease

A polypeptide or polynucleotide of the present invention can be used to treat or detect infectious agents. For example, by increasing the immune response, particularly increasing the proliferation and differentiation of B and/or T cells, infectious diseases may be treated. The immune response may be increased by either enhancing an existing immune response, or by initiating a new immune response. Alternatively, the polypeptide or polynucleotide of the present invention may also directly inhibit the infectious agent, without necessarily eliciting an immune response.

Viruses are one example of an infectious agent that can cause disease or symptoms that can be treated or detected by a polynucleotide or polypeptide of the present invention. Examples of viruses, include, but are not limited to the following DNA and RNA viral families: Arbovirus, Adenoviridae, Arenaviridae, Arterivirus, Birnaviridae, Bunyaviridae, Caliciviridae, Circoviridae, Coronaviridae, Flaviviridae, Hepadnaviridae (Hepatitis), Herpesviridae (such as, Cytomegalovirus, Herpes

Simplex, Herpes Zoster), Mononegavirus (e.g., Paramyxoviridae, Morbillivirus, Rhabdoviridae), Orthomyxoviridae (e.g., Influenza), Papovaviridae, Parvoviridae, Picornaviridae, Poxviridae (such as Smallpox or Vaccinia), Reoviridae (e.g., Rotavirus), Retroviridae (HTLV-I, HTLV-II, Lentivirus), and Togaviridae (e.g., Rubivirus). Viruses falling within these families can cause a variety of diseases or symptoms, including, but not limited to: arthritis, bronchiolitis, encephalitis, eye infections (e.g., conjunctivitis, keratitis), chronic fatigue syndrome, hepatitis (A, B, C, E, Chronic Active, Delta), meningitis, opportunistic infections (e.g., AIDS), pneumonia, Burkitt's Lymphoma, chickenpox, hemorrhagic fever, Measles, Mumps, Parainfluenza, Rabies, the common cold, Polio, leukemia, Rubella, sexually transmitted diseases, skin diseases (e.g., Kaposi's, warts), and viremia. A polypeptide or polynucleotide of the present invention can be used to treat or detect any of these symptoms or diseases.

Similarly, bacterial or fungal agents that can cause disease or symptoms and that can be treated or detected by a polynucleotide or polypeptide of the present invention include, but not limited to, the following Gram-Negative and Gram-positive bacterial families and fungi: Actinomycetales (e.g., Corynebacterium, Mycobacterium, Norcardia), Aspergillosis, Bacillaceae (e.g., Anthrax, Clostridium), Bacteroidaceae, Blastomycosis, Bordetella, Borrelia, Brucellosis, Candidiasis, Campylobacter, Coccidioidomycosis, Cryptococcosis, Dermatocycoses, Enterobacteriaceae (Klebsiella, Salmonella, Serratia, Yersinia), Erysipelothrix, Helicobacter, Legionellosis, Leptospirosis, Listeria, Mycoplasmatales, Neisseriaceae (e.g., Acinetobacter, Gonorrhea, Meningococcal), Pasteurellaceae Infections (e.g., Actinobacillus, Haemophilus, Pasteurella), Pseudomonas, Rickettsiaceae, Chlamydiaceae, Syphilis, and Staphylococcal. These bacterial or fungal families can cause the following diseases or symptoms, including, but not limited to: bacteremia, endocarditis, eye infections (conjunctivitis, tuberculosis, uveitis), gingivitis, opportunistic infections (e.g., AIDS related infections), paronychia, prosthesis-related infections, Reiter's Disease, respiratory tract infections, such as Whooping Cough or Empyema, sepsis, Lyme Disease, Cat-Scratch Disease, Dysentery, Paratyphoid Fever, food poisoning, Typhoid, pneumonia, Gonorrhea, meningitis, Chlamydia, Syphilis, Diphtheria, Leprosy, Paratuberculosis, Tuberculosis, Lupus, Botulism, gangrene, tetanus, impetigo, Rheumatic Fever, Scarlet Fever, sexually transmitted diseases, skin diseases (e.g., cellulitis, dermatocycoses), toxemia, urinary tract infections, wound infections. A polypeptide or polynucleotide of the present invention can be used to treat or detect any of these symptoms or diseases.

Moreover, parasitic agents causing disease or symptoms that can be treated or detected by a polynucleotide or polypeptide of the present invention include, but not limited to, the following families: Amebiasis, Babesiosis, Coccidiosis, Cryptosporidiosis, Dientamoebiasis, Dourine, Ectoparasitic, Giardiasis, Helminthiasis, Leishmaniasis, Theileriasis, Toxoplasmosis, Trypanosomiasis, and Trichomonas. These parasites can cause a variety of diseases or symptoms, including, but not limited to: Scabies, Trombiculiasis, eye infections, intestinal disease (e.g., dysentery, giardiasis), liver disease, lung disease, opportunistic infections (e.g., AIDS related), Malaria, pregnancy complications, and toxoplasmosis. A polypeptide or polynucleotide of the present invention can be used to treat or detect any of these symptoms or diseases.

Preferably, treatment using a polypeptide or polynucleotide of the present invention could either be by administering an effective amount of a polypeptide to the patient, or by removing cells from the patient, supplying the cells with a polynucleotide of the present invention, and returning the engineered cells to the patient (ex vivo therapy). Moreover, the polypeptide or polynucleotide of the present invention can be used as an antigen in a vaccine to raise an immune response against infectious disease.

Regeneration

A polynucleotide or polypeptide of the present invention can be used to differentiate, proliferate, and attract cells, leading to the regeneration of tissues. (See, Science 276:59-87 (1997).) The regeneration of tissues could be used to repair, replace, or protect tissue damaged by congenital defects, trauma (wounds, burns, incisions, or ulcers), age, disease (e.g. osteoporosis, osteoarthritis, periodontal disease, liver failure), surgery, including cosmetic plastic surgery, fibrosis, reperfusion injury, or systemic cytokine damage.

Tissues that could be regenerated using the present invention include organs (e.g., pancreas, liver, intestine, kidney, skin, endothelium), muscle (smooth, skeletal or cardiac), vascular (including vascular endothelium), nervous, hematopoietic, and skeletal (bone, cartilage, tendon, and ligament) tissue. Preferably, regeneration occurs without or decreased scarring. Regeneration also may include angiogenesis.

Moreover, a polynucleotide or polypeptide of the present invention may increase regeneration of tissues difficult to heal. For example, increased tendon/ligament regeneration would quicken recovery time after damage. A polynucleotide or polypeptide of the present invention could also be used prophylactically in an effort to avoid damage. Specific diseases that could be treated include of tendinitis, carpal tunnel syndrome, and other tendon or ligament defects. A further example of tissue

regeneration of non-healing wounds includes pressure ulcers, ulcers associated with vascular insufficiency, surgical, and traumatic wounds.

Similarly, nerve and brain tissue could also be regenerated by using a polynucleotide or polypeptide of the present invention to proliferate and differentiate
5 nerve cells. Diseases that could be treated using this method include central and peripheral nervous system diseases, neuropathies, or mechanical and traumatic disorders (e.g., spinal cord disorders, head trauma, cerebrovascular disease, and stoke). Specifically, diseases associated with peripheral nerve injuries, peripheral
10 neuropathy (e.g., resulting from chemotherapy or other medical therapies), localized neuropathies, and central nervous system diseases (e.g., Alzheimer's disease, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, and Shy-Drager syndrome), could all be treated using the polynucleotide or polypeptide of the present invention.

15 **Chemotaxis**

A polynucleotide or polypeptide of the present invention may have chemotaxis activity. A chemotactic molecule attracts or mobilizes cells (e.g., monocytes, fibroblasts, neutrophils, T-cells, mast cells, eosinophils, epithelial and/or endothelial cells) to a particular site in the body, such as inflammation, infection, or site of
20 hyperproliferation. The mobilized cells can then fight off and/or heal the particular trauma or abnormality.

A polynucleotide or polypeptide of the present invention may increase chemotactic activity of particular cells. These chemotactic molecules can then be used to treat inflammation, infection, hyperproliferative disorders, or any immune system
25 disorder by increasing the number of cells targeted to a particular location in the body. For example, chemotactic molecules can be used to treat wounds and other trauma to tissues by attracting immune cells to the injured location. Chemotactic molecules of the present invention can also attract fibroblasts, which can be used to treat wounds.

It is also contemplated that a polynucleotide or polypeptide of the present
30 invention may inhibit chemotactic activity. These molecules could also be used to treat disorders. Thus, a polynucleotide or polypeptide of the present invention could be used as an inhibitor of chemotaxis.

Binding Activity

35 A polypeptide of the present invention may be used to screen for molecules that bind to the polypeptide or for molecules to which the polypeptide binds. The binding of the polypeptide and the molecule may activate (agonist), increase, inhibit

(antagonist), or decrease activity of the polypeptide or the molecule bound. Examples of such molecules include antibodies, oligonucleotides, proteins (e.g., receptors), or small molecules.

Preferably, the molecule is closely related to the natural ligand of the polypeptide, e.g., a fragment of the ligand, or a natural substrate, a ligand, a structural or functional mimetic. (See, Coligan et al., Current Protocols in Immunology 1(2):Chapter 5 (1991).) Similarly, the molecule can be closely related to the natural receptor to which the polypeptide binds, or at least, a fragment of the receptor capable of being bound by the polypeptide (e.g., active site). In either case, the molecule can be rationally designed using known techniques.

Preferably, the screening for these molecules involves producing appropriate cells which express the polypeptide, either as a secreted protein or on the cell membrane. Preferred cells include cells from mammals, yeast, *Drosophila*, or *E. coli*. Cells expressing the polypeptide (or cell membrane containing the expressed polypeptide) are then preferably contacted with a test compound potentially containing the molecule to observe binding, stimulation, or inhibition of activity of either the polypeptide or the molecule.

The assay may simply test binding of a candidate compound to the polypeptide, wherein binding is detected by a label, or in an assay involving competition with a labeled competitor. Further, the assay may test whether the candidate compound results in a signal generated by binding to the polypeptide.

Alternatively, the assay can be carried out using cell-free preparations, polypeptide/molecule affixed to a solid support, chemical libraries, or natural product mixtures. The assay may also simply comprise the steps of mixing a candidate compound with a solution containing a polypeptide, measuring polypeptide/molecule activity or binding, and comparing the polypeptide/molecule activity or binding to a standard.

Preferably, an ELISA assay can measure polypeptide level or activity in a sample (e.g., biological sample) using a monoclonal or polyclonal antibody. The antibody can measure polypeptide level or activity by either binding, directly or indirectly, to the polypeptide or by competing with the polypeptide for a substrate.

All of these above assays can be used as diagnostic or prognostic markers. The molecules discovered using these assays can be used to treat disease or to bring about a particular result in a patient (e.g., blood vessel growth) by activating or inhibiting the polypeptide/molecule. Moreover, the assays can discover agents which may inhibit or enhance the production of the polypeptide from suitably manipulated cells or tissues.

Therefore, the invention includes a method of identifying compounds which bind to a polypeptide of the invention comprising the steps of: (a) incubating a candidate binding compound with a polypeptide of the invention; and (b) determining if binding has occurred. Moreover, the invention includes a method of identifying agonists/antagonists comprising the steps of: (a) incubating a candidate compound with a polypeptide of the invention, (b) assaying a biological activity, and (b) determining if a biological activity of the polypeptide has been altered.

Other Activities

10 A polypeptide or polynucleotide of the present invention may also increase or decrease the differentiation or proliferation of embryonic stem cells, besides, as discussed above, hematopoietic lineage.

A polypeptide or polynucleotide of the present invention may also be used to modulate mammalian characteristics, such as body height, weight, hair color, eye color, skin, percentage of adipose tissue, pigmentation, size, and shape (e.g., cosmetic surgery). Similarly, a polypeptide or polynucleotide of the present invention may be used to modulate mammalian metabolism affecting catabolism, anabolism, processing, utilization, and storage of energy.

20 A polypeptide or polynucleotide of the present invention may be used to change a mammal's mental state or physical state by influencing biorhythms, circadian rhythms, depression (including depressive disorders), tendency for violence, tolerance for pain, reproductive capabilities (preferably by Activin or Inhibin-like activity), hormonal or endocrine levels, appetite, libido, memory, stress, or other cognitive qualities.

25 A polypeptide or polynucleotide of the present invention may also be used as a food additive or preservative, such as to increase or decrease storage capabilities, fat content, lipid, protein, carbohydrate, vitamins, minerals, cofactors or other nutritional components.

Other Preferred Embodiments

Other preferred embodiments of the claimed invention include an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least about 50 contiguous nucleotides in the nucleotide sequence of SEQ ID NO:X wherein X is any integer as defined in Table 1.

35 Also preferred is a nucleic acid molecule wherein said sequence of contiguous nucleotides is included in the nucleotide sequence of SEQ ID NO:X in the range of

positions beginning with the nucleotide at about the position of the 5' Nucleotide of the Clone Sequence and ending with the nucleotide at about the position of the 3' Nucleotide of the Clone Sequence as defined for SEQ ID NO:X in Table 1.

Also preferred is a nucleic acid molecule wherein said sequence of contiguous
5 nucleotides is included in the nucleotide sequence of SEQ ID NO:X in the range of positions beginning with the nucleotide at about the position of the 5' Nucleotide of the Start Codon and ending with the nucleotide at about the position of the 3' Nucleotide of the Clone Sequence as defined for SEQ ID NO:X in Table 1.

Similarly preferred is a nucleic acid molecule wherein said sequence of
10 contiguous nucleotides is included in the nucleotide sequence of SEQ ID NO:X in the range of positions beginning with the nucleotide at about the position of the 5' Nucleotide of the First Amino Acid of the Signal Peptide and ending with the nucleotide at about the position of the 3' Nucleotide of the Clone Sequence as defined for SEQ ID NO:X in Table 1.

Also preferred is an isolated nucleic acid molecule comprising a nucleotide
15 sequence which is at least 95% identical to a sequence of at least about 150 contiguous nucleotides in the nucleotide sequence of SEQ ID NO:X.

Further preferred is an isolated nucleic acid molecule comprising a nucleotide
sequence which is at least 95% identical to a sequence of at least about 500 contiguous
20 nucleotides in the nucleotide sequence of SEQ ID NO:X.

A further preferred embodiment is a nucleic acid molecule comprising a
nucleotide sequence which is at least 95% identical to the nucleotide sequence of SEQ
ID NO:X beginning with the nucleotide at about the position of the 5' Nucleotide of the
First Amino Acid of the Signal Peptide and ending with the nucleotide at about the
25 position of the 3' Nucleotide of the Clone Sequence as defined for SEQ ID NO:X in
Table 1.

A further preferred embodiment is an isolated nucleic acid molecule comprising
a nucleotide sequence which is at least 95% identical to the complete nucleotide
sequence of SEQ ID NO:X.

Also preferred is an isolated nucleic acid molecule which hybridizes under
30 stringent hybridization conditions to a nucleic acid molecule, wherein said nucleic acid molecule which hybridizes does not hybridize under stringent hybridization conditions to a nucleic acid molecule having a nucleotide sequence consisting of only A residues or of only T residues.

Also preferred is a composition of matter comprising a DNA molecule which
35 comprises a human cDNA clone identified by a cDNA Clone Identifier in Table 1, which DNA molecule is contained in the material deposited with the American Type

Culture Collection and given the ATCC Deposit Number shown in Table 1 for said cDNA Clone Identifier.

Also preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least 50 contiguous nucleotides in the nucleotide sequence of a human cDNA clone identified by a cDNA Clone Identifier in Table 1, which DNA molecule is contained in the deposit given the ATCC Deposit Number shown in Table 1.

Also preferred is an isolated nucleic acid molecule, wherein said sequence of at least 50 contiguous nucleotides is included in the nucleotide sequence of the complete open reading frame sequence encoded by said human cDNA clone.

Also preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to sequence of at least 150 contiguous nucleotides in the nucleotide sequence encoded by said human cDNA clone.

A further preferred embodiment is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to sequence of at least 500 contiguous nucleotides in the nucleotide sequence encoded by said human cDNA clone.

A further preferred embodiment is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to the complete nucleotide sequence encoded by said human cDNA clone.

A further preferred embodiment is a method for detecting in a biological sample a nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X wherein X is any integer as defined in Table 1; and a nucleotide sequence encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1; which method comprises a step of comparing a nucleotide sequence of at least one nucleic acid molecule in said sample with a sequence selected from said group and determining whether the sequence of said nucleic acid molecule in said sample is at least 95% identical to said selected sequence.

Also preferred is the above method wherein said step of comparing sequences comprises determining the extent of nucleic acid hybridization between nucleic acid molecules in said sample and a nucleic acid molecule comprising said sequence selected from said group. Similarly, also preferred is the above method wherein said step of comparing sequences is performed by comparing the nucleotide sequence determined from a nucleic acid molecule in said sample with said sequence selected from said group. The nucleic acid molecules can comprise DNA molecules or RNA molecules.

A further preferred embodiment is a method for identifying the species, tissue or cell type of a biological sample which method comprises a step of detecting nucleic acid molecules in said sample, if any, comprising a nucleotide sequence that is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X wherein X is any integer as defined in Table 1; and a nucleotide sequence encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

The method for identifying the species, tissue or cell type of a biological sample can comprise a step of detecting nucleic acid molecules comprising a nucleotide sequence in a panel of at least two nucleotide sequences, wherein at least one sequence in said panel is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from said group.

Also preferred is a method for diagnosing in a subject a pathological condition associated with abnormal structure or expression of a gene encoding a secreted protein identified in Table 1, which method comprises a step of detecting in a biological sample obtained from said subject nucleic acid molecules, if any, comprising a nucleotide sequence that is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X wherein X is any integer as defined in Table 1; and a nucleotide sequence encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

The method for diagnosing a pathological condition can comprise a step of detecting nucleic acid molecules comprising a nucleotide sequence in a panel of at least two nucleotide sequences, wherein at least one sequence in said panel is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from said group.

Also preferred is a composition of matter comprising isolated nucleic acid molecules wherein the nucleotide sequences of said nucleic acid molecules comprise a panel of at least two nucleotide sequences, wherein at least one sequence in said panel is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X wherein X is any integer as defined in Table 1; and a nucleotide sequence encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1. The nucleic acid molecules can comprise DNA molecules or RNA molecules.

Also preferred is an isolated polypeptide comprising an amino acid sequence at least 90% identical to a sequence of at least about 10 contiguous amino acids in the amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1.

Also preferred is a polypeptide, wherein said sequence of contiguous amino acids is included in the amino acid sequence of SEQ ID NO:Y in the range of positions beginning with the residue at about the position of the First Amino Acid of the Secreted Portion and ending with the residue at about the Last Amino Acid of the Open Reading Frame as set forth for SEQ ID NO:Y in Table 1.

Also preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence of at least about 30 contiguous amino acids in the amino acid sequence of SEQ ID NO:Y.

Further preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence of at least about 100 contiguous amino acids in the amino acid sequence of SEQ ID NO:Y.

Further preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to the complete amino acid sequence of SEQ ID NO:Y.

Further preferred is an isolated polypeptide comprising an amino acid sequence at least 90% identical to a sequence of at least about 10 contiguous amino acids in the complete amino acid sequence of a secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is a polypeptide wherein said sequence of contiguous amino acids is included in the amino acid sequence of a secreted portion of the secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence of at least about 30 contiguous amino acids in the amino acid sequence of the secreted portion of the protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence of at least about 100 contiguous amino acids in the amino acid sequence of the secreted portion of the protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to the amino acid sequence of the secreted portion of the protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Further preferred is an isolated antibody which binds specifically to a polypeptide comprising an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Further preferred is a method for detecting in a biological sample a polypeptide comprising an amino acid sequence which is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1; which method comprises a step of comparing an amino acid sequence of at least one polypeptide molecule in said sample with a sequence selected from said group and determining whether the sequence of said polypeptide molecule in said sample is at least 90% identical to said sequence of at least 10 contiguous amino acids.

Also preferred is the above method wherein said step of comparing an amino acid sequence of at least one polypeptide molecule in said sample with a sequence selected from said group comprises determining the extent of specific binding of polypeptides in said sample to an antibody which binds specifically to a polypeptide comprising an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is the above method wherein said step of comparing sequences is performed by comparing the amino acid sequence determined from a polypeptide molecule in said sample with said sequence selected from said group.

Also preferred is a method for identifying the species, tissue or cell type of a biological sample which method comprises a step of detecting polypeptide molecules in said sample, if any, comprising an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is the above method for identifying the species, tissue or cell type of a biological sample, which method comprises a step of detecting polypeptide molecules comprising an amino acid sequence in a panel of at least two amino acid sequences, wherein at least one sequence in said panel is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the above group.

Also preferred is a method for diagnosing in a subject a pathological condition associated with abnormal structure or expression of a gene encoding a secreted protein identified in Table 1, which method comprises a step of detecting in a biological sample obtained from said subject polypeptide molecules comprising an amino acid sequence in a panel of at least two amino acid sequences, wherein at least one sequence in said panel is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

In any of these methods, the step of detecting said polypeptide molecules includes using an antibody.

Also preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a nucleotide sequence encoding a polypeptide wherein said polypeptide comprises an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is an isolated nucleic acid molecule, wherein said nucleotide sequence encoding a polypeptide has been optimized for expression of said polypeptide in a prokaryotic host.

Also preferred is an isolated nucleic acid molecule, wherein said polypeptide
5 comprises an amino acid sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

10 Further preferred is a method of making a recombinant vector comprising inserting any of the above isolated nucleic acid molecule into a vector. Also preferred is the recombinant vector produced by this method. Also preferred is a method of making a recombinant host cell comprising introducing the vector into a host cell, as well as the recombinant host cell produced by this method.

15 Also preferred is a method of making an isolated polypeptide comprising culturing this recombinant host cell under conditions such that said polypeptide is expressed and recovering said polypeptide. Also preferred is this method of making an isolated polypeptide, wherein said recombinant host cell is a eukaryotic cell and said polypeptide is a secreted portion of a human secreted protein comprising an amino acid
20 sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y beginning with the residue at the position of the First Amino Acid of the Secreted Portion of SEQ ID NO:Y wherein Y is an integer set forth in Table 1 and said position of the First Amino Acid of the Secreted Portion of SEQ ID NO:Y is defined in Table 1; and an amino acid sequence of a secreted portion of a protein encoded by a human
25 cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1. The isolated polypeptide produced by this method is also preferred.

Also preferred is a method of treatment of an individual in need of an increased level of a secreted protein activity, which method comprises administering to such an
30 individual a pharmaceutical composition comprising an amount of an isolated polypeptide, polynucleotide, or antibody of the claimed invention effective to increase the level of said protein activity in said individual.

Having generally described the invention, the same will be more readily understood by reference to the following examples, which are provided by way of
35 illustration and are not intended as limiting.

Examples

Example 1: Isolation of a Selected cDNA Clone From the Deposited Sample

5 Each cDNA clone in a cited ATCC deposit is contained in a plasmid vector. Table 1 identifies the vectors used to construct the cDNA library from which each clone was isolated. In many cases, the vector used to construct the library is a phage vector from which a plasmid has been excised. The table immediately below correlates the related plasmid for each phage vector used in constructing the cDNA library. For
10 example, where a particular clone is identified in Table 1 as being isolated in the vector "Lambda Zap," the corresponding deposited clone is in "pBluescript."

	<u>Vector Used to Construct Library</u>	<u>Corresponding Deposited Plasmid</u>
	Lambda Zap	pBluescript (pBS)
	Uni-Zap XR	pBluescript (pBS)
15	Zap Express	pBK
	lafmid BA	plafmid BA
	pSport1	pSport1
	pCMVSPORT 2.0	pCMVSPORT 2.0
	pCMVSPORT 3.0	pCMVSPORT 3.0
20	pCR [®] 2.1	pCR [®] 2.1

Vectors Lambda Zap (U.S. Patent Nos. 5,128,256 and 5,286,636), Uni-Zap XR (U.S. Patent Nos. 5,128, 256 and 5,286,636), Zap Express (U.S. Patent Nos. 5,128,256 and 5,286,636), pBluescript (pBS) (Short, J. M. et al., Nucleic Acids Res. 16:7583-7600 (1988); Alting-Mees, M. A. and Short, J. M., Nucleic Acids Res. 25 17:9494 (1989)) and pBK (Alting-Mees, M. A. et al., Strategies 5:58-61 (1992)) are commercially available from Stratagene Cloning Systems, Inc., 11011 N. Torrey Pines Road, La Jolla, CA, 92037. pBS contains an ampicillin resistance gene and pBK contains a neomycin resistance gene. Both can be transformed into E. coli strain XL-1 Blue, also available from Stratagene. pBS comes in 4 forms SK+, SK-, KS+ and KS. 30 The S and K refers to the orientation of the polylinker to the T7 and T3 primer sequences which flank the polylinker region ("S" is for SacI and "K" is for KpnI which are the first sites on each respective end of the linker). "+" or "-" refer to the orientation of the f1 origin of replication ("ori"), such that in one orientation, single stranded rescue initiated from the f1 ori generates sense strand DNA and in the other, antisense.

35 Vectors pSport1, pCMVSPORT 2.0 and pCMVSPORT 3.0, were obtained from Life Technologies, Inc., P. O. Box 6009, Gaithersburg, MD 20897. All Sport vectors contain an ampicillin resistance gene and may be transformed into E. coli strain

DH10B, also available from Life Technologies. (See, for instance, Gruber, C. E., et al., *Focus* 15:59 (1993).) Vector lafmid BA (Bento Soares, Columbia University, NY) contains an ampicillin resistance gene and can be transformed into *E. coli* strain XL-1 Blue. Vector pCR[®]2.1, which is available from Invitrogen, 1600 Faraday Avenue, Carlsbad, CA 92008, contains an ampicillin resistance gene and may be transformed into *E. coli* strain DH10B, available from Life Technologies. (See, for instance, Clark, J. M., *Nuc. Acids Res.* 16:9677-9686 (1988) and Mead, D. et al., *Bio/Technology* 9: (1991).) Preferably, a polynucleotide of the present invention does not comprise the phage vector sequences identified for the particular clone in Table 1, as well as the corresponding plasmid vector sequences designated above.

The deposited material in the sample assigned the ATCC Deposit Number cited in Table 1 for any given cDNA clone also may contain one or more additional plasmids, each comprising a cDNA clone different from that given clone. Thus, deposits sharing the same ATCC Deposit Number contain at least a plasmid for each cDNA clone identified in Table 1. Typically, each ATCC deposit sample cited in Table 1 comprises a mixture of approximately equal amounts (by weight) of about 50 plasmid DNAs, each containing a different cDNA clone; but such a deposit sample may include plasmids for more or less than 50 cDNA clones, up to about 500 cDNA clones.

Two approaches can be used to isolate a particular clone from the deposited sample of plasmid DNAs cited for that clone in Table 1. First, a plasmid is directly isolated by screening the clones using a polynucleotide probe corresponding to SEQ ID NO:X.

Particularly, a specific polynucleotide with 30-40 nucleotides is synthesized using an Applied Biosystems DNA synthesizer according to the sequence reported. The oligonucleotide is labeled, for instance, with ³²P-γ-ATP using T4 polynucleotide kinase and purified according to routine methods. (E.g., Maniatis et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Press, Cold Spring, NY (1982).) The plasmid mixture is transformed into a suitable host, as indicated above (such as XL-1 Blue (Stratagene)) using techniques known to those of skill in the art, such as those provided by the vector supplier or in related publications or patents cited above. The transformants are plated on 1.5% agar plates (containing the appropriate selection agent, e.g., ampicillin) to a density of about 150 transformants (colonies) per plate. These plates are screened using Nylon membranes according to routine methods for bacterial colony screening (e.g., Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2nd Edit., (1989), Cold Spring Harbor Laboratory Press, pages 1.93 to 1.104), or other techniques known to those of skill in the art.

Alternatively, two primers of 17-20 nucleotides derived from both ends of the SEQ ID NO:X (i.e., within the region of SEQ ID NO:X bounded by the 5' NT and the 3' NT of the clone defined in Table 1) are synthesized and used to amplify the desired cDNA using the deposited cDNA plasmid as a template. The polymerase chain reaction is carried out under routine conditions, for instance, in 25 μ l of reaction mixture with 0.5 μ g of the above cDNA template. A convenient reaction mixture is 1.5-5 mM $MgCl_2$, 0.01% (w/v) gelatin, 20 μ M each of dATP, dCTP, dGTP, dTTP, 25 pmol of each primer and 0.25 Unit of Taq polymerase. Thirty five cycles of PCR (denaturation at 94°C for 1 min; annealing at 55°C for 1 min; elongation at 72°C for 1 min) are performed with a Perkin-Elmer Cetus automated thermal cycler. The amplified product is analyzed by agarose gel electrophoresis and the DNA band with expected molecular weight is excised and purified. The PCR product is verified to be the selected sequence by subcloning and sequencing the DNA product.

Several methods are available for the identification of the 5' or 3' non-coding portions of a gene which may not be present in the deposited clone. These methods include but are not limited to, filter probing, clone enrichment using specific probes, and protocols similar or identical to 5' and 3' "RACE" protocols which are well known in the art. For instance, a method similar to 5' RACE is available for generating the missing 5' end of a desired full-length transcript. (Fromont-Racine et al., Nucleic Acids Res. 21(7):1683-1684 (1993).)

Briefly, a specific RNA oligonucleotide is ligated to the 5' ends of a population of RNA presumably containing full-length gene RNA transcripts. A primer set containing a primer specific to the ligated RNA oligonucleotide and a primer specific to a known sequence of the gene of interest is used to PCR amplify the 5' portion of the desired full-length gene. This amplified product may then be sequenced and used to generate the full length gene.

This above method starts with total RNA isolated from the desired source, although poly-A⁺ RNA can be used. The RNA preparation can then be treated with phosphatase if necessary to eliminate 5' phosphate groups on degraded or damaged RNA which may interfere with the later RNA ligase step. The phosphatase should then be inactivated and the RNA treated with tobacco acid pyrophosphatase in order to remove the cap structure present at the 5' ends of messenger RNAs. This reaction leaves a 5' phosphate group at the 5' end of the cap cleaved RNA which can then be ligated to an RNA oligonucleotide using T4 RNA ligase.

This modified RNA preparation is used as a template for first strand cDNA synthesis using a gene specific oligonucleotide. The first strand synthesis reaction is

used as a template for PCR amplification of the desired 5' end using a primer specific to the ligated RNA oligonucleotide and a primer specific to the known sequence of the gene of interest. The resultant product is then sequenced and analyzed to confirm that the 5' end sequence belongs to the desired gene.

5

Example 2: Isolation of Genomic Clones Corresponding to a Polynucleotide

A human genomic P1 library (Genomic Systems, Inc.) is screened by PCR using primers selected for the cDNA sequence corresponding to SEQ ID NO:X.,
10 according to the method described in Example 1. (See also, Sambrook.)

Example 3: Tissue Distribution of Polypeptide

Tissue distribution of mRNA expression of polynucleotides of the present invention is determined using protocols for Northern blot analysis, described by,
15 among others, Sambrook et al. For example, a cDNA probe produced by the method described in Example 1 is labeled with P³² using the rediprime™ DNA labeling system (Amersham Life Science), according to manufacturer's instructions. After labeling, the probe is purified using CHROMA SPIN-100™ column (Clontech Laboratories, Inc.), according to manufacturer's protocol number PT1200-1. The purified labeled probe is
20 then used to examine various human tissues for mRNA expression.

Multiple Tissue Northern (MTN) blots containing various human tissues (H) or human immune system tissues (IM) (Clontech) are examined with the labeled probe using ExpressHyb™ hybridization solution (Clontech) according to manufacturer's protocol number PT1190-1. Following hybridization and washing, the blots are
25 mounted and exposed to film at -70°C overnight, and the films developed according to standard procedures.

Example 4: Chromosomal Mapping of the Polynucleotides

An oligonucleotide primer set is designed according to the sequence at the 5'
30 end of SEQ ID NO:X. This primer preferably spans about 100 nucleotides. This primer set is then used in a polymerase chain reaction under the following set of conditions : 30 seconds, 95°C; 1 minute, 56°C; 1 minute, 70°C. This cycle is repeated
32 times followed by one 5 minute cycle at 70°C. Human, mouse, and hamster DNA is used as template in addition to a somatic cell hybrid panel containing individual
35 chromosomes or chromosome fragments (Bios, Inc). The reactions is analyzed on

either 8% polyacrylamide gels or 3.5 % agarose gels. Chromosome mapping is determined by the presence of an approximately 100 bp PCR fragment in the particular somatic cell hybrid.

5 **Example 5: Bacterial Expression of a Polypeptide**

A polynucleotide encoding a polypeptide of the present invention is amplified using PCR oligonucleotide primers corresponding to the 5' and 3' ends of the DNA sequence, as outlined in Example 1, to synthesize insertion fragments. The primers used to amplify the cDNA insert should preferably contain restriction sites, such as
10 BamHI and XbaI, at the 5' end of the primers in order to clone the amplified product into the expression vector. For example, BamHI and XbaI correspond to the restriction enzyme sites on the bacterial expression vector pQE-9. (Qiagen, Inc., Chatsworth, CA). This plasmid vector encodes antibiotic resistance (Amp^r), a bacterial origin of replication (ori), an IPTG-regulatable promoter/operator (P/O), a ribosome binding site
15 (RBS), a 6-histidine tag (6-His), and restriction enzyme cloning sites.

The pQE-9 vector is digested with BamHI and XbaI and the amplified fragment is ligated into the pQE-9 vector maintaining the reading frame initiated at the bacterial RBS. The ligation mixture is then used to transform the E. coli strain M15/rep4 (Qiagen, Inc.) which contains multiple copies of the plasmid pREP4, which expresses
20 the lacI repressor and also confers kanamycin resistance (Kan^r). Transformants are identified by their ability to grow on LB plates and ampicillin/kanamycin resistant colonies are selected. Plasmid DNA is isolated and confirmed by restriction analysis.

Clones containing the desired constructs are grown overnight (O/N) in liquid culture in LB media supplemented with both Amp (100 ug/ml) and Kan (25 ug/ml).
25 The O/N culture is used to inoculate a large culture at a ratio of 1:100 to 1:250. The cells are grown to an optical density 600 (O.D.⁶⁰⁰) of between 0.4 and 0.6. IPTG (Isopropyl-B-D-thiogalacto pyranoside) is then added to a final concentration of 1 mM. IPTG induces by inactivating the lacI repressor, clearing the P/O leading to increased gene expression.

30 Cells are grown for an extra 3 to 4 hours. Cells are then harvested by centrifugation (20 mins at 6000Xg). The cell pellet is solubilized in the chaotropic agent 6 Molar Guanidine HCl by stirring for 3-4 hours at 4°C. The cell debris is removed by centrifugation, and the supernatant containing the polypeptide is loaded onto a nickel-nitrilo-tri-acetic acid ("Ni-NTA") affinity resin column (available from
35 QIAGEN, Inc., *supra*). Proteins with a 6 x His tag bind to the Ni-NTA resin with high

affinity and can be purified in a simple one-step procedure (for details see: The QIAexpressionist (1995) QIAGEN, Inc., *supra*).

Briefly, the supernatant is loaded onto the column in 6 M guanidine-HCl, pH 8, the column is first washed with 10 volumes of 6 M guanidine-HCl, pH 8, then washed
5 with 10 volumes of 6 M guanidine-HCl pH 6, and finally the polypeptide is eluted with 6 M guanidine-HCl, pH 5.

The purified protein is then renatured by dialyzing it against phosphate-buffered saline (PBS) or 50 mM Na-acetate, pH 6 buffer plus 200 mM NaCl. Alternatively, the protein can be successfully refolded while immobilized on the Ni-NTA column. The
10 recommended conditions are as follows: renature using a linear 6M-1M urea gradient in 500 mM NaCl, 20% glycerol, 20 mM Tris/HCl pH 7.4, containing protease inhibitors. The renaturation should be performed over a period of 1.5 hours or more. After renaturation the proteins are eluted by the addition of 250 mM imidazole. Imidazole is removed by a final dialyzing step against PBS or 50 mM sodium acetate pH 6 buffer
15 plus 200 mM NaCl. The purified protein is stored at 4° C or frozen at -80° C.

In addition to the above expression vector, the present invention further includes an expression vector comprising phage operator and promoter elements operatively linked to a polynucleotide of the present invention, called pHE4a. (ATCC Accession Number 209645, deposited on February 25, 1998.) This vector contains: 1) a
20 neomycinphosphotransferase gene as a selection marker, 2) an E. coli origin of replication, 3) a T5 phage promoter sequence, 4) two lac operator sequences, 5) a Shine-Delgarno sequence, and 6) the lactose operon repressor gene (*lacIq*). The origin of replication (*oriC*) is derived from pUC19 (LTI, Gaithersburg, MD). The promoter sequence and operator sequences are made synthetically.

DNA can be inserted into the pHEa by restricting the vector with NdeI and
25 XbaI, BamHI, XhoI, or Asp718, running the restricted product on a gel, and isolating the larger fragment (the stuffer fragment should be about 310 base pairs). The DNA insert is generated according to the PCR protocol described in Example 1, using PCR primers having restriction sites for NdeI (5' primer) and XbaI, BamHI, XhoI, or
30 Asp718 (3' primer). The PCR insert is gel purified and restricted with compatible enzymes. The insert and vector are ligated according to standard protocols.

The engineered vector could easily be substituted in the above protocol to express protein in a bacterial system.

35 **Example 6: Purification of a Polypeptide from an Inclusion Body**

The following alternative method can be used to purify a polypeptide expressed in *E. coli* when it is present in the form of inclusion bodies. Unless otherwise specified, all of the following steps are conducted at 4-10°C.

Upon completion of the production phase of the *E. coli* fermentation, the cell
5 culture is cooled to 4-10°C and the cells harvested by continuous centrifugation at 15,000 rpm (Heraeus Sepatech). On the basis of the expected yield of protein per unit weight of cell paste and the amount of purified protein required, an appropriate amount of cell paste, by weight, is suspended in a buffer solution containing 100 mM Tris, 50 mM EDTA, pH 7.4. The cells are dispersed to a homogeneous suspension using a
10 high shear mixer.

The cells are then lysed by passing the solution through a microfluidizer (Microfluidics, Corp. or APV Gaulin, Inc.) twice at 4000-6000 psi. The homogenate is then mixed with NaCl solution to a final concentration of 0.5 M NaCl, followed by centrifugation at 7000 xg for 15 min. The resultant pellet is washed again using 0.5M
15 NaCl, 100 mM Tris, 50 mM EDTA, pH 7.4.

The resulting washed inclusion bodies are solubilized with 1.5 M guanidine hydrochloride (GuHCl) for 2-4 hours. After 7000 xg centrifugation for 15 min., the pellet is discarded and the polypeptide containing supernatant is incubated at 4°C overnight to allow further GuHCl extraction.

20 Following high speed centrifugation (30,000 xg) to remove insoluble particles, the GuHCl solubilized protein is refolded by quickly mixing the GuHCl extract with 20 volumes of buffer containing 50 mM sodium, pH 4.5, 150 mM NaCl, 2 mM EDTA by vigorous stirring. The refolded diluted protein solution is kept at 4°C without mixing for 12 hours prior to further purification steps.

25 To clarify the refolded polypeptide solution, a previously prepared tangential filtration unit equipped with 0.16 µm membrane filter with appropriate surface area (e.g., Filtron), equilibrated with 40 mM sodium acetate, pH 6.0 is employed. The filtered sample is loaded onto a cation exchange resin (e.g., Poros HS-50, Perseptive Biosystems). The column is washed with 40 mM sodium acetate, pH 6.0 and eluted
30 with 250 mM, 500 mM, 1000 mM, and 1500 mM NaCl in the same buffer, in a stepwise manner. The absorbance at 280 nm of the effluent is continuously monitored. Fractions are collected and further analyzed by SDS-PAGE.

Fractions containing the polypeptide are then pooled and mixed with 4 volumes of water. The diluted sample is then loaded onto a previously prepared set of tandem

columns of strong anion (Poros HQ-50, Perseptive Biosystems) and weak anion (Poros CM-20, Perseptive Biosystems) exchange resins. The columns are equilibrated with 40 mM sodium acetate, pH 6.0. Both columns are washed with 40 mM sodium acetate, pH 6.0, 200 mM NaCl. The CM-20 column is then eluted using a 10 column
5 volume linear gradient ranging from 0.2 M NaCl, 50 mM sodium acetate, pH 6.0 to 1.0 M NaCl, 50 mM sodium acetate, pH 6.5. Fractions are collected under constant A_{280} monitoring of the effluent. Fractions containing the polypeptide (determined, for instance, by 16% SDS-PAGE) are then pooled.

The resultant polypeptide should exhibit greater than 95% purity after the above
10 refolding and purification steps. No major contaminant bands should be observed from Commassie blue stained 16% SDS-PAGE gel when 5 μ g of purified protein is loaded. The purified protein can also be tested for endotoxin/LPS contamination, and typically the LPS content is less than 0.1 ng/ml according to LAL assays.

15 Example 7: Cloning and Expression of a Polypeptide in a Baculovirus Expression System

In this example, the plasmid shuttle vector pA2 is used to insert a polynucleotide into a baculovirus to express a polypeptide. This expression vector contains the strong polyhedrin promoter of the *Autographa californica* nuclear polyhedrosis virus
20 (AcMNPV) followed by convenient restriction sites such as BamHI, Xba I and Asp718. The polyadenylation site of the simian virus 40 ("SV40") is used for efficient polyadenylation. For easy selection of recombinant virus, the plasmid contains the beta-galactosidase gene from *E. coli* under control of a weak *Drosophila* promoter in the same orientation, followed by the polyadenylation signal of the polyhedrin gene. The
25 inserted genes are flanked on both sides by viral sequences for cell-mediated homologous recombination with wild-type viral DNA to generate a viable virus that express the cloned polynucleotide.

Many other baculovirus vectors can be used in place of the vector above, such as pAc373, pVL941, and pAcIM1, as one skilled in the art would readily appreciate, as
30 long as the construct provides appropriately located signals for transcription, translation, secretion and the like, including a signal peptide and an in-frame AUG as required. Such vectors are described, for instance, in Luckow et al., *Virology* 170:31-39 (1989).

Specifically, the cDNA sequence contained in the deposited clone, including the
35 AUG initiation codon and the naturally associated leader sequence identified in Table 1, is amplified using the PCR protocol described in Example 1. If the naturally occurring

signal sequence is used to produce the secreted protein, the pA2 vector does not need a second signal peptide. Alternatively, the vector can be modified (pA2 GP) to include a baculovirus leader sequence, using the standard methods described in Summers et al., "A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures,"

5 Texas Agricultural Experimental Station Bulletin No. 1555 (1987).

The amplified fragment is isolated from a 1% agarose gel using a commercially available kit ("GeneClean," BIO 101 Inc., La Jolla, Ca.). The fragment then is digested with appropriate restriction enzymes and again purified on a 1% agarose gel.

10 The plasmid is digested with the corresponding restriction enzymes and optionally, can be dephosphorylated using calf intestinal phosphatase, using routine procedures known in the art. The DNA is then isolated from a 1% agarose gel using a commercially available kit ("GeneClean" BIO 101 Inc., La Jolla, Ca.).

The fragment and the dephosphorylated plasmid are ligated together with T4 DNA ligase. *E. coli* HB101 or other suitable *E. coli* hosts such as XL-1 Blue
15 (Stratagene Cloning Systems, La Jolla, CA) cells are transformed with the ligation mixture and spread on culture plates. Bacteria containing the plasmid are identified by digesting DNA from individual colonies and analyzing the digestion product by gel electrophoresis. The sequence of the cloned fragment is confirmed by DNA sequencing.

20 Five µg of a plasmid containing the polynucleotide is co-transfected with 1.0 µg of a commercially available linearized baculovirus DNA ("BaculoGold™ baculovirus DNA", Pharmingen, San Diego, CA), using the lipofection method described by Felgner et al., Proc. Natl. Acad. Sci. USA 84:7413-7417 (1987). One µg of BaculoGold™ virus DNA and 5 µg of the plasmid are mixed in a sterile well of a
25 microtiter plate containing 50 µl of serum-free Grace's medium (Life Technologies Inc., Gaithersburg, MD). Afterwards, 10 µl Lipofectin plus 90 µl Grace's medium are added, mixed and incubated for 15 minutes at room temperature. Then the transfection mixture is added drop-wise to Sf9 insect cells (ATCC CRL 1711) seeded in a 35 mm tissue culture plate with 1 ml Grace's medium without serum. The plate is then
30 incubated for 5 hours at 27° C. The transfection solution is then removed from the plate and 1 ml of Grace's insect medium supplemented with 10% fetal calf serum is added. Cultivation is then continued at 27° C for four days.

After four days the supernatant is collected and a plaque assay is performed, as described by Summers and Smith, *supra*. An agarose gel with "Blue Gal" (Life
35 Technologies Inc., Gaithersburg) is used to allow easy identification and isolation of gal-expressing clones, which produce blue-stained plaques. (A detailed description of a "plaque assay" of this type can also be found in the user's guide for insect cell culture

and baculovirology distributed by Life Technologies Inc., Gaithersburg, page 9-10.) After appropriate incubation, blue stained plaques are picked with the tip of a micropipettor (e.g., Eppendorf). The agar containing the recombinant viruses is then resuspended in a microcentrifuge tube containing 200 µl of Grace's medium and the suspension containing the recombinant baculovirus is used to infect Sf9 cells seeded in 35 mm dishes. Four days later the supernatants of these culture dishes are harvested and then they are stored at 4° C.

To verify the expression of the polypeptide, Sf9 cells are grown in Grace's medium supplemented with 10% heat-inactivated FBS. The cells are infected with the recombinant baculovirus containing the polynucleotide at a multiplicity of infection ("MOI") of about 2. If radiolabeled proteins are desired, 6 hours later the medium is removed and is replaced with SF900 II medium minus methionine and cysteine (available from Life Technologies Inc., Rockville, MD). After 42 hours, 5 µCi of ³⁵S-methionine and 5 µCi ³⁵S-cysteine (available from Amersham) are added. The cells are further incubated for 16 hours and then are harvested by centrifugation. The proteins in the supernatant as well as the intracellular proteins are analyzed by SDS-PAGE followed by autoradiography (if radiolabeled).

Microsequencing of the amino acid sequence of the amino terminus of purified protein may be used to determine the amino terminal sequence of the produced protein.

Example 8: Expression of a Polypeptide in Mammalian Cells

The polypeptide of the present invention can be expressed in a mammalian cell. A typical mammalian expression vector contains a promoter element, which mediates the initiation of transcription of mRNA, a protein coding sequence, and signals required for the termination of transcription and polyadenylation of the transcript. Additional elements include enhancers, Kozak sequences and intervening sequences flanked by donor and acceptor sites for RNA splicing. Highly efficient transcription is achieved with the early and late promoters from SV40, the long terminal repeats (LTRs) from Retroviruses, e.g., RSV, HTLV, HIV and the early promoter of the cytomegalovirus (CMV). However, cellular elements can also be used (e.g., the human actin promoter).

Suitable expression vectors for use in practicing the present invention include, for example, vectors such as pSVL and pMSG (Pharmacia, Uppsala, Sweden), pRSVcat (ATCC 37152), pSV2dhfr (ATCC 37146), pBC12MI (ATCC 67109), pCMVSPORT 2.0, and pCMVSPORT 3.0. Mammalian host cells that could be used include, human HeLa, 293, H9 and Jurkat cells, mouse NIH3T3 and C127 cells, Cos 1, Cos 7 and CV1, quail QC1-3 cells, mouse L cells and Chinese hamster ovary (CHO) cells.

Alternatively, the polypeptide can be expressed in stable cell lines containing the polynucleotide integrated into a chromosome. The co-transfection with a selectable marker such as dhfr, gpt, neomycin, hygromycin allows the identification and isolation of the transfected cells.

5 The transfected gene can also be amplified to express large amounts of the encoded protein. The DHFR (dihydrofolate reductase) marker is useful in developing cell lines that carry several hundred or even several thousand copies of the gene of interest. (See, e.g., Alt, F. W., et al., J. Biol. Chem. 253:1357-1370 (1978); Hamlin, J. L. and Ma, C., Biochem. et Biophys. Acta, 1097:107-143 (1990); Page, M. J. and
10 Sydenham, M. A., Biotechnology 9:64-68 (1991).) Another useful selection marker is the enzyme glutamine synthase (GS) (Murphy et al., Biochem J. 227:277-279 (1991); Bebbington et al., Bio/Technology 10:169-175 (1992). Using these markers, the mammalian cells are grown in selective medium and the cells with the highest resistance are selected. These cell lines contain the amplified gene(s) integrated into a
15 chromosome. Chinese hamster ovary (CHO) and NSO cells are often used for the production of proteins.

Derivatives of the plasmid pSV2-dhfr (ATCC Accession No. 37146), the expression vectors pC4 (ATCC Accession No. 209646) and pC6 (ATCC Accession No.209647) contain the strong promoter (LTR) of the Rous Sarcoma Virus (Cullen et
20 al., Molecular and Cellular Biology, 438-447 (March, 1985)) plus a fragment of the CMV-enhancer (Boshart et al., Cell 41:521-530 (1985).) Multiple cloning sites, e.g., with the restriction enzyme cleavage sites BamHI, XbaI and Asp718, facilitate the cloning of the gene of interest. The vectors also contain the 3' intron, the polyadenylation and termination signal of the rat preproinsulin gene, and the mouse
25 DHFR gene under control of the SV40 early promoter.

Specifically, the plasmid pC6, for example, is digested with appropriate restriction enzymes and then dephosphorylated using calf intestinal phosphates by procedures known in the art. The vector is then isolated from a 1% agarose gel.

A polynucleotide of the present invention is amplified according to the protocol
30 outlined in Example 1. If the naturally occurring signal sequence is used to produce the secreted protein, the vector does not need a second signal peptide. Alternatively, if the naturally occurring signal sequence is not used, the vector can be modified to include a heterologous signal sequence. (See, e.g., WO 96/34891.)

The amplified fragment is isolated from a 1% agarose gel using a commercially
35 available kit ("GeneClean," BIO 101 Inc., La Jolla, Ca.). The fragment then is digested with appropriate restriction enzymes and again purified on a 1% agarose gel.

The amplified fragment is then digested with the same restriction enzyme and purified on a 1% agarose gel. The isolated fragment and the dephosphorylated vector are then ligated with T4 DNA ligase. *E. coli* HB101 or XL-1 Blue cells are then transformed and bacteria are identified that contain the fragment inserted into plasmid pC6 using, for instance, restriction enzyme analysis.

Chinese hamster ovary cells lacking an active DHFR gene is used for transfection. Five μ g of the expression plasmid pC6 is cotransfected with 0.5 μ g of the plasmid pSVneo using lipofectin (Felgner et al., *supra*). The plasmid pSV2-neo contains a dominant selectable marker, the *neo* gene from Tn5 encoding an enzyme that confers resistance to a group of antibiotics including G418. The cells are seeded in alpha minus MEM supplemented with 1 mg/ml G418. After 2 days, the cells are trypsinized and seeded in hybridoma cloning plates (Greiner, Germany) in alpha minus MEM supplemented with 10, 25, or 50 ng/ml of methotrexate plus 1 mg/ml G418. After about 10-14 days single clones are trypsinized and then seeded in 6-well petri dishes or 10 ml flasks using different concentrations of methotrexate (50 nM, 100 nM, 200 nM, 400 nM, 800 nM). Clones growing at the highest concentrations of methotrexate are then transferred to new 6-well plates containing even higher concentrations of methotrexate (1 μ M, 2 μ M, 5 μ M, 10 mM, 20 mM). The same procedure is repeated until clones are obtained which grow at a concentration of 100 - 200 μ M. Expression of the desired gene product is analyzed, for instance, by SDS-PAGE and Western blot or by reversed phase HPLC analysis.

Example 9: Protein Fusions

The polypeptides of the present invention are preferably fused to other proteins. These fusion proteins can be used for a variety of applications. For example, fusion of the present polypeptides to His-tag, HA-tag, protein A, IgG domains, and maltose binding protein facilitates purification. (See Example 5; see also EP A 394,827; Traunecker, et al., Nature 331:84-86 (1988).) Similarly, fusion to IgG-1, IgG-3, and albumin increases the halflife time in vivo. Nuclear localization signals fused to the polypeptides of the present invention can target the protein to a specific subcellular localization, while covalent heterodimer or homodimers can increase or decrease the activity of a fusion protein. Fusion proteins can also create chimeric molecules having more than one function. Finally, fusion proteins can increase solubility and/or stability of the fused protein compared to the non-fused protein. All of the types of fusion proteins described above can be made by modifying the following protocol, which outlines the fusion of a polypeptide to an IgG molecule, or the protocol described in Example 5.

Briefly, the human Fc portion of the IgG molecule can be PCR amplified, using primers that span the 5' and 3' ends of the sequence described below. These primers also should have convenient restriction enzyme sites that will facilitate cloning into an expression vector, preferably a mammalian expression vector.

5 For example, if pC4 (Accession No. 209646) is used, the human Fc portion can be ligated into the BamHI cloning site. Note that the 3' BamHI site should be destroyed. Next, the vector containing the human Fc portion is re-restricted with BamHI, linearizing the vector, and a polynucleotide of the present invention, isolated by the PCR protocol described in Example 1, is ligated into this BamHI site. Note that
10 the polynucleotide is cloned without a stop codon, otherwise a fusion protein will not be produced.

If the naturally occurring signal sequence is used to produce the secreted protein, pC4 does not need a second signal peptide. Alternatively, if the naturally occurring signal sequence is not used, the vector can be modified to include a
15 heterologous signal sequence. (See, e.g., WO 96/34891.)

Human IgG Fc region:

GGGATCCGGAGCCCAAATCTTCTGACAAACTCACACATGCCCACCGTGCC
CAGCACCTGAATTTCGAGGGTGCACCGTCAGTCTTCCTCTTCCCCCAAAC
20 CAAGGACACCCTCATGATCTCCCGGACTCCTGAGGTCACATGCGTGGTGGT
GGACGTAAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACG
GCGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAAC
AGCACGTACCGTGTGGTTCAGCGTCCTCACCGTCCTGCACCAGGACTGGCTG
AATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCCCAACCCCC
25 ATCGAGAAAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGAACCACAGGT
GTACACCCTGCCCCCATCCCGGGATGAGCTGACCAAGAACCAGGTCAGCCT
GACCTGCCTGGTCAAAGGCTTCTATCCAAGCGACATCGCCGTGGAGTGGGA
GAGCAATGGGCAGCCGGAGAACAACACTACAAGACCACGCCTCCCGTGCTGG
ACTCCGACGGCTCCTTCTTCCTCTACAGCAAGCTCACCGTGGACAAGAGCA
30 GGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGC
ACAACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAATGAGTGC
GACGGCCGCGACTCTAGAGGAT (SEQ ID NO:1)

Example 10: Production of an Antibody from a Polypeptide

35 The antibodies of the present invention can be prepared by a variety of methods. (See, Current Protocols, Chapter 2.) For example, cells expressing a polypeptide of the present invention is administered to an animal to induce the production of sera

containing polyclonal antibodies. In a preferred method, a preparation of the secreted protein is prepared and purified to render it substantially free of natural contaminants. Such a preparation is then introduced into an animal in order to produce polyclonal antisera of greater specific activity.

5 In the most preferred method, the antibodies of the present invention are monoclonal antibodies (or protein binding fragments thereof). Such monoclonal antibodies can be prepared using hybridoma technology. (Köhler et al., *Nature* 256:495 (1975); Köhler et al., *Eur. J. Immunol.* 6:511 (1976); Köhler et al., *Eur. J. Immunol.* 6:292 (1976); Hammerling et al., in: *Monoclonal Antibodies and T-Cell*
10 *Hybridomas*, Elsevier, N.Y., pp. 563-681 (1981).) In general, such procedures involve immunizing an animal (preferably a mouse) with polypeptide or, more preferably, with a secreted polypeptide-expressing cell. Such cells may be cultured in any suitable tissue culture medium; however, it is preferable to culture cells in Earle's modified Eagle's medium supplemented with 10% fetal bovine serum (inactivated at
15 about 56°C), and supplemented with about 10 g/l of nonessential amino acids, about 1,000 U/ml of penicillin, and about 100 µg/ml of streptomycin.

 The splenocytes of such mice are extracted and fused with a suitable myeloma cell line. Any suitable myeloma cell line may be employed in accordance with the present invention; however, it is preferable to employ the parent myeloma cell line
20 (SP2O), available from the ATCC. After fusion, the resulting hybridoma cells are selectively maintained in HAT medium, and then cloned by limiting dilution as described by Wands et al. (*Gastroenterology* 80:225-232 (1981).) The hybridoma cells obtained through such a selection are then assayed to identify clones which secrete antibodies capable of binding the polypeptide.

25 Alternatively, additional antibodies capable of binding to the polypeptide can be produced in a two-step procedure using anti-idiotypic antibodies. Such a method makes use of the fact that antibodies are themselves antigens, and therefore, it is possible to obtain an antibody which binds to a second antibody. In accordance with this method, protein specific antibodies are used to immunize an animal, preferably a
30 mouse. The splenocytes of such an animal are then used to produce hybridoma cells, and the hybridoma cells are screened to identify clones which produce an antibody whose ability to bind to the protein-specific antibody can be blocked by the polypeptide. Such antibodies comprise anti-idiotypic antibodies to the protein-specific antibody and can be used to immunize an animal to induce formation of further protein-specific
35 antibodies.

It will be appreciated that Fab and F(ab')₂ and other fragments of the antibodies of the present invention may be used according to the methods disclosed herein. Such fragments are typically produced by proteolytic cleavage, using enzymes such as papain (to produce Fab fragments) or pepsin (to produce F(ab')₂ fragments). Alternatively, 5 secreted protein-binding fragments can be produced through the application of recombinant DNA technology or through synthetic chemistry.

For in vivo use of antibodies in humans, it may be preferable to use "humanized" chimeric monoclonal antibodies. Such antibodies can be produced using genetic constructs derived from hybridoma cells producing the monoclonal antibodies 10 described above. Methods for producing chimeric antibodies are known in the art. (See, for review, Morrison, Science 229:1202 (1985); Oi et al., BioTechniques 4:214 (1986); Cabilly et al., U.S. Patent No. 4,816,567; Taniguchi et al., EP 171496; Morrison et al., EP 173494; Neuberger et al., WO 8601533; Robinson et al., WO 8702671; Boulianne et al., Nature 312:643 (1984); Neuberger et al., Nature 314:268 15 (1985).)

Example 11: Production Of Secreted Protein For High-Throughput Screening Assays

The following protocol produces a supernatant containing a polypeptide to be 20 tested. This supernatant can then be used in the Screening Assays described in Examples 13-20.

First, dilute Poly-D-Lysine (644 587 Boehringer-Mannheim) stock solution (1mg/ml in PBS) 1:20 in PBS (w/o calcium or magnesium 17-516F Biowhittaker) for a working solution of 50ug/ml. Add 200 ul of this solution to each well (24 well plates) 25 and incubate at RT for 20 minutes. Be sure to distribute the solution over each well (note: a 12-channel pipetter may be used with tips on every other channel). Aspirate off the Poly-D-Lysine solution and rinse with 1ml PBS (Phosphate Buffered Saline). The PBS should remain in the well until just prior to plating the cells and plates may be poly-lysine coated in advance for up to two weeks.

30 Plate 293T cells (do not carry cells past P+20) at 2×10^5 cells/well in .5ml DMEM(Dulbecco's Modified Eagle Medium)(with 4.5 G/L glucose and L-glutamine (12-604F Biowhittaker))/10% heat inactivated FBS(14-503F Biowhittaker)/1x Penstrep(17-602E Biowhittaker). Let the cells grow overnight.

The next day, mix together in a sterile solution basin: 300 ul Lipofectamine 35 (18324-012 Gibco/BRL) and 5ml Optimem I (31985070 Gibco/BRL)/96-well plate. With a small volume multi-channel pipetter, aliquot approximately 2ug of an expression vector containing a polynucleotide insert, produced by the methods described in

Examples 8 or 9, into an appropriately labeled 96-well round bottom plate. With a multi-channel pipetter, add 50ul of the Lipofectamine/Optimem I mixture to each well. Pipette up and down gently to mix. Incubate at RT 15-45 minutes. After about 20 minutes, use a multi-channel pipetter to add 150ul Optimem I to each well. As a control, one plate of vector DNA lacking an insert should be transfected with each set of transfections.

Preferably, the transfection should be performed by tag-teaming the following tasks. By tag-teaming, hands on time is cut in half, and the cells do not spend too much time on PBS. First, person A aspirates off the media from four 24-well plates of cells, and then person B rinses each well with .5-1ml PBS. Person A then aspirates off PBS rinse, and person B, using a 12-channel pipetter with tips on every other channel, adds the 200ul of DNA/Lipofectamine/Optimem I complex to the odd wells first, then to the even wells, to each row on the 24-well plates. Incubate at 37°C for 6 hours.

While cells are incubating, prepare appropriate media, either 1%BSA in DMEM with 1x penstrep, or CHO-5 media (116.6 mg/L of CaCl₂ (anhyd); 0.00130 mg/L CuSO₄·5H₂O; 0.050 mg/L of Fe(NO₃)₃·9H₂O; 0.417 mg/L of FeSO₄·7H₂O; 311.80 mg/L of KCl; 28.64 mg/L of MgCl₂; 48.84 mg/L of MgSO₄; 6995.50 mg/L of NaCl; 2400.0 mg/L of NaHCO₃; 62.50 mg/L of NaH₂PO₄·H₂O; 71.02 mg/L of Na₂HPO₄; .4320 mg/L of ZnSO₄·7H₂O; .002 mg/L of Arachidonic Acid; 1.022 mg/L of Cholesterol; .070 mg/L of DL-alpha-Tocopherol-Acetate; 0.0520 mg/L of Linoleic Acid; 0.010 mg/L of Linolenic Acid; 0.010 mg/L of Myristic Acid; 0.010 mg/L of Oleic Acid; 0.010 mg/L of Palmitic Acid; 0.010 mg/L of Palmitic Acid; 100 mg/L of Pluronic F-68; 0.010 mg/L of Stearic Acid; 2.20 mg/L of Tween 80; 4551 mg/L of D-Glucose; 130.85 mg/ml of L- Alanine; 147.50 mg/ml of L-Arginine-HCL; 7.50 mg/ml of L-Asparagine-H₂O; 6.65 mg/ml of L-Aspartic Acid; 29.56 mg/ml of L-Cystine-2HCL-H₂O; 31.29 mg/ml of L-Cystine-2HCL; 7.35 mg/ml of L-Glutamic Acid; 365.0 mg/ml of L-Glutamine; 18.75 mg/ml of Glycine; 52.48 mg/ml of L-Histidine-HCL-H₂O; 106.97 mg/ml of L-Isoleucine; 111.45 mg/ml of L-Leucine; 163.75 mg/ml of L-Lysine HCL; 32.34 mg/ml of L-Methionine; 68.48 mg/ml of L-Phenylalanine; 40.0 mg/ml of L-Proline; 26.25 mg/ml of L-Serine; 101.05 mg/ml of L-Threonine; 19.22 mg/ml of L-Tryptophan; 91.79 mg/ml of L-Tyrosine-2Na-2H₂O; 99.65 mg/ml of L-Valine; 0.0035 mg/L of Biotin; 3.24 mg/L of D-Ca Pantothenate; 11.78 mg/L of Choline Chloride; 4.65 mg/L of Folic Acid; 15.60 mg/L of i-Inositol; 3.02 mg/L of Niacinamide; 3.00 mg/L of Pyridoxal HCL; 0.031 mg/L of Pyridoxine HCL; 0.319 mg/L of Riboflavin; 3.17 mg/L of Thiamine HCL; 0.365 mg/L of Thymidine; and 0.680 mg/L of Vitamin B₁₂; 25 mM of HEPES Buffer; 2.39 mg/L of Na Hypoxanthine;

0.105 mg/L of Lipoic Acid; 0.081 mg/L of Sodium Putrescine-2HCL; 55.0 mg/L of Sodium Pyruvate; 0.0067 mg/L of Sodium Selenite; 20uM of Ethanolamine; 0.122 mg/L of Ferric Citrate; 41.70 mg/L of Methyl-B-Cyclodextrin complexed with Linoleic Acid; 33.33 mg/L of Methyl-B-Cyclodextrin complexed with Oleic Acid; and 10 mg/L of Methyl-B-Cyclodextrin complexed with Retinal) with 2mm glutamine and 1x penstrep. (BSA (81-068-3 Bayer) 100gm dissolved in 1L DMEM for a 10% BSA stock solution). Filter the media and collect 50 ul for endotoxin assay in 15ml polystyrene conical.

The transfection reaction is terminated, preferably by tag-teaming, at the end of the incubation period. Person A aspirates off the transfection media, while person B adds 1.5ml appropriate media to each well. Incubate at 37°C for 45 or 72 hours depending on the media used: 1%BSA for 45 hours or CHO-5 for 72 hours.

On day four, using a 300ul multichannel pipetter, aliquot 600ul in one 1ml deep well plate and the remaining supernatant into a 2ml deep well. The supernatants from each well can then be used in the assays described in Examples 13-20.

It is specifically understood that when activity is obtained in any of the assays described below using a supernatant, the activity originates from either the polypeptide directly (e.g., as a secreted protein) or by the polypeptide inducing expression of other proteins, which are then secreted into the supernatant. Thus, the invention further provides a method of identifying the protein in the supernatant characterized by an activity in a particular assay.

Example 12: Construction of GAS Reporter Construct

One signal transduction pathway involved in the differentiation and proliferation of cells is called the Jaks-STATs pathway. Activated proteins in the Jaks-STATs pathway bind to gamma activation site "GAS" elements or interferon-sensitive responsive element ("ISRE"), located in the promoter of many genes. The binding of a protein to these elements alter the expression of the associated gene.

GAS and ISRE elements are recognized by a class of transcription factors called Signal Transducers and Activators of Transcription, or "STATs." There are six members of the STATs family. Stat1 and Stat3 are present in many cell types, as is Stat2 (as response to IFN-alpha is widespread). Stat4 is more restricted and is not in many cell types though it has been found in T helper class I, cells after treatment with IL-12. Stat5 was originally called mammary growth factor, but has been found at higher concentrations in other cells including myeloid cells. It can be activated in tissue culture cells by many cytokines.

The STATs are activated to translocate from the cytoplasm to the nucleus upon tyrosine phosphorylation by a set of kinases known as the Janus Kinase ("Jaks") family. Jaks represent a distinct family of soluble tyrosine kinases and include Tyk2, Jak1, Jak2, and Jak3. These kinases display significant sequence similarity and are generally catalytically inactive in resting cells.

The Jaks are activated by a wide range of receptors summarized in the Table below. (Adapted from review by Schidler and Darnell, *Ann. Rev. Biochem.* 64:621-51 (1995).) A cytokine receptor family, capable of activating Jaks, is divided into two groups: (a) Class 1 includes receptors for IL-2, IL-3, IL-4, IL-6, IL-7, IL-9, IL-11, IL-12, IL-15, Epo, PRL, GH, G-CSF, GM-CSF, LIF, CNTF, and thrombopoietin; and (b) Class 2 includes IFN- α , IFN- γ , and IL-10. The Class 1 receptors share a conserved cysteine motif (a set of four conserved cysteines and one tryptophan) and a WSXWS motif (a membrane proximal region encoding Trp-Ser-Xxx-Trp-Ser (SEQ ID NO:2)).

Thus, on binding of a ligand to a receptor, Jaks are activated, which in turn activate STATs, which then translocate and bind to GAS elements. This entire process is encompassed in the Jaks-STATs signal transduction pathway.

Therefore, activation of the Jaks-STATs pathway, reflected by the binding of the GAS or the ISRE element, can be used to indicate proteins involved in the proliferation and differentiation of cells. For example, growth factors and cytokines are known to activate the Jaks-STATs pathway. (See Table below.) Thus, by using GAS elements linked to reporter molecules, activators of the Jaks-STATs pathway can be identified.

	Ligand	tyk2	JAKs		252 STATS	GAS(elements) or ISRE
			Jak1	Jak2	Jak3	
<u>IFN family</u>						
5	IFN-a/B	+	+	-	1,2,3	ISRE
	IFN-g		+	+	1	GAS (IRF1>Lys6>IFP)
	IL-10	+	?	?	- 1,3	
<u>gp130 family</u>						
10	IL-6 (Pleiotrohic)	+	+	+	? 1,3	GAS (IRF1>Lys6>IFP)
	IL-11(Pleiotrohic)	?	+	?	? 1,3	
	OnM(Pleiotrohic)	?	+	+	? 1,3	
	LIF(Pleiotrohic)	?	+	+	? 1,3	
	CNTF(Pleiotrohic)	-/+	+	+	? 1,3	
15	G-CSF(Pleiotrohic)	?	+	?	? 1,3	
	IL-12(Pleiotrohic)	+	-	+	+	1,3
<u>g-C family</u>						
20	IL-2 (lymphocytes)	-	+	-	+	1,3,5 GAS
	IL-4 (lymph/myeloid)	-	+	-	+	6 GAS (IRF1 = IFP >>Ly6)(IgH)
	IL-7 (lymphocytes)	-	+	-	+	5 GAS
	IL-9 (lymphocytes)	-	+	-	+	5 GAS
	IL-13 (lymphocyte)	-	+	?	?	6 GAS
	IL-15	?	+	?	+	5 GAS
25	<u>gp140 family</u>					
30	IL-3 (myeloid)	-	-	+	-	5 GAS (IRF1>IFP>>Ly6)
	IL-5 (myeloid)	-	-	+	-	5 GAS
	GM-CSF (myeloid)	-	-	+	-	5 GAS
<u>Growth hormone family</u>						
35	GH	?	-	+	-	5
	PRL	?	+/-	+	-	1,3,5
	EPO	?	-	+	-	5
	CAS>IRF1=IFP>>Ly6)					GAS(B-
<u>Receptor Tyrosine Kinases</u>						
40	EGF	?	+	+	-	1,3 GAS (IRF1)
	PDGF	?	+	+	-	1,3
	CSF-1	?	+	+	-	1,3 GAS (not IRF1)

To construct a synthetic GAS containing promoter element, which is used in the Biological Assays described in Examples 13-14, a PCR based strategy is employed to generate a GAS-SV40 promoter sequence. The 5' primer contains four tandem copies of the GAS binding site found in the IRF1 promoter and previously demonstrated to bind STATs upon induction with a range of cytokines (Rothman et al., Immunity 1:457-468 (1994).), although other GAS or ISRE elements can be used instead. The 5' primer also contains 18bp of sequence complementary to the SV40 early promoter sequence and is flanked with an XhoI site. The sequence of the 5' primer is:

5':GCGCCTCGAGATTTCCCCGAAATCTAGATTTCCCCGAAATGATTTCCCCG
AAATGATTTCCCCGAAATATCTGCCATCTCAATTAG:3' (SEQ ID NO:3)

The downstream primer is complementary to the SV40 promoter and is flanked with a Hind III site: 5':GCGGCAAGCTTTTGGCAAAGCCTAGGC:3' (SEQ ID NO:4)

PCR amplification is performed using the SV40 promoter template present in the B-gal:promoter plasmid obtained from Clontech. The resulting PCR fragment is digested with XhoI/Hind III and subcloned into BLSK2-. (Stratagene.) Sequencing with forward and reverse primers confirms that the insert contains the following sequence:

5':CTCGAGATTTCCCCGAAATCTAGATTTCCCCGAAATGATTTCCCCGAAATG
ATTTCCCCGAAATATCTGCCATCTCAATTAGTCAGCAACCATAGTCCCGCCC
CTAACTCCGCCCATCCCGCCCCTAACTCCGCCCAGTTCCGCCCATTCTCCGC
CCCATGGCTGACTAATTTTTTTTATTTATGCAGAGGCCGAGGCCGCCTCGGC
CTCTGAGCTATTCCAGAAGTAGTGAGGAGGCTTTTTTGGAGGCCTAGGCTTT
TGCAAAAAGCTT:3' (SEQ ID NO:5)

With this GAS promoter element linked to the SV40 promoter, a GAS:SEAP2 reporter construct is next engineered. Here, the reporter molecule is a secreted alkaline phosphatase, or "SEAP." Clearly, however, any reporter molecule can be instead of SEAP, in this or in any of the other Examples. Well known reporter molecules that can be used instead of SEAP include chloramphenicol acetyltransferase (CAT), luciferase, alkaline phosphatase, B-galactosidase, green fluorescent protein (GFP), or any protein detectable by an antibody.

The above sequence confirmed synthetic GAS-SV40 promoter element is subcloned into the pSEAP-Promoter vector obtained from Clontech using HindIII and XhoI, effectively replacing the SV40 promoter with the amplified GAS:SV40 promoter element, to create the GAS-SEAP vector. However, this vector does not contain a neomycin resistance gene, and therefore, is not preferred for mammalian expression systems.

Thus, in order to generate mammalian stable cell lines expressing the GAS-SEAP reporter, the GAS-SEAP cassette is removed from the GAS-SEAP vector using SalI and NotI, and inserted into a backbone vector containing the neomycin resistance gene, such as pGFP-1 (Clontech), using these restriction sites in the multiple cloning
5 site, to create the GAS-SEAP/Neo vector. Once this vector is transfected into mammalian cells, this vector can then be used as a reporter molecule for GAS binding as described in Examples 13-14.

Other constructs can be made using the above description and replacing GAS with a different promoter sequence. For example, construction of reporter molecules
10 containing NFK-B and EGR promoter sequences are described in Examples 15 and 16. However, many other promoters can be substituted using the protocols described in these Examples. For instance, SRE, IL-2, NFAT, or Osteocalcin promoters can be substituted, alone or in combination (e.g., GAS/NF-KB/EGR, GAS/NF-KB, IL-2/NFAT, or NF-KB/GAS). Similarly, other cell lines can be used to test reporter
15 construct activity, such as HELA (epithelial), HUVEC (endothelial), Reh (B-cell), Saos-2 (osteoblast), HUVAC (aortic), or Cardiomyocyte.

Example 13: High-Throughput Screening Assay for T-cell Activity.

The following protocol is used to assess T-cell activity by identifying factors,
20 such as growth factors and cytokines, that may proliferate or differentiate T-cells. T-cell activity is assessed using the GAS/SEAP/Neo construct produced in Example 12. Thus, factors that increase SEAP activity indicate the ability to activate the Jaks-STATS signal transduction pathway. The T-cell used in this assay is Jurkat T-cells (ATCC Accession No. TIB-152), although Molt-3 cells (ATCC Accession No. CRL-1552) and
25 Molt-4 cells (ATCC Accession No. CRL-1582) cells can also be used.

Jurkat T-cells are lymphoblastic CD4+ Th1 helper cells. In order to generate stable cell lines, approximately 2 million Jurkat cells are transfected with the GAS-SEAP/neo vector using DMRIE-C (Life Technologies)(transfection procedure described below). The transfected cells are seeded to a density of approximately
30 20,000 cells per well and transfectants resistant to 1 mg/ml gentamicin selected. Resistant colonies are expanded and then tested for their response to increasing concentrations of interferon gamma. The dose response of a selected clone is demonstrated.

Specifically, the following protocol will yield sufficient cells for 75 wells containing 200 ul of cells. Thus, it is either scaled up, or performed in multiple to
35 generate sufficient cells for multiple 96 well plates. Jurkat cells are maintained in RPMI + 10% serum with 1%Pen-Strep. Combine 2.5 mls of OPTI-MEM (Life Technologies)

with 10 ug of plasmid DNA in a T25 flask. Add 2.5 ml OPTI-MEM containing 50 ul of DMRIE-C and incubate at room temperature for 15-45 mins.

During the incubation period, count cell concentration, spin down the required number of cells (10^7 per transfection), and resuspend in OPTI-MEM to a final
5 concentration of 10^7 cells/ml. Then add 1ml of 1×10^7 cells in OPTI-MEM to T25 flask and incubate at 37°C for 6 hrs. After the incubation, add 10 ml of RPMI + 15% serum.

The Jurkat:GAS-SEAP stable reporter lines are maintained in RPMI + 10% serum, 1 mg/ml Genticin, and 1% Pen-Strep. These cells are treated with supernatants containing a polypeptide as produced by the protocol described in Example 11.

10 On the day of treatment with the supernatant, the cells should be washed and resuspended in fresh RPMI + 10% serum to a density of 500,000 cells per ml. The exact number of cells required will depend on the number of supernatants being screened. For one 96 well plate, approximately 10 million cells (for 10 plates, 100 million cells) are required.

15 Transfer the cells to a triangular reservoir boat, in order to dispense the cells into a 96 well dish, using a 12 channel pipette. Using a 12 channel pipette, transfer 200 ul of cells into each well (therefore adding 100,000 cells per well).

After all the plates have been seeded, 50 ul of the supernatants are transferred directly from the 96 well plate containing the supernatants into each well using a 12
20 channel pipette. In addition, a dose of exogenous interferon gamma (0.1, 1.0, 10 ng) is added to wells H9, H10, and H11 to serve as additional positive controls for the assay.

The 96 well dishes containing Jurkat cells treated with supernatants are placed in an incubator for 48 hrs (note: this time is variable between 48-72 hrs). 35 ul samples
25 from each well are then transferred to an opaque 96 well plate using a 12 channel pipette. The opaque plates should be covered (using sellophane covers) and stored at -20°C until SEAP assays are performed according to Example 17. The plates containing the remaining treated cells are placed at 4°C and serve as a source of material for repeating the assay on a specific well if desired.

30 As a positive control, 100 Unit/ml interferon gamma can be used which is known to activate Jurkat T cells. Over 30 fold induction is typically observed in the positive control wells.

Example 14: High-Throughput Screening Assay Identifying Myeloid

35 Activity

The following protocol is used to assess myeloid activity by identifying factors, such as growth factors and cytokines, that may proliferate or differentiate myeloid cells. Myeloid cell activity is assessed using the GAS/SEAP/Neo construct produced in Example 12. Thus, factors that increase SEAP activity indicate the ability to activate the Jaks-STATS signal transduction pathway. The myeloid cell used in this assay is U937, a pre-monocyte cell line, although TF-1, HL60, or KG1 can be used.

To transiently transfect U937 cells with the GAS/SEAP/Neo construct produced in Example 12, a DEAE-Dextran method (Kharbanda et. al., 1994, Cell Growth & Differentiation, 5:259-265) is used. First, harvest 2×10^7 U937 cells and wash with PBS. The U937 cells are usually grown in RPMI 1640 medium containing 10% heat-inactivated fetal bovine serum (FBS) supplemented with 100 units/ml penicillin and 100 mg/ml streptomycin.

Next, suspend the cells in 1 ml of 20 mM Tris-HCl (pH 7.4) buffer containing 0.5 mg/ml DEAE-Dextran, 8 ug GAS-SEAP2 plasmid DNA, 140 mM NaCl, 5 mM KCl, 375 uM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 1 mM MgCl_2 , and 675 uM CaCl_2 . Incubate at 37°C for 45 min.

Wash the cells with RPMI 1640 medium containing 10% FBS and then resuspend in 10 ml complete medium and incubate at 37°C for 36 hr.

The GAS-SEAP/U937 stable cells are obtained by growing the cells in 400 ug/ml G418. The G418-free medium is used for routine growth but every one to two months, the cells should be re-grown in 400 ug/ml G418 for couple of passages.

These cells are tested by harvesting 1×10^8 cells (this is enough for ten 96-well plates assay) and wash with PBS. Suspend the cells in 200 ml above described growth medium, with a final density of 5×10^5 cells/ml. Plate 200 ul cells per well in the 96-well plate (or 1×10^5 cells/well).

Add 50 ul of the supernatant prepared by the protocol described in Example 11. Incubate at 37°C for 48 to 72 hr. As a positive control, 100 Unit/ml interferon gamma can be used which is known to activate U937 cells. Over 30 fold induction is typically observed in the positive control wells. SEAP assay the supernatant according to the protocol described in Example 17.

Example 15: High-Throughput Screening Assay Identifying Neuronal Activity.

When cells undergo differentiation and proliferation, a group of genes are activated through many different signal transduction pathways. One of these genes, EGR1 (early growth response gene 1), is induced in various tissues and cell types upon

activation. The promoter of EGR1 is responsible for such induction. Using the EGR1 promoter linked to reporter molecules, activation of cells can be assessed.

Particularly, the following protocol is used to assess neuronal activity in PC12 cell lines. PC12 cells (rat pheochromocytoma cells) are known to proliferate and/or differentiate by activation with a number of mitogens, such as TPA (tetradecanoyl phorbol acetate), NGF (nerve growth factor), and EGF (epidermal growth factor). The EGR1 gene expression is activated during this treatment. Thus, by stably transfecting PC12 cells with a construct containing an EGR promoter linked to SEAP reporter, activation of PC12 cells can be assessed.

The EGR/SEAP reporter construct can be assembled by the following protocol. The EGR-1 promoter sequence (-633 to +1)(Sakamoto K et al., Oncogene 6:867-871 (1991)) can be PCR amplified from human genomic DNA using the following primers:

5' GCGCTCGAGGGATGACAGCGATAGAACCCCGG -3' (SEQ ID NO:6)

5' GCGAAGCTTCGCGACTCCCCGGATCCGCCTC-3' (SEQ ID NO:7)

Using the GAS:SEAP/Neo vector produced in Example 12, EGR1 amplified product can then be inserted into this vector. Linearize the GAS:SEAP/Neo vector using restriction enzymes XhoI/HindIII, removing the GAS/SV40 stuffer. Restrict the EGR1 amplified product with these same enzymes. Ligate the vector and the EGR1 promoter.

To prepare 96 well-plates for cell culture, two mls of a coating solution (1:30 dilution of collagen type I (Upstate Biotech Inc. Cat#08-115) in 30% ethanol (filter sterilized)) is added per one 10 cm plate or 50 ml per well of the 96-well plate, and allowed to air dry for 2 hr.

PC12 cells are routinely grown in RPMI-1640 medium (Bio Whittaker) containing 10% horse serum (JRH BIOSCIENCES, Cat. # 12449-78P), 5% heat-inactivated fetal bovine serum (FBS) supplemented with 100 units/ml penicillin and 100 ug/ml streptomycin on a precoated 10 cm tissue culture dish. One to four split is done every three to four days. Cells are removed from the plates by scraping and resuspended with pipetting up and down for more than 15 times.

Transfect the EGR/SEAP/Neo construct into PC12 using the Lipofectamine protocol described in Example 11. EGR-SEAP/PC12 stable cells are obtained by growing the cells in 300 ug/ml G418. The G418-free medium is used for routine growth but every one to two months, the cells should be re-grown in 300 ug/ml G418 for couple of passages.

To assay for neuronal activity, a 10 cm plate with cells around 70 to 80% confluent is screened by removing the old medium. Wash the cells once with PBS

(Phosphate buffered saline). Then starve the cells in low serum medium (RPMI-1640 containing 1% horse serum and 0.5% FBS with antibiotics) overnight.

The next morning, remove the medium and wash the cells with PBS. Scrape off the cells from the plate, suspend the cells well in 2 ml low serum medium. Count
5 the cell number and add more low serum medium to reach final cell density as 5×10^5 cells/ml.

Add 200 ul of the cell suspension to each well of 96-well plate (equivalent to 1×10^5 cells/well). Add 50 ul supernatant produced by Example 11, 37°C for 48 to 72 hr. As a positive control, a growth factor known to activate PC12 cells through EGR
10 can be used, such as 50 ng/ul of Neuronal Growth Factor (NGF). Over fifty-fold induction of SEAP is typically seen in the positive control wells. SEAP assay the supernatant according to Example 17.

Example 16: High-Throughput Screening Assay for T-cell Activity

15 NF- κ B (Nuclear Factor κ B) is a transcription factor activated by a wide variety of agents including the inflammatory cytokines IL-1 and TNF, CD30 and CD40, lymphotoxin-alpha and lymphotoxin-beta, by exposure to LPS or thrombin, and by expression of certain viral gene products. As a transcription factor, NF- κ B regulates the expression of genes involved in immune cell activation, control of apoptosis (NF-
20 κ B appears to shield cells from apoptosis), B and T-cell development, anti-viral and antimicrobial responses, and multiple stress responses.

In non-stimulated conditions, NF- κ B is retained in the cytoplasm with I- κ B (Inhibitor κ B). However, upon stimulation, I- κ B is phosphorylated and degraded, causing NF- κ B to shuttle to the nucleus, thereby activating transcription of target
25 genes. Target genes activated by NF- κ B include IL-2, IL-6, GM-CSF, ICAM-1 and class 1 MHC.

Due to its central role and ability to respond to a range of stimuli, reporter constructs utilizing the NF- κ B promoter element are used to screen the supernatants produced in Example 11. Activators or inhibitors of NF- κ B would be useful in treating
30 diseases. For example, inhibitors of NF- κ B could be used to treat those diseases related to the acute or chronic activation of NF- κ B, such as rheumatoid arthritis.

To construct a vector containing the NF- κ B promoter element, a PCR based strategy is employed. The upstream primer contains four tandem copies of the NF- κ B binding site (GGGGACTTTCCC) (SEQ ID NO:8), 18 bp of sequence complementary to the 5' end of the SV40 early promoter sequence, and is flanked with an XhoI site:

5 5':GCGGCCTCGAGGGGACTTTCCCGGGGACTTTCCGGGGACTTTCCGGGAC
TTCCATCCTGCCATCTCAATTAG:3' (SEQ ID NO:9)

The downstream primer is complementary to the 3' end of the SV40 promoter and is flanked with a Hind III site:

5':GCGGCAAGCTTTTTGCAAAGCCTAGGC:3' (SEQ ID NO:4)

10 PCR amplification is performed using the SV40 promoter template present in the pB-gal:promoter plasmid obtained from Clontech. The resulting PCR fragment is digested with XhoI and Hind III and subcloned into BLSK2-. (Stratagene) Sequencing with the T7 and T3 primers confirms the insert contains the following sequence:

15 5':CTCGAGGGGACTTTCCCGGGGACTTTCCGGGGACTTTCCGGGACTTTCC
ATCTGCCATCTCAATTAGTCAGCAACCATAGTCCCGCCCCTAACTCCGCCCCA
TCCCGCCCCTAACTCCGCCCAGTTCCGCCCATTCTCCGCCCCATGGCTGACT
AATTTTTTTTATTTATGCAGAGGCCGAGGCCGCCTCGGCCTCTGAGCTATTC
20 CAGAAGTAGTGAGGAGGCTTTTTTGGAGGCCTAGGCTTTTGCAAAAAGCTT:
3' (SEQ ID NO:10)

Next, replace the SV40 minimal promoter element present in the pSEAP2-promoter plasmid (Clontech) with this NF- κ B/SV40 fragment using XhoI and HindIII.

25 However, this vector does not contain a neomycin resistance gene, and therefore, is not preferred for mammalian expression systems.

In order to generate stable mammalian cell lines, the NF- κ B/SV40/SEAP cassette is removed from the above NF- κ B/SEAP vector using restriction enzymes SalI and NotI, and inserted into a vector containing neomycin resistance. Particularly, the

30 NF- κ B/SV40/SEAP cassette was inserted into pGFP-1 (Clontech), replacing the GFP gene, after restricting pGFP-1 with SalI and NotI.

Once NF- κ B/SV40/SEAP/Neo vector is created, stable Jurkat T-cells are created and maintained according to the protocol described in Example 13. Similarly, the method for assaying supernatants with these stable Jurkat T-cells is also described

in Example 13. As a positive control, exogenous TNF alpha (0.1, 1, 10 ng) is added to wells H9, H10, and H11, with a 5-10 fold activation typically observed.

Example 17: Assay for SEAP Activity

5 As a reporter molecule for the assays described in Examples 13-16, SEAP activity is assayed using the Tropix Phospho-light Kit (Cat. BP-400) according to the following general procedure. The Tropix Phospho-light Kit supplies the Dilution, Assay, and Reaction Buffers used below.

10 Prime a dispenser with the 2.5x Dilution Buffer and dispense 15 μ l of 2.5x dilution buffer into Optiplates containing 35 μ l of a supernatant. Seal the plates with a plastic sealer and incubate at 65°C for 30 min. Separate the Optiplates to avoid uneven heating.

Cool the samples to room temperature for 15 minutes. Empty the dispenser and prime with the Assay Buffer. Add 50 μ l Assay Buffer and incubate at room temperature 5 min. Empty the dispenser and prime with the Reaction Buffer (see the table below). Add 50 μ l Reaction Buffer and incubate at room temperature for 20 minutes. Since the intensity of the chemiluminescent signal is time dependent, and it takes about 10 minutes to read 5 plates on luminometer, one should treat 5 plates at each time and start the second set 10 minutes later.

20 Read the relative light unit in the luminometer. Set H12 as blank, and print the results. An increase in chemiluminescence indicates reporter activity.

Reaction Buffer Formulation:

# of plates	Rxn buffer diluent (ml)	CSPD (ml)
10	60	3
11	65	3.25
12	70	3.5
13	75	3.75
14	80	4
15	85	4.25
16	90	4.5
17	95	4.75
18	100	5
19	105	5.25
20	110	5.5
21	115	5.75
22	120	6
23	125	6.25
24	130	6.5
25	135	6.75
26	140	7
27	145	7.25

28	150	7.5
29	155	7.75
30	160	8
31	165	8.25
32	170	8.5
33	175	8.75
34	180	9
35	185	9.25
36	190	9.5
37	195	9.75
38	200	10
39	205	10.25
40	210	10.5
41	215	10.75
42	220	11
43	225	11.25
44	230	11.5
45	235	11.75
46	240	12
47	245	12.25
48	250	12.5
49	255	12.75
50	260	13

Example 18: High-Throughput Screening Assay Identifying Changes in Small Molecule Concentration and Membrane Permeability

Binding of a ligand to a receptor is known to alter intracellular levels of small molecules, such as calcium, potassium, sodium, and pH, as well as alter membrane potential. These alterations can be measured in an assay to identify supernatants which bind to receptors of a particular cell. Although the following protocol describes an assay for calcium, this protocol can easily be modified to detect changes in potassium, sodium, pH, membrane potential, or any other small molecule which is detectable by a fluorescent probe.

The following assay uses Fluorometric Imaging Plate Reader ("FLIPR") to measure changes in fluorescent molecules (Molecular Probes) that bind small molecules. Clearly, any fluorescent molecule detecting a small molecule can be used instead of the calcium fluorescent molecule, fluo-3, used here.

For adherent cells, seed the cells at 10,000 -20,000 cells/well in a Co-star black 96-well plate with clear bottom. The plate is incubated in a CO₂ incubator for 20 hours. The adherent cells are washed two times in Biotek washer with 200 ul of HBSS (Hank's Balanced Salt Solution) leaving 100 ul of buffer after the final wash.

A stock solution of 1 mg/ml fluo-3 is made in 10% pluronic acid DMSO. To load the cells with fluo-3, 50 ul of 12 ug/ml fluo-3 is added to each well. The plate is

incubated at 37°C in a CO₂ incubator for 60 min. The plate is washed four times in the Biotek washer with HBSS leaving 100 ul of buffer.

For non-adherent cells, the cells are spun down from culture media. Cells are re-suspended to 2-5x10⁶ cells/ml with HBSS in a 50-ml conical tube. 4 ul of 1 mg/ml fluo-3 solution in 10% pluronic acid DMSO is added to each ml of cell suspension. The tube is then placed in a 37°C water bath for 30-60 min. The cells are washed twice with HBSS, resuspended to 1x10⁶ cells/ml, and dispensed into a microplate, 100 ul/well. The plate is centrifuged at 1000 rpm for 5 min. The plate is then washed once in Denley CellWash with 200 ul, followed by an aspiration step to 100 ul final volume.

For a non-cell based assay, each well contains a fluorescent molecule, such as fluo-3. The supernatant is added to the well, and a change in fluorescence is detected.

To measure the fluorescence of intracellular calcium, the FLIPR is set for the following parameters: (1) System gain is 300-800 mW; (2) Exposure time is 0.4 second; (3) Camera F/stop is F/2; (4) Excitation is 488 nm; (5) Emission is 530 nm; and (6) Sample addition is 50 ul. Increased emission at 530 nm indicates an extracellular signaling event which has resulted in an increase in the intracellular Ca⁺⁺ concentration.

Example 19: High-Throughput Screening Assay Identifying Tyrosine Kinase Activity

The Protein Tyrosine Kinases (PTK) represent a diverse group of transmembrane and cytoplasmic kinases. Within the Receptor Protein Tyrosine Kinase (RPTK) group are receptors for a range of mitogenic and metabolic growth factors including the PDGF, FGF, EGF, NGF, HGF and Insulin receptor subfamilies. In addition there are a large family of RPTKs for which the corresponding ligand is unknown. Ligands for RPTKs include mainly secreted small proteins, but also membrane-bound and extracellular matrix proteins.

Activation of RPTK by ligands involves ligand-mediated receptor dimerization, resulting in transphosphorylation of the receptor subunits and activation of the cytoplasmic tyrosine kinases. The cytoplasmic tyrosine kinases include receptor associated tyrosine kinases of the src-family (e.g., src, yes, lck, lyn, fyn) and non-receptor linked and cytosolic protein tyrosine kinases, such as the Jak family, members of which mediate signal transduction triggered by the cytokine superfamily of receptors (e.g., the Interleukins, Interferons, GM-CSF, and Leptin).

Because of the wide range of known factors capable of stimulating tyrosine kinase activity, the identification of novel human secreted proteins capable of activating

tyrosine kinase signal transduction pathways are of interest. Therefore, the following protocol is designed to identify those novel human secreted proteins capable of activating the tyrosine kinase signal transduction pathways.

Seed target cells (e.g., primary keratinocytes) at a density of approximately
5 25,000 cells per well in a 96 well Loprodyne Silent Screen Plates purchased from Nalge Nunc (Naperville, IL). The plates are sterilized with two 30 minute rinses with 100% ethanol, rinsed with water and dried overnight. Some plates are coated for 2 hr with 100 ml of cell culture grade type I collagen (50 mg/ml), gelatin (2%) or polylysine (50 mg/ml), all of which can be purchased from Sigma Chemicals (St. Louis, MO) or
10 10% Matrigel purchased from Becton Dickinson (Bedford, MA), or calf serum, rinsed with PBS and stored at 4°C. Cell growth on these plates is assayed by seeding 5,000 cells/well in growth medium and indirect quantitation of cell number through use of alamarBlue as described by the manufacturer Alamar Biosciences, Inc. (Sacramento, CA) after 48 hr. Falcon plate covers #3071 from Becton Dickinson (Bedford, MA) are
15 used to cover the Loprodyne Silent Screen Plates. Falcon Microtest III cell culture plates can also be used in some proliferation experiments.

To prepare extracts, A431 cells are seeded onto the nylon membranes of Loprodyne plates (20,000/200ml/well) and cultured overnight in complete medium. Cells are quiesced by incubation in serum-free basal medium for 24 hr. After 5-20
20 minutes treatment with EGF (60ng/ml) or 50 ul of the supernatant produced in Example 11, the medium was removed and 100 ml of extraction buffer ((20 mM HEPES pH 7.5, 0.15 M NaCl, 1% Triton X-100, 0.1% SDS, 2 mM Na₃VO₄, 2 mM Na₄P₂O₇ and a cocktail of protease inhibitors (# 1836170) obtained from Boehringer Mannheim (Indianapolis, IN) is added to each well and the plate is shaken on a rotating shaker for
25 5 minutes at 4°C. The plate is then placed in a vacuum transfer manifold and the extract filtered through the 0.45 mm membrane bottoms of each well using house vacuum. Extracts are collected in a 96-well catch/assay plate in the bottom of the vacuum manifold and immediately placed on ice. To obtain extracts clarified by centrifugation, the content of each well, after detergent solubilization for 5 minutes, is removed and
30 centrifuged for 15 minutes at 4°C at 16,000 x g.

Test the filtered extracts for levels of tyrosine kinase activity. Although many methods of detecting tyrosine kinase activity are known, one method is described here.

Generally, the tyrosine kinase activity of a supernatant is evaluated by determining its ability to phosphorylate a tyrosine residue on a specific substrate (a
35 biotinylated peptide). Biotinylated peptides that can be used for this purpose include PSK1 (corresponding to amino acids 6-20 of the cell division kinase cdc2-p34) and

PSK2 (corresponding to amino acids 1-17 of gastrin). Both peptides are substrates for a range of tyrosine kinases and are available from Boehringer Mannheim.

The tyrosine kinase reaction is set up by adding the following components in order. First, add 10ul of 5uM Biotinylated Peptide, then 10ul ATP/Mg₂⁺ (5mM
5 ATP/50mM MgCl₂), then 10ul of 5x Assay Buffer (40mM imidazole hydrochloride, pH7.3, 40 mM beta-glycerophosphate, 1mM EGTA, 100mM MgCl₂, 5 mM MnCl₂, 0.5 mg/ml BSA), then 5ul of Sodium Vanadate(1mM), and then 5ul of water. Mix the components gently and preincubate the reaction mix at 30°C for 2 min. Initiate the reaction by adding 10ul of the control enzyme or the filtered supernatant.

10 The tyrosine kinase assay reaction is then terminated by adding 10 ul of 120mM EDTA and place the reactions on ice.

Tyrosine kinase activity is determined by transferring 50 ul aliquot of reaction mixture to a microtiter plate (MTP) module and incubating at 37°C for 20 min. This allows the streptavidin coated 96 well plate to associate with the biotinylated peptide.
15 Wash the MTP module with 300ul/well of PBS four times. Next add 75 ul of anti-phosphotyrosine antibody conjugated to horse radish peroxidase(anti-P-Tyr-POD(0.5u/ml)) to each well and incubate at 37°C for one hour. Wash the well as above.

Next add 100ul of peroxidase substrate solution (Boehringer Mannheim) and
20 incubate at room temperature for at least 5 mins (up to 30 min). Measure the absorbance of the sample at 405 nm by using ELISA reader. The level of bound peroxidase activity is quantitated using an ELISA reader and reflects the level of tyrosine kinase activity.

25 **Example 20: High-Throughput Screening Assay Identifying Phosphorylation Activity**

As a potential alternative and/or complement to the assay of protein tyrosine kinase activity described in Example 19, an assay which detects activation (phosphorylation) of major intracellular signal transduction intermediates can also be
30 used. For example, as described below one particular assay can detect tyrosine phosphorylation of the Erk-1 and Erk-2 kinases. However, phosphorylation of other molecules, such as Raf, JNK, p38 MAP, Map kinase kinase (MEK), MEK kinase, Src, Muscle specific kinase (MuSK), IRAK, Tec, and Janus, as well as any other phosphoserine, phosphotyrosine, or phosphothreonine molecule, can be detected by
35 substituting these molecules for Erk-1 or Erk-2 in the following assay.

Specifically, assay plates are made by coating the wells of a 96-well ELISA plate with 0.1ml of protein G (1ug/ml) for 2 hr at room temp, (RT). The plates are then rinsed with PBS and blocked with 3% BSA/PBS for 1 hr at RT. The protein G plates are then treated with 2 commercial monoclonal antibodies (100ng/well) against Erk-1 and Erk-2 (1 hr at RT) (Santa Cruz Biotechnology). (To detect other molecules, this step can easily be modified by substituting a monoclonal antibody detecting any of the above described molecules.) After 3-5 rinses with PBS, the plates are stored at 4°C until use.

A431 cells are seeded at 20,000/well in a 96-well Loprodyne filterplate and cultured overnight in growth medium. The cells are then starved for 48 hr in basal medium (DMEM) and then treated with EGF (6ng/well) or 50 ul of the supernatants obtained in Example 11 for 5-20 minutes. The cells are then solubilized and extracts filtered directly into the assay plate.

After incubation with the extract for 1 hr at RT, the wells are again rinsed. As a positive control, a commercial preparation of MAP kinase (10ng/well) is used in place of A431 extract. Plates are then treated with a commercial polyclonal (rabbit) antibody (1ug/ml) which specifically recognizes the phosphorylated epitope of the Erk-1 and Erk-2 kinases (1 hr at RT). This antibody is biotinylated by standard procedures. The bound polyclonal antibody is then quantitated by successive incubations with Europium-streptavidin and Europium fluorescence enhancing reagent in the Wallac DELFIA instrument (time-resolved fluorescence). An increased fluorescent signal over background indicates a phosphorylation.

Example 21: Method of Determining Alterations in a Gene

Corresponding to a Polynucleotide

RNA isolated from entire families or individual patients presenting with a phenotype of interest (such as a disease) is isolated. cDNA is then generated from these RNA samples using protocols known in the art. (See, Sambrook.) The cDNA is then used as a template for PCR, employing primers surrounding regions of interest in SEQ ID NO:X. Suggested PCR conditions consist of 35 cycles at 95°C for 30 seconds; 60-120 seconds at 52-58°C; and 60-120 seconds at 70°C, using buffer solutions described in Sidransky, D., et al., Science 252:706 (1991).

PCR products are then sequenced using primers labeled at their 5' end with T4 polynucleotide kinase, employing SequiTherm Polymerase. (Epicentre Technologies). The intron-exon borders of selected exons is also determined and genomic PCR

products analyzed to confirm the results. PCR products harboring suspected mutations is then cloned and sequenced to validate the results of the direct sequencing.

PCR products is cloned into T-tailed vectors as described in Holton, T.A. and Graham, M.W., Nucleic Acids Research, 19:1156 (1991) and sequenced with T7
5 polymerase (United States Biochemical). Affected individuals are identified by mutations not present in unaffected individuals.

Genomic rearrangements are also observed as a method of determining alterations in a gene corresponding to a polynucleotide. Genomic clones isolated according to Example 2 are nick-translated with digoxigenin deoxy-uridine 5'-
10 triphosphate (Boehringer Mannheim), and FISH performed as described in Johnson, Cg. et al., Methods Cell Biol. 35:73-99 (1991). Hybridization with the labeled probe is carried out using a vast excess of human cot-1 DNA for specific hybridization to the corresponding genomic locus.

Chromosomes are counterstained with 4,6-diamino-2-phenylidole and
15 propidium iodide, producing a combination of C- and R-bands. Aligned images for precise mapping are obtained using a triple-band filter set (Chroma Technology, Brattleboro, VT) in combination with a cooled charge-coupled device camera (Photometrics, Tucson, AZ) and variable excitation wavelength filters. (Johnson, Cv. et al., Genet. Anal. Tech. Appl., 8:75 (1991).) Image collection, analysis and
20 chromosomal fractional length measurements are performed using the ISee Graphical Program System. (Inovision Corporation, Durham, NC.) Chromosome alterations of the genomic region hybridized by the probe are identified as insertions, deletions, and translocations. These alterations are used as a diagnostic marker for an associated disease.

25

Example 22: Method of Detecting Abnormal Levels of a Polypeptide in a Biological Sample

A polypeptide of the present invention can be detected in a biological sample, and if an increased or decreased level of the polypeptide is detected, this polypeptide is
30 a marker for a particular phenotype. Methods of detection are numerous, and thus, it is understood that one skilled in the art can modify the following assay to fit their particular needs.

For example, antibody-sandwich ELISAs are used to detect polypeptides in a sample, preferably a biological sample. Wells of a microtiter plate are coated with
35 specific antibodies, at a final concentration of 0.2 to 10 ug/ml. The antibodies are either monoclonal or polyclonal and are produced by the method described in Example 10.

The wells are blocked so that non-specific binding of the polypeptide to the well is reduced.

The coated wells are then incubated for > 2 hours at RT with a sample containing the polypeptide. Preferably, serial dilutions of the sample should be used to validate results. The plates are then washed three times with deionized or distilled water to remove unbounded polypeptide.

Next, 50 ul of specific antibody-alkaline phosphatase conjugate, at a concentration of 25-400 ng, is added and incubated for 2 hours at room temperature. The plates are again washed three times with deionized or distilled water to remove unbounded conjugate.

Add 75 ul of 4-methylumbelliferyl phosphate (MUP) or p-nitrophenyl phosphate (NPP) substrate solution to each well and incubate 1 hour at room temperature. Measure the reaction by a microtiter plate reader. Prepare a standard curve, using serial dilutions of a control sample, and plot polypeptide concentration on the X-axis (log scale) and fluorescence or absorbance of the Y-axis (linear scale). Interpolate the concentration of the polypeptide in the sample using the standard curve.

Example 23: Formulating a Polypeptide

The secreted polypeptide composition will be formulated and dosed in a fashion consistent with good medical practice, taking into account the clinical condition of the individual patient (especially the side effects of treatment with the secreted polypeptide alone), the site of delivery, the method of administration, the scheduling of administration, and other factors known to practitioners. The "effective amount" for purposes herein is thus determined by such considerations.

As a general proposition, the total pharmaceutically effective amount of secreted polypeptide administered parenterally per dose will be in the range of about 1 µg/kg/day to 10 mg/kg/day of patient body weight, although, as noted above, this will be subject to therapeutic discretion. More preferably, this dose is at least 0.01 mg/kg/day, and most preferably for humans between about 0.01 and 1 mg/kg/day for the hormone. If given continuously, the secreted polypeptide is typically administered at a dose rate of about 1 µg/kg/hour to about 50 µg/kg/hour, either by 1-4 injections per day or by continuous subcutaneous infusions, for example, using a mini-pump. An intravenous bag solution may also be employed. The length of treatment needed to observe changes and the interval following treatment for responses to occur appears to vary depending on the desired effect.

Pharmaceutical compositions containing the secreted protein of the invention are administered orally, rectally, parenterally, intracisternally, intravaginally,

intraperitoneally, topically (as by powders, ointments, gels, drops or transdermal patch), buccally, or as an oral or nasal spray. "Pharmaceutically acceptable carrier" refers to a non-toxic solid, semisolid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type. The term "parenteral" as used herein refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion.

The secreted polypeptide is also suitably administered by sustained-release systems. Suitable examples of sustained-release compositions include semi-permeable polymer matrices in the form of shaped articles, e.g., films, or microcapsules. Sustained-release matrices include polylactides (U.S. Pat. No. 3,773,919, EP 58,481), copolymers of L-glutamic acid and gamma-ethyl-L-glutamate (Sidman, U. et al., Biopolymers 22:547-556 (1983)), poly (2- hydroxyethyl methacrylate) (R. Langer et al., J. Biomed. Mater. Res. 15:167-277 (1981), and R. Langer, Chem. Tech. 12:98-105 (1982)), ethylene vinyl acetate (R. Langer et al.) or poly-D- (-)-3-hydroxybutyric acid (EP 133,988). Sustained-release compositions also include liposomally entrapped polypeptides. Liposomes containing the secreted polypeptide are prepared by methods known per se: DE 3,218,121; Epstein et al., Proc. Natl. Acad. Sci. USA 82:3688-3692 (1985); Hwang et al., Proc. Natl. Acad. Sci. USA 77:4030-4034 (1980); EP 52,322; EP 36,676; EP 88,046; EP 143,949; EP 142,641; Japanese Pat. Appl. 83-118008; U.S. Pat. Nos. 4,485,045 and 4,544,545; and EP 102,324. Ordinarily, the liposomes are of the small (about 200-800 Angstroms) unilamellar type in which the lipid content is greater than about 30 mol. percent cholesterol, the selected proportion being adjusted for the optimal secreted polypeptide therapy.

For parenteral administration, in one embodiment, the secreted polypeptide is formulated generally by mixing it at the desired degree of purity, in a unit dosage injectable form (solution, suspension, or emulsion), with a pharmaceutically acceptable carrier, i.e., one that is non-toxic to recipients at the dosages and concentrations employed and is compatible with other ingredients of the formulation. For example, the formulation preferably does not include oxidizing agents and other compounds that are known to be deleterious to polypeptides.

Generally, the formulations are prepared by contacting the polypeptide uniformly and intimately with liquid carriers or finely divided solid carriers or both. Then, if necessary, the product is shaped into the desired formulation. Preferably the carrier is a parenteral carrier, more preferably a solution that is isotonic with the blood of the recipient. Examples of such carrier vehicles include water, saline, Ringer's solution, and dextrose solution. Non-aqueous vehicles such as fixed oils and ethyl oleate are also useful herein, as well as liposomes.

The carrier suitably contains minor amounts of additives such as substances that enhance isotonicity and chemical stability. Such materials are non-toxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, succinate, acetic acid, and other organic acids or their salts; antioxidants such as ascorbic acid; low molecular weight (less than about ten residues) polypeptides, e.g., polyarginine or tripeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids, such as glycine, glutamic acid, aspartic acid, or arginine; monosaccharides, disaccharides, and other carbohydrates including cellulose or its derivatives, glucose, manose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; counterions such as sodium; and/or nonionic surfactants such as polysorbates, poloxamers, or PEG.

The secreted polypeptide is typically formulated in such vehicles at a concentration of about 0.1 mg/ml to 100 mg/ml, preferably 1-10 mg/ml, at a pH of about 3 to 8. It will be understood that the use of certain of the foregoing excipients, carriers, or stabilizers will result in the formation of polypeptide salts.

Any polypeptide to be used for therapeutic administration can be sterile. Sterility is readily accomplished by filtration through sterile filtration membranes (e.g., 0.2 micron membranes). Therapeutic polypeptide compositions generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

Polypeptides ordinarily will be stored in unit or multi-dose containers, for example, sealed ampoules or vials, as an aqueous solution or as a lyophilized formulation for reconstitution. As an example of a lyophilized formulation, 10-ml vials are filled with 5 ml of sterile-filtered 1% (w/v) aqueous polypeptide solution, and the resulting mixture is lyophilized. The infusion solution is prepared by reconstituting the lyophilized polypeptide using bacteriostatic Water-for-Injection.

The invention also provides a pharmaceutical pack or kit comprising one or more containers filled with one or more of the ingredients of the pharmaceutical compositions of the invention. Associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration. In addition, the polypeptides of the present invention may be employed in conjunction with other therapeutic compounds.

Example 24: Method of Treating Decreased Levels of the Polypeptide

It will be appreciated that conditions caused by a decrease in the standard or normal expression level of a secreted protein in an individual can be treated by administering the polypeptide of the present invention, preferably in the secreted form. Thus, the invention also provides a method of treatment of an individual in need of an increased level of the polypeptide comprising administering to such an individual a pharmaceutical composition comprising an amount of the polypeptide to increase the activity level of the polypeptide in such an individual.

For example, a patient with decreased levels of a polypeptide receives a daily dose 0.1-100 ug/kg of the polypeptide for six consecutive days. Preferably, the polypeptide is in the secreted form. The exact details of the dosing scheme, based on administration and formulation, are provided in Example 23.

Example 25: Method of Treating Increased Levels of the Polypeptide

Antisense technology is used to inhibit production of a polypeptide of the present invention. This technology is one example of a method of decreasing levels of a polypeptide, preferably a secreted form, due to a variety of etiologies, such as cancer.

For example, a patient diagnosed with abnormally increased levels of a polypeptide is administered intravenously antisense polynucleotides at 0.5, 1.0, 1.5, 2.0 and 3.0 mg/kg day for 21 days. This treatment is repeated after a 7-day rest period if the treatment was well tolerated. The formulation of the antisense polynucleotide is provided in Example 23.

Example 26: Method of Treatment Using Gene Therapy

One method of gene therapy transplants fibroblasts, which are capable of expressing a polypeptide, onto a patient. Generally, fibroblasts are obtained from a subject by skin biopsy. The resulting tissue is placed in tissue-culture medium and separated into small pieces. Small chunks of the tissue are placed on a wet surface of a tissue culture flask, approximately ten pieces are placed in each flask. The flask is turned upside down, closed tight and left at room temperature over night. After 24 hours at room temperature, the flask is inverted and the chunks of tissue remain fixed to the bottom of the flask and fresh media (e.g., Ham's F12 media, with 10% FBS, penicillin and streptomycin) is added. The flasks are then incubated at 37°C for approximately one week.

At this time, fresh media is added and subsequently changed every several days. After an additional two weeks in culture, a monolayer of fibroblasts emerge. The monolayer is trypsinized and scaled into larger flasks.

pMV-7 (Kirschmeier, P.T. et al., DNA, 7:219-25 (1988)), flanked by the long terminal repeats of the Moloney murine sarcoma virus, is digested with EcoRI and HindIII and subsequently treated with calf intestinal phosphatase. The linear vector is fractionated on agarose gel and purified, using glass beads.

5 The cDNA encoding a polypeptide of the present invention can be amplified using PCR primers which correspond to the 5' and 3' end sequences respectively as set forth in Example 1. Preferably, the 5' primer contains an EcoRI site and the 3' primer includes a HindIII site. Equal quantities of the Moloney murine sarcoma virus linear backbone and the amplified EcoRI and HindIII fragment are added together, in the
10 presence of T4 DNA ligase. The resulting mixture is maintained under conditions appropriate for ligation of the two fragments. The ligation mixture is then used to transform bacteria HB101, which are then plated onto agar containing kanamycin for the purpose of confirming that the vector has the gene of interest properly inserted.

15 The amphotropic pA317 or GP+am12 packaging cells are grown in tissue culture to confluent density in Dulbecco's Modified Eagles Medium (DMEM) with 10% calf serum (CS), penicillin and streptomycin. The MSV vector containing the gene is then added to the media and the packaging cells transduced with the vector. The packaging cells now produce infectious viral particles containing the gene (the packaging cells are now referred to as producer cells).

20 Fresh media is added to the transduced producer cells, and subsequently, the media is harvested from a 10 cm plate of confluent producer cells. The spent media, containing the infectious viral particles, is filtered through a millipore filter to remove detached producer cells and this media is then used to infect fibroblast cells. Media is removed from a sub-confluent plate of fibroblasts and quickly replaced with the media
25 from the producer cells. This media is removed and replaced with fresh media. If the titer of virus is high, then virtually all fibroblasts will be infected and no selection is required. If the titer is very low, then it is necessary to use a retroviral vector that has a selectable marker, such as neo or his. Once the fibroblasts have been efficiently infected, the fibroblasts are analyzed to determine whether protein is produced.

30 The engineered fibroblasts are then transplanted onto the host, either alone or after having been grown to confluence on cytodex 3 microcarrier beads.

It will be clear that the invention may be practiced otherwise than as particularly described in the foregoing description and examples. Numerous modifications and variations of the present invention are possible in light of the above teachings and,
35 therefore, are within the scope of the appended claims.

The entire disclosure of each document cited (including patents, patent applications, journal articles, abstracts, laboratory manuals, books, or other

disclosures) in the Background of the Invention, Detailed Description, and Examples is hereby incorporated herein by reference. Further, the hard copy of the sequence listing submitted herewith and the corresponding computer readable form are both incorporated herein by reference in their entireties.

What Is Claimed Is:

1. An isolated nucleic acid molecule comprising a polynucleotide having a nucleotide sequence at least 95% identical to a sequence selected from the group consisting of:

(a) a polynucleotide fragment of SEQ ID NO:X or a polynucleotide fragment of the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X;

(b) a polynucleotide encoding a polypeptide fragment of SEQ ID NO:Y or a polypeptide fragment encoded by the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X;

(c) a polynucleotide encoding a polypeptide domain of SEQ ID NO:Y or a polypeptide domain encoded by the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X;

(d) a polynucleotide encoding a polypeptide epitope of SEQ ID NO:Y or a polypeptide epitope encoded by the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X;

(e) a polynucleotide encoding a polypeptide of SEQ ID NO:Y or the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X, having biological activity;

(f) a polynucleotide which is a variant of SEQ ID NO:X;

(g) a polynucleotide which is an allelic variant of SEQ ID NO:X;

(h) a polynucleotide which encodes a species homologue of the SEQ ID NO:Y;

(i) a polynucleotide capable of hybridizing under stringent conditions to any one of the polynucleotides specified in (a)-(h), wherein said polynucleotide does not hybridize under stringent conditions to a nucleic acid molecule having a nucleotide sequence of only A residues or of only T residues.

2. The isolated nucleic acid molecule of claim 1, wherein the polynucleotide fragment comprises a nucleotide sequence encoding a secreted protein.

3. The isolated nucleic acid molecule of claim 1, wherein the polynucleotide fragment comprises a nucleotide sequence encoding the sequence identified as SEQ ID NO:Y or the polypeptide encoded by the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X.

4. The isolated nucleic acid molecule of claim 1, wherein the polynucleotide fragment comprises the entire nucleotide sequence of SEQ ID NO:X or the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X.
5. The isolated nucleic acid molecule of claim 2, wherein the nucleotide sequence comprises sequential nucleotide deletions from either the C-terminus or the N-terminus.
6. The isolated nucleic acid molecule of claim 3, wherein the nucleotide sequence comprises sequential nucleotide deletions from either the C-terminus or the N-terminus.
7. A recombinant vector comprising the isolated nucleic acid molecule of claim 1.
8. A method of making a recombinant host cell comprising the isolated nucleic acid molecule of claim 1.
9. A recombinant host cell produced by the method of claim 8.
10. The recombinant host cell of claim 9 comprising vector sequences.
11. An isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence selected from the group consisting of:
 - (a) a polypeptide fragment of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z;
 - (b) a polypeptide fragment of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z, having biological activity;
 - (c) a polypeptide domain of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z;
 - (d) a polypeptide epitope of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z;
 - (e) a secreted form of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z;
 - (f) a full length protein of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z;

- (g) a variant of SEQ ID NO:Y;
- (h) an allelic variant of SEQ ID NO:Y; or
- (i) a species homologue of the SEQ ID NO:Y.

12. The isolated polypeptide of claim 11, wherein the secreted form or the full length protein comprises sequential amino acid deletions from either the C-terminus or the N-terminus.

13. An isolated antibody that binds specifically to the isolated polypeptide of claim 11.

14. A recombinant host cell that expresses the isolated polypeptide of claim 11.

15. A method of making an isolated polypeptide comprising:
(a) culturing the recombinant host cell of claim 14 under conditions such that said polypeptide is expressed; and
(b) recovering said polypeptide.

16. The polypeptide produced by claim 15.

17. A method for preventing, treating, or ameliorating a medical condition, comprising administering to a mammalian subject a therapeutically effective amount of the polypeptide of claim 11 or the polynucleotide of claim 1.

18. A method of diagnosing a pathological condition or a susceptibility to a pathological condition in a subject comprising:

- (a) determining the presence or absence of a mutation in the polynucleotide of claim 1; and
- (b) diagnosing a pathological condition or a susceptibility to a pathological condition based on the presence or absence of said mutation.

19. A method of diagnosing a pathological condition or a susceptibility to a pathological condition in a subject comprising:

- (a) determining the presence or amount of expression of the polypeptide of claim 11 in a biological sample; and
- (b) diagnosing a pathological condition or a susceptibility to a pathological condition based on the presence or amount of expression of the polypeptide.

20. A method for identifying a binding partner to the polypeptide of claim 11 comprising:

- (a) contacting the polypeptide of claim 11 with a binding partner; and
- (b) determining whether the binding partner effects an activity of the polypeptide.

21. The gene corresponding to the cDNA sequence of SEQ ID NO:Y..

22. A method of identifying an activity in a biological assay, wherein the method comprises:

- (a) expressing SEQ ID NO:X in a cell;
- (b) isolating the supernatant;
- (c) detecting an activity in a biological assay; and
- (d) identifying the protein in the supernatant having the activity.

23. The product produced by the method of claim 20.

<110> Rosen et al.
Human Genome Sciences, Inc.

<120> 125 Human Secreted Proteins

<130> P2020.PCT

<140> Unassigned

<141> 1998-11-04

<150> 60/064,911

<151> 1997-11-07

<150> 60/064,912

<151> 1997-11-07

<150> 60/064,983

<151> 1997-11-07

<150> 60/064,900

<151> 1997-11-07

<150> 60/064,988

<151> 1997-11-07

<150> 60/064,987

<151> 1997-11-07

<150> 60/064,908

<151> 1997-11-07

<150> 60/064,984

<151> 1997-11-07

<150> 60/064,985

<151> 1997-11-07

<150> 60/066,094

<151> 1997-11-17

<150> 60/066,100

<151> 1997-11-17

<150> 60/066,089

<151> 1997-11-17

<150> 60/066,095

<151> 1997-11-17

<150> 60/066,090

<151> 1997-11-17

<160> 502

<170> PatentIn Ver. 2.0

<210> 1
 <211> 733
 <212> DNA
 <213> Homo sapiens

<400> 1
 gggatccgga gcccaaactct tctgacaaaa ctcacacatg cccaccgtgc ccagcacctg 60
 aattcgaggg tgcaccgtca gtcttcctct tcccccaaa acccaaggac accctcatga 120
 tctcccgga tcttgagggtc acatgcgtgg tgggtggacgt aagccacgaa gaccctgagg 180
 tcaagttcaa ctggtacgtg gacggcgtgg aggtgcataa tgccaagaca aagccgcggg 240
 aggagcagta caacagcacg taccgtgtgg tcagcgtcct caccgtcctg caccaggact 300
 ggctgaatgg caaggagtac aagtgcgaagg tctccaacaa agccctccca acccccatcg 360
 agaaaacat ctccaaagcc aaagggcagc cccgagaacc acaggtgtac accctgcccc 420
 catcccgga tgagctgacc aagaaccagg tcagcctgac ctgcctgggtc aaaggcttct 480
 atccaagcga catcgccgtg gagtgggaga gcaatgggca gccggagAAC aactacaaga 540
 ccacgcctcc cgtgctggac tccgacggct ccttcttcct ctacagcaag ctcaccgtgg 600
 acaagagcag gtggcagcag gggaacgtct tctcatgctc cgtgatgcat gaggtctctg 660
 acaaccacta cagcagaag agcctctccc tgtctccggg taaatgagtg cgacggccgc 720
 gactctagag gat 733

<210> 2
 <211> 5
 <212> PRT
 <213> Homo sapiens

<220>
 <221> Site
 <222> (3)
 <223> Xaa equals any of the twenty naturally occurring L-amino acids

<400> 2
 Trp Ser Xaa Trp Ser
 1 5

<210> 3
 <211> 86
 <212> DNA
 <213> Homo sapiens

<400> 3
 gcgcctcgag atttccccga aatctagatt tccccgaaat gatttccccg aaatgatttc 60
 cccgaaatat ctgccatctc aattag 86

<210> 4
 <211> 27
 <212> DNA
 <213> Homo sapiens

<400> 4
 gcggcaagct ttttgcaaag cctaggc 27

<210> 5
 <211> 271
 <212> DNA
 <213> Homo sapiens

3

<400> 5
ctcgagattt ccccgaaatc tagatttccc cgaaatgatt tccccgaaat gatttccccg 60
aaatatctgc catctcaatt agtcagcaac catagtcccc cccctaactc cgcccatccc 120
gcccctaact cgcgccagtt ccgcccattc tccgccccat ggctgactaa tttttttat 180
ttatgcagag gccgaggccg cctcggcctc tgagctattc cagaagtagt gaggaggctt 240
ttttggaggc ctaggctttt gcaaaaagct t 271

<210> 6
<211> 32
<212> DNA
<213> Homo sapiens

<400> 6
gcgctcgagg gatgacagcg atagaacccc gg 32

<210> 7
<211> 31
<212> DNA
<213> Homo sapiens

<400> 7
gcgaagcttc gcgactcccc ggatccgcct c 31

<210> 8
<211> 12
<212> DNA
<213> Homo sapiens

<400> 8
ggggactttc cc 12

<210> 9
<211> 73
<212> DNA
<213> Homo sapiens

<400> 9
gcggcctcga ggggactttc ccggggactt tccggggact ttccgggact ttccatcctg 60
ccatctcaat tag 73

<210> 10
<211> 256
<212> DNA
<213> Homo sapiens

<400> 10
ctcgagggga ctttcccggg gactttccgg ggactttccg ggactttcca tctgccatct 60
caattagtca gcaaccatag tcccgcacct aactccgccc atcccgcccc taactccgcc 120
cagttccgcc cattctccgc cccatggctg actaattttt tttatttatg cagaggccga 180
ggccgcctcg gcctctgagc tattccagaa gtagtgagga ggcttttttg gaggcctagg 240
cttttgcaaa aagctt 256

4

<210> 11
 <211> 1010
 <212> DNA
 <213> Homo sapiens

<400> 11
 gcgtccgtat gttccagtgt gggttattgc agcagctttg tactatccta atggctactg 60
 ggggttcctgc tgatatcctg actgagacca taaatactgt atcagaagtt attcgaggct 120
 gccaaagtaaa ccaagactac tttgcatctg taaatgcacc ttcaaaccga ccaagaccgg 180
 caattgtagt acttctcatg tccatgggta atgaaaggca gccatttggt ttgcgctgtg 240
 ctgttctcta ttgtttccag tggttcttgt ataaaaacca aaaaggacaa ggagaaatcg 300
 tgtcaacact tttaccttct accattgatg caacaggtaa ttcagtttca gctggccagt 360
 tattatgtgg aggtttgttt tctactgatt cactttcaaa ctggtgtgct gctgtggccc 420
 ttgcccattgc gttgcaagaa aatgccaccc agaaagaaca gttgctcagg gttcaacttg 480
 ctacaagtat tggcaaccct ccagtttctt tacttcaaca gtgcaccaat attctttcac 540
 agggaaagcaa aatacaaaaca agagttggat tattaatgtt gctttgtacc tggctaagca 600
 attgtcccat tgcagtaacg cattttcttc acaattcagc caatgttcca ttccttacag 660
 gacaaattgc agaaaatctt ggagaagaag agcagtttgt ccaaggctta tgtgcccttt 720
 tggtgggcat ttcgatttat ttcaatgata actcacttga gagctacatg aaagagaagc 780
 taaaacaact gattgagaag aggattggca aagagaattt catagagaaa ctaggattta 840
 ttagcaaaaca tgagttgtat tccagagcat ctcagaaacc ccagccaaac tttcccagtc 900
 cagaatacat gatatttgat catgagttta cgaagctggt aaaagaactt gaagggtgta 960
 taactaaggc tatttataag tccagtgaag aagataaaaa aaaaaaaaaa 1010

<210> 12
 <211> 1559
 <212> DNA
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (1547)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (1549)
 <223> n equals a,t,g, or c

<400> 12
 ccggctacta ggaactagtg gatcccccg gctgccaggt ttttattctt tatatataaa 60
 atgtctccca gatacaaaa gatcaaataa aaatcccacc cctcctactg cgttcctagg 120
 ttggctgtat acttatttga ggaatgtttc acagcccttg tttcagtgca gtattttctg 180
 gtttatgggg ctggaatcaa agggaccgct attccctctg cccaacaca cagacttacg 240
 aatcagtgtc ccctatagca ccatactgca ggattgggtg gcatgactgt ggcattccatc 300
 agacataattc tggtagagat ctgggtaccg atagccttag ccatgggcac ccgaggcctc 360
 actcaaatgt tagctgtcat tcaatccaga agccaatggg cactctcttg aagtctgctt 420
 ggctcttctg ttctcttttt tagtaggtct gatttattaa tatgatcaat acagggttaac 480
 ttgtgaagtc ctatatacaa cactcttctt cctctagaca actttgctat tgctacttag 540
 aaatgatgta aatcattgga ggtgtgataa atttacaat atttagttt taaccatag 600
 caagggagat attgggctag agtgtaaaac acagagcatt taatatacat aggaaattta 660
 ggagtaaaat tttgatgcta attttccaaa atcatttcta ctatttctcg tcattataag 720
 attgctcaca tattttgaat ggcattccac agtagtcaca ggcaaattag aataacatat 780
 gctttacaaa acaaaaaggc atatcagaaa atgagtcaaa gtaaaaaaaa tggcaggtag 840
 catccattct ccagtaata atgaatatgg agtgatttta attacaaatt tccaaatata 900
 tattcgcata ataaagacaa ggaaaatgaa acaaaatcaa atgtgttttg ttgaaataag 960

5

cgcatgagct	tgaaggatcm	agtactgatg	gctccagggtg	ttttccatgt	tagattccct	1020
ttgaaagtac	tgatgccatc	agcacttgtc	agctgggtacc	caatgatgta	aattttgagc	1080
ttgaatactt	ttcatrggar	rgcatcacat	tcctaaagga	gacaaatgac	cactaaaaac	1140
taaccattaa	tatagattac	ttatggaggg	ttttatcttg	actactaatt	aagggagtaa	1200
ttgtatctct	aaactttata	tcagattgcc	agtggatttg	ccagattatt	aatgcttttt	1260
tagaatacat	gggggggtggg	gttataatca	cataacacag	tttttcatat	tttttaaaag	1320
gggttgacc	aagacataag	tttcattgac	aatcttaatg	ttttcactgt	agacaatatg	1380
ttagaataac	tgcagtgagt	cacacagcat	ataacttctt	aatggtaa	tcagaaccag	1440
tcttccagat	aagcaagtat	aacatttttag	ttaaaaaaaa	aaaaaaaaaa	aaaaaaaaaca	1500
aaaaaaaaaa	aaaaaactcg	aggggggggtc	cgtacccaa	tctcctnana	tgcacgtga	1559

<210> 13

<211> 1589

<212> DNA

<213> Homo sapiens

<400> 13

ggcagagctt	ttgctctgag	aacaatgcta	cggtagttt	ccagtgtgtt	tgctttgcct	60
ttttatttga	attttagaat	ttattat	ttt	cttattttaa	tggtattcat	120
ttttcttcta	ctaatttga	gtatcacagc	ttcgtactcc	ttgacttgca	ttctctcagg	180
agctggggag	ccaagctcgg	tctccgcttc	ggtaggttca	ggccccgggt	tctgtctggc	240
ggctctgctt	ctaagcgga	ctggcggtt	tgcagcaacg	cttttgccag	tagcgccac	300
tgagcggtt	ttcagttgct	gcaccgttct	tagcgcccaa	cggaaacgtt	cccgtacg	360
gagtcataa	gttgctctg	tcgtcacttt	ctcggttctg	gcacgctag	tatctcctag	420
gaacttcta	aamccgtt	acgggtgcaga	acggtaattg	caccgagcct	ctgtgccaac	480
tagtacttcc	gtaataccgt	tagccggctc	ttttcctgtc	acaacacatt	aaaacgtgtt	540
aaaacacatt	aaacggatta	aacacattar	acggatta	acacattaaa	acgtcacac	600
acatttaaac	agagtcggtc	cattcytcca	gacagtatgt	ttcgggatgg	aagcaccct	660
cctaagaatc	cgcagtttct	ttstaggcagc	accgttatgg	cgcyttaa	gatatttg	720
tggcaatacc	agcgctatcc	gctaggtgcc	ggcgcttgct	gagtcctcagc	gccgccagat	780
tcttgctagc	gaggtccaga	gcagtgcaga	gacctagcgg	acagttttcc	ggtggcagca	840
atgctcattt	cccggacaca	gatgggtttt	ggscctgaa	gaactcttta	atccctgcgt	900
tttgtagcgc	tcgttctctg	taagggttgc	ttartatccg	tttcatttcg	acttcttttc	960
tcgctgtttt	tctgtcggaa	ttacgatttg	ttttgtttct	atgtgtyctc	tagaatttta	1020
ccggttttca	tttgctact	aattttcgtg	cattcggttac	tattgagttt	cataatatct	1080
gggtgccmtc	tgccackag	ctccggagag	cataaatacc	caggctgatg	gtagtgttaa	1140
aggttctgct	tcattctcgtc	aatcagcgca	aacgcttatg	tcctcctgct	ctgaggctta	1200
attaagcgat	ggagcaacat	tttcttggtc	atgtgaagcc	tgcttgagat	tccgcagagg	1260
tggtgcctga	gctccagcct	cggcctat	tgaattcaga	agactgtgat	attactgtgt	1320
gagccaccac	tttgggaggc	cagcatgggc	agatcacttg	aggtcaggag	ttggagacca	1380
gcctggccca	agtggtgaaa	ccccatctct	actaaaaata	caaaaattag	ccggtgatga	1440
tgcagtcctg	caatcccagc	tactcgggag	gctgaggcag	gagaatctct	tgagctcgga	1500
ggcagaggtt	gcagtgagct	cagatcgtgc	caccgtctc	cagcctgggc	agactgagac	1560
tccatcaaaa	aaaaaaaaaa	aaaactcga				1589

<210> 14

<211> 1255

<212> DNA

<213> Homo sapiens

<400> 14

gctgaaatgt	ctcatttagc	taaatggaaa	cattcaccag	acacttctag	ctagctctcc	60
cacatcactt	tgagccacac	aggcacttcc	atgtttggga	ctgaattgcc	tgggaaatccg	120
tatacccaaa	atctgagccc	tcattcttac	ccttgggtga	tgctctcca	tagccctcag	180
caatgctctc	cactcgtctg	atggggctac	atctcgtgca	gattttgtgg	cgctgttggga	240
ccagttcggc	aaccattaca	tccaggaagc	tatctacggc	tttgaggagt	cctgttctat	300

ctggtaccca aacaagcagg tccagcggcg actctggctg ⁶gagtatgaag acatcagtaa 360
 aggcaactcc ccatcagatg agtctgagga gcgggaaaga gacccaagt gctgacattc 420
 ccagaataca tcaccagctt gtcagactcc ggcaccaagc gcatggcggc tggagtcgac 480
 atggagtgcc agagcaaggg acgatgcccc tcgtcctgcc ccctgtgtca tgtgacatcc 540
 agccctgaca cccctgctga gccggttctg ctggaggtga ccaaagcagc ccccatctat 600
 gaactagtga ccaacaacca gaccagagg ctcttgagg aggctaccat gagctctctc 660
 tgggtgctcag ggactggaga tgtcatcgag gactgggtgc gatgtgactc cactgctttt 720
 ggagctgatg gactccccac ctgtgcgcct cccccacagc ctgtgtatgg ttctctttct 780
 ctctttcagc attactctgg aaacagataa tgagcgtcta ctatttgcaa agaaataggc 840
 tgggtactgg gaaacaaaga aaaatatgac acaatacctg cctttagaga gctttcaatc 900
 taccaagggg ggcagcacag atgggttttag tagagaaagg gtgtgacaag tgtaacaag 960
 aggcatttaa agaaagtgtg aagagggaat ggtcaagtct gaactgacgc catggtgacg 1020
 gctctgtgga ggaggtggcg cttgaacttc tccttgaagg tggagaaagt aggacagcat 1080
 aactagaaga caatggggag aaaagtgcc ttgtaatgga agaagttaga gatctacact 1140
 gagagcttaa ctagaccaa aacataacca ctcatgtaac acttactggg tgtccaggca 1200
 ttgtatggag cagttagata ctgctcattt aatagacaaa aaaaaaaaaa aaaaa 1255

<210> 15

<211> 1191

<212> DNA

<213> Homo sapiens

<400> 15

gaattcggca cgagcggagg ccgaagagtt gagtcggtg ctaagcaacg aacttcacag 60
 acagcgatcc ccaggtgttt catttggttt atcagtggtt aatttgatga atgccatcat 120
 gggaagtggc atccttggtt tagcttatgt tatggctaata accggtgtct ttggatttag 180
 cttcttgctg ctgacagttg ctctcctggc ttcttactca gtccatcttc tgccttagtat 240
 gtgtattcag acagctgtaa catcttatga agatcttggc ctctttgcat ttggattacc 300
 tggaaagtgt gtggtggcag gcaccataat aattcagaat attggagcta tgtcatctta 360
 tcttttaatt attaaaacag agcttctctg tgctattgca gaatttttga ctggagacta 420
 tagtagatat tggatcttg atggacaaac actactaata atcatatgtg ttggcattgt 480
 gtccctctt gcacttcttc ccaaaatagg ctttcttggc tacacaagta gtttatcatt 540
 ttyctttatg atgttcttg ctcttggtgt aataattaaa aaatgggtcca tcccttgctc 600
 tctgacatta aattatgtag agaaaggctt ccagatttca aaygttacag atgattgtaa 660
 gccaaagctc tttcatttct ccaaagagag tgcttatgcc ttaccaacca tggctttttc 720
 atttctctgc catacctcaa tattgcccat atactgtgaa cttcaaagtc cttcaaagaa 780
 aagaatgcag aatgttacca atacagcaat tgctttaagt tttctcattt attttatatc 840
 tgcactcttt gggtagctca ctttttatgg gtctcattct gtcgcacagg ttggcgtgca 900
 gtggtgtgat ctgagctcat tgcaacctct gcctcccgga ctcaagcaat cctcccacct 960
 cagcctccag agtagctgag actacagaaa aggaagagat accatggaga tgtgcacca 1020
 gaggaaggc cagcaagga cacagcaaga aggcaactgt ttacaagcca agggaagagg 1080
 cctcaggaga accaaacgtg tccacacctt gatcttgac ttcccaacct ccagaactgt 1140
 gagcaaataa atgatgttgt ttaatcaaaa aaaaaaaaaa caagggggcc g 1191

<210> 16

<211> 1186

<212> DNA

<213> Homo sapiens

<400> 16

gaattcggca cgagattgaa tgttccagat aatccctttc ccagtcctgc ctgacatctg 60
 ggtagggggg ttgtccctgg aattctggga cactggctgg gggttgagga gagaagccag 120
 tacctacctg gctgcaggat gaagctggcc agtggtctt tggttttgtg gctcagcctt 180
 ggggggtggc tggctcagag cgacacgagc cctgacacgg aggagtcta ttcagactgg 240
 ggctctcgcc acctccgggg aagctttgaa tccgtcaata gctacttcga ttcttttctg 300
 gagctgctgg gaggaagaa tggagtctgt cagtacaggt gccgatatgg aaaggcacca 360

atgcccagac	ctggetacaa	gccccaaagag	cccaatggct	gcggtccta	tttcctgggt	420
ctcaaggtag	cagaaagtat	ggacttgggc	attccagcaa	tgacaaagt	ctgcaaccag	480
ctggatgtct	gttatgacac	ttgcgggtgcc	aacaaatate	gctgtgatgc	aaaattccga	540
tgggtgtctcc	amtcgatctg	ctctgacctt	aagcggagtc	tgggctttgt	ctccaaagt	600
gaagcctgtg	attccctggg	tgacactgtg	ttcaacaccg	tgtggacctt	gggctgccgc	660
cccttttatga	atagtcagcg	ggcagcttgc	atctgtgcag	aggaggagaa	ggaagagtta	720
tgaggaagaa	gtgattcctt	cctgggtttg	agtgcacca	cagctgtcag	ccttcaagat	780
gtcaagtctt	cgartcagcg	tgactcattc	gttcttccaa	cagtttgagc	accacaaagc	840
aggagaaagg	gaacatTTTT	ctacagctgg	aaagtgtgac	ctatcctttg	aggaaatttg	900
aaaaaagaca	tggagtgggt	tgaaagctac	tcttcattta	agactgctct	ccccaaccaa	960
gacacatttg	cctggaaatt	cagttcttag	cttaaagact	aaaatgcaag	caaaccctgc	1020
aattcctgga	cctgatagtt	atattcatga	gtgaaattgt	ggggagtcca	gccatttggg	1080
aggcaatgac	tttctgctgg	cccatgtttc	agttgccagt	aagcttctca	catttaataa	1140
agtgtacttt	ttagaacatt	tggaaaaaaa	aaaaaaaaaa	actcga		1186

<210> 17

<211> 1182

<212> DNA

<213> Homo sapiens

<400> 17

cccacgcgtc	cggtaaaata	taaagaaact	gaaccagtgt	gtctttttcac	catagatata	60
agagtccgga	ccgccagca	cacaaggcca	gcagctgtgt	cctctgtcac	gctctcgcta	120
tagctgttgt	ccagatcggt	atcttctcag	aaagctgggc	atttgccaag	aacatcaact	180
tctataatgt	gaggcctcct	ctcgacctta	caccatttcc	aaatagcttc	aagtgttcta	240
cttgtgaaaa	cgcaggggat	aattataact	gcaatcgatg	ggcagaagac	aaatgggtgc	300
cacaaaaatac	acagtactgt	ttgacagttc	atcacttcac	cagccacgga	agaagcacat	360
ccatcaccaa	aaagtgtgcc	tccagaagtg	aatgtcattt	tgtcgggttg	caccacagcc	420
gagattctga	acatacggag	tgtaggtctt	gctgtgaagg	aatgatctgc	aatgtagaat	480
taccaccaa	tcacactaat	gcagtgtttg	ccgtaatgca	cgctcagaga	acatctggca	540
gcagtgcctc	cacactctac	cttaccagt	cttgccctgg	tctttgtgct	tccattgctg	600
tgatgccacc	attcctagga	gaggcagaga	ccagcctcta	aagcacaagc	caaaaactgt	660
gtgaacgggtg	aactttggag	tgaagatcaa	tcttgcaatt	ggtgaagagt	gcacattgga	720
cctcaaggcg	aaagccagtg	gtttgcttgg	ataaaatgtt	cccgcagtag	gccacaggac	780
tgaggatggg	aatttggcag	ggcctgagaa	gatggtctga	cttccaggct	tcctgggtcaa	840
agagagctac	gtttgggcag	ttctgcagag	aggatcctgg	caactagtcc	cacctgacta	900
ggccttttagc	tgaaggatt	tcttgacctc	cttgactgcc	tcagaggctg	ccagggtcaa	960
ccctcttgtt	tatgtgatta	gctcagagca	tctctatgaa	atctaaccct	tcccctcatg	1020
agaaagcagt	tttccccacc	aacagcatag	tcaatgagaa	aggcaactgt	acgaagaaaa	1080
cttcagtggt	aactaatatg	aaatctattt	gcaatttatg	gggggaaata	aagcttttaa	1140
attatacaat	gtaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aa		1182

<210> 18

<211> 1171

<212> DNA

<213> Homo sapiens

<400> 18

gcttcagacc	tttgtgatac	accatgctgc	gtgggacgat	gacggcgtgg	agaggaatga	60
ggcctgaggt	cacactggct	tgctcctcc	tagccacagc	aggctgcttt	gctgacttga	120
acgaggtecc	tcaggtcacc	gtccagcctg	cgccaccgt	ccagaagccc	ggaggcactg	180
tgatcttggg	ctgcgtgggt	gaacctccaa	ggatgaatgt	aacctggcgc	ctgaatggaa	240
aggagctgaa	tggctcggat	gatgctctgg	gtgtcctcat	caccacacggg	accctcgtca	300
tactgcctt	taacaaccac	actgtgggac	ggtaccagt	tgtggcccg	atgcctgcgg	360
gggctgtggc	cacgtgccag	ccactgtgac	actagccagt	gagtctgctc	ctttgcctcc	420
ctgccatggg	gcggctccctc	ctcatctctc	ccacctgaa	gccccacca	ttcatgctgc	480

8

ctcttgttac	tcttagcata	aaatgggcct	taactgcaga	aatgtcaa	cagaacagta	540
gctgcctagt	aatgcccagt	gatgggggac	ccttgtgccc	ttggaaaacc	tcactccaag	600
tagaggctgt	atctggagtg	agtgtctaca	gagaggggaa	ttggtcagtg	catggcagaa	660
cttgacatgg	cagaactgtt	ccgtgggccc	cagagcaggg	cctgccgtgc	cttcctactt	720
ggtcatectg	tggctgtagg	tgcactgtcc	caactgctca	cacactttcc	agtcccctcc	780
ctccattctc	accacagtta	cagtcggagc	aggagagagc	aggacaccaa	gagacatggg	840
cagggccacg	tgtagtcttg	acagagccca	cccagcccag	acttggcctc	ctgtctctct	900
gtgaggggca	agcatagtcc	gaaggcctgg	attataaata	ttagaacata	atgaaaagga	960
aactgtgtgg	caaagggata	aagggatata	gagaaaagaa	acaaggaaga	gatgaggtga	1020
cagtttggac	caaggaaaat	gccaagagaa	gacttcacac	tacctttttt	tttttctggt	1080
tttgccattc	ttttattcat	gttgttacac	tcagtatagg	atagttttat	aaaatcatta	1140
tgtctgtaga	aaaaaaaaaa	aaaaaaaaaa	a			1171

<210> 19

<211> 1337

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (22)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (1318)

<223> n equals a,t,g, or c

<400> 19

cggggcttcg	gcgccaggcc	angcgctagt	cggtctggta	aggattttaca	aaaggtgcag	60
gtatgagcag	gtctgaagac	taacattttg	tgaagttgta	aaacagaaaa	cctgttagaa	120
atgtgggtgt	ttcagcaagg	cctcagtttc	cttccttcag	cccttgtaat	ttggacatct	180
gctgctttca	tattttcata	cattactgca	gtaacactcc	accatataga	ccgggcttta	240
ccttatatca	gtgacactgg	tacagtagct	ccagaaaaat	gcttattttg	ggcaatgcta	300
aatattgcgg	cagttttatg	cattgctacc	atttatgttc	gttataagca	agttcatgct	360
ctgagtcctg	aagagaacgt	tatcatcaaa	ttaaacaagg	ctggccttgt	acttgggaata	420
ctgagttggt	taggactttc	tattgtggca	aacttccaga	aaacaaccct	ttttgctgca	480
catgtaagtg	gagctgtgct	tacctttggt	atgggctcat	tatatatggt	tgttcagacc	540
atcctttcct	accaaagtca	gcccaaaatc	catggcaaac	aagtcttctg	gatcagactg	600
ttgttggtta	tctggtgtgg	agtaagtgca	cttagcatgc	tgacttgctc	atcagttttg	660
cacagtggca	attttgggac	tgatttagaa	cagaaactcc	attggaaccc	cgaggacaaa	720
ggttatgtgc	ttcacatgat	cactactgca	gcagaatggt	ctatgtcatt	ttccttcttt	780
ggttttttcc	tgacttacat	tcgtgatttt	cagaaaattt	ctttacgggt	ggaagccaat	840
ttacatggat	taaccctcta	tgacactgca	ccttgcccta	ttaacaatga	acgaacacgg	900
ctactttcca	gagatatattg	atgaaaggat	aaaatatattc	tgtaatgatt	atgattctca	960
gggattgggg	aaaggttcac	agaagttgct	tattcttctc	tgaaattttc	aaccacttaa	1020
tcaaggctga	cagtaacact	gatgaatgct	gataatcagg	aaacatgaaa	gaagccattt	1080
gatagattat	tctaaaggat	atcatcaaga	agactattaa	aaacacctat	gcctatactt	1140
ttttatctca	gaaaataaag	tcraaagact	atgawawmaw	agttttttat	accttattta	1200
agagaaacaa	cctgacgtgc	accawtcagt	ctgcacatcc	caacccttca	cattttataa	1260
attattgtag	atcatgtttt	gttagggagcc	cttttatgga	gaggacattt	tcccatgnct	1320
taagtaatcc	agcctttt					1337

<210> 20

<211> 1162

<212> DNA

<213> Homo sapiens

9

<400> 20

ggcagcagagc	gccccggact	cttctcagtt	gagagtgcgg	ttcctgggca	ggtttccaca	60
ccagttcctt	tccgcgtcct	tcggccctgg	ctctggctgc	ctggcggagg	tggggtagca	120
tttgtcattt	gcacactgct	ggctttatct	ttggggctgc	accccgaggc	aacaaatgca	180
ggatgctctg	tcacccacat	gtccaccacc	atctggtttg	ccttttggct	actttgactt	240
tctccttaaa	tgttctctgt	gctgagcaaa	cattccacag	ccagcagagc	aatggagagt	300
tcatggccac	tcttcccagt	atcagcaagc	aatttggggg	gacggtttgg	aagcctcaga	360
ggaaagatgt	catcaggctt	cctgtggctt	tgctcttcag	catggggctc	ggcttgcttt	420
cacctgcctt	aggaagattt	ctggcttctg	agctctgata	tggggagaag	ataagggctg	480
ggatctttga	gtctgccccct	agctgggtat	gtgcgtccgg	tgtgcggggc	ttggagtttt	540
tggtaatgac	tcacttgtgc	tctttctggg	atctgtctcc	ctcccacatg	accccggtgg	600
gtccctgaat	gactgtttta	gagtacccat	gtgggttccc	tgagtcacag	caggggatgt	660
ttaataagga	ggtagcact	gagcttgggg	acgtgctgtc	acaccagcag	gacgtgcag	720
gaaggagcag	gctacttccct	ttcttgacgt	gcaaataact	cgtataggct	aatcaacagg	780
cttataagtt	aaaagggcta	ccgctcggcc	ccttggggat	tccatccctc	cctctgtaac	840
ttggagatgt	ttgtttctgc	tcgagactca	gagggttcg	atgaagagt	gtgggactga	900
gttgagaagc	ttatcccttc	gctgggtggg	aggtttctaa	ttgccctgtt	ctttggggga	960
tccttaagtc	cagcttccag	gtgggggcag	cgataggacc	aagttctcct	agtagtctct	1020
gggaagccac	ttgagggaag	ctgccggtca	tgcccatgca	cccattgggt	ttctgccagc	1080
aggccctgta	ggtcgtgcca	tgttccatgt	ccttctgggt	tcttggggga	gaaggaagct	1140
gttgaaaaaa	aaaaaaaaaa	aa				1162

<210> 21

<211> 1837

<212> DNA

<213> Homo sapiens

<400> 21

aggaaagaaat	gaataatgtg	agcattaggg	gcggcagtg	gattactcag	tgttgagaga	60
aacaacacgg	aaagcccaga	gcagaggagg	gagaaggccc	cattctctca	gctttaatct	120
actgcacctc	ggagggcgaga	gctctaacca	gacaccctgg	ataagagtcc	gctggctcct	180
ggaagctccc	tggtagaccc	ccagatctct	ccttgggtgc	tgatggccat	tctgctggcc	240
tgcttcacag	ccgtcttggc	cttcatctgc	ctccagttct	ggtgtgtccg	ttgccatgag	300
ccgcgatgg	cttacagggc	tggccacatg	gaggaggcca	atgggttgg	gagatggcca	360
gaggaggccc	cggatcttgg	tcagaggagg	gaagacctgc	aggggctccc	cctgggtggaa	420
atgccacgca	agaactccag	agatggagct	gaactggatc	ccgaagccaa	ccaggatgcc	480
cctgatgcgg	gtgccttaca	gagggggggg	ggtgaccac	ccgctatact	gcctcattgt	540
ggggaatgag	agtggggagg	agagcgccc	catcatgtag	cctaaagctt	tccacacagt	600
aggtcgtttg	tacaaatgtg	tctatagact	acccatttct	ctcccatcaa	acgtcactgc	660
tattgtagg	cacctgggtt	ggatgaatgc	cccatagaca	agcttctcaa	gctggaaaaat	720
gtatccccca	gggtatgcca	aggcacaac	raggcatctc	agcagcagcr	tccactcgga	780
ggacacagtt	aagaattgaa	agcattgttt	taacattaaa	acaaacatgg	atatgttata	840
aagtagaaag	caaaaatttg	catgacttaa	ataaaacaca	gcacttgaag	gaaacacttg	900
agtwacaagt	aggctctcag	argtacactt	ttgggcaaa	gtagatgtcc	agtttctctt	960
gcccttgagg	ggattatttg	gaaataattt	gggattattt	ggaaataatt	tgtgagaacc	1020
cttgctctgg	agtgttttcc	aacttttagg	caatcacata	ccaccgytyt	ctatttttta	1080
aaaccatgga	ttatcttcac	tattattaat	aatactttcc	tttaaattga	ytcatTTTTT	1140
aagcgtaaac	ttattttcaa	agacagcctt	atattactcc	ataagtga	aaaccagcacc	1200
ctcytgctgc	aaacagaagg	gaagagacat	aagaataaac	atcatgaaaa	caagataata	1260
ttaaataatc	tgacttagct	tctattgcct	gccagtgtat	ctgagcttaa	tgcttgcctg	1320
cttgctcttg	ataaaaagg	agatccagtg	ttggagaggt	aataaagaca	tattagcacc	1380
aatatttg	tttctccttg	atatagcata	agatttgaaa	gagaatcgaa	aagggaagt	1440
ctgtttttac	tttgcgagtc	agtgttagtt	aatgccatg	tctgtgaacc	acctcaaagt	1500
ctgtcagtc	gcttcccttg	atttgagaaa	attgtcttga	ccagactagg	caggctgttc	1560
tggagaagac	gcctgtactc	tccaagcatc	atccaagacc	ttcccagtag	cctcttatat	1620

ctatagagca	caaaatccca	gaatcacagg	acgacactgc	agtgatcaa	ctaagaaaca	1680
gcaggagctg	agaagccaag	acgagaagcc	ccacatccct	attcccttgc	ctacctcatg	1740
cattccctgc	tcggcaccca	gactttttgcc	cccattcctg	ctacttgtga	acaaataaag	1800
attcatatac	tcacaaaaaa	aaaaaaaaaa	aactcga			1837

<210> 22

<211> 1054

<212> DNA

<213> Homo sapiens

<400> 22

cacgcgtccg	cggcacgctg	ggaaagaatc	ctccagtttt	agctcctact	tcaacagcat	60
ttccttatct	atacagtaac	ccaagtggga	tgtctcctta	tgcttctcag	ggttttccat	120
ttcttctctc	atatcctcca	caagaagcaa	acaggagtat	cacttcttta	tctgttgctg	180
acactgtttc	ttcttcaaca	acaagtcata	ccacagccaa	gcctgccgct	ccttcatttg	240
gtgtcctttc	aaatctgcca	ttacccattc	ccacagtggg	tgcttcaata	ccgacaagcc	300
aaaatggttt	tgggtacaag	atgccagatg	tccctgatgc	atttccagaa	ctctcagaac	360
taagtgtgtc	acaactcaca	gatatgaatg	aacaagagga	ggtattacta	gaacagtttc	420
tgactttgcc	tcaactaaaa	caaattatta	ccgacaaaga	tgacttagta	aaaagtattg	480
aggaactagc	aagaaaaaat	ctccttttgg	agcccagctt	ggaagccaaa	agacaaactg	540
ttttagataa	gtatgaatta	cttacacaga	tgaagtccac	tttcgaaaag	aagatgcaaa	600
ggcagcatga	acttagtgag	agctgtagtg	caagtgcctt	tcaggcaaga	ttgaaagttag	660
ctgcacatga	agctgaggaa	gaatctgata	atattgcaga	agacttcttg	gagggaaaga	720
tggaaaataga	tgattttctc	agtagcttca	tggaaaagag	aacaatttgc	cactgtagaa	780
gagccaagga	agagaaaactt	cagcaggcga	tagcaatgca	cagccaattt	catgctccac	840
tataggtaaa	ttgtatttca	agtttgagtc	tcaagggtgat	tgcatcagtg	ttctttaaat	900
agacatgttg	ttaacggtgc	ctgttcatca	gccttaagca	taattctgtc	attatagtta	960
ctgtgctatg	taacatagaa	tgctttgtat	tatataataa	gcataatata	aacatataac	1020
tagtagaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaa			1054

<210> 23

<211> 1066

<212> DNA

<213> Homo sapiens

<400> 23

gaattcggca	cgagaaaaag	caaaaagtaaa	ataaatatta	aggaagaaat	gtcaagagta	60
gtctctctct	tctttttcat	tctcttttcc	tttttcttct	ttgctttctc	cttatcgtct	120
tctctttctt	ttgtacacta	tgagaaaacta	gtgcaagtga	aagaatgttt	ggattcattt	180
ctcaaaaaaa	taaaaattaa	ggaatacaag	actagacaat	gttatcattt	aataagggtg	240
gagaacaatg	gggccaagtt	gcaaaagtga	acacagggtg	aagcatcagc	aaagcagagc	300
cgtagaagaa	gtcaagcaac	tttgtccaag	gcagggtttt	aagctcacat	taggcagggtg	360
ggggctaagc	agagctcctc	agagggtcag	atgcatttgt	agatacctta	ctgctctact	420
gtctcctgtt	ccagaaaata	cctatgcata	taacattcat	aaaactttta	tatacgtaca	480
aaactagtat	cttgactgta	gtgctagtgt	gattcttatg	ttttgcaatt	catattttgt	540
aattgaaaaat	atgaaagagt	gtgtgtatag	aagaaagaag	agagaaggaa	ggaagggaagg	600
aagaagaaaag	agaaaggaga	gaaagaaaaa	gagggaaaaa	tgagattaag	gttcaagtga	660
ttgagtttgg	cgtctcctcc	caagggattt	tacattggct	ttggttcaac	tctgatctta	720
aattgcaact	gcctataaaa	tgtattttat	aaattggggc	aggggggtat	aaaatgttat	780
tttagcttgt	agaattctag	tggattggag	tgtagtaca	gatcgaaaga	actatagatg	840
ggagataatg	agaattggaa	ttaagccatg	caaagtaaaa	atctaagagg	cagctctatg	900
aaatatgact	ctactgaaca	ggtttaggag	acagcagctg	gtgaggcaaa	ctcctgctag	960
gaagaatttt	ctaattctaa	gctactttca	aatttgatag	ggctgaaaaa	tatctctctg	1020
aatcaaagtc	agacattaaa	aaaaaaaaaa	aaaaaaaaag	ccgctc		1066

<210> 24
 <211> 928
 <212> DNA
 <213> Homo sapiens

<400> 24
 ggcacttcat ctaaagtagt aactcagaaa gtgcacttat cttctgttga atttcctttc 60
 atgtcagcct ctctaaaaaa ccacctgacc cactgctttt tgctattgct gcttaaagag 120
 cttgtttccc ccaccatgat tagctttgtg cctacactaa ggcactccta cagattcttc 180
 aacctcttct catgtgatgc agaaagtaca aaggagagcc ctggccgaac tgtccagttt 240
 agtaaaacac ccagaggagt aactatgttt atttaggggtg tttcagagct tgctgctctg 300
 cctagtctat tgtaagagct ctgaaattat tagagctttt cctaaaaata tccacacttg 360
 gatgtgactc agtcatagaa aaacaactga cccaagaaac agagagtctg agttttgggt 420
 cttggtttgt tttgttctta ttgtgttttc attttgttgc atgaaacatg aaacctccgg 480
 tcttgggata gatttgagtc ctgactcagc cagtataag ttgtttgacc ttgggtgagt 540
 tactaaactt ctctggggct aaatttcttc atctaagaag tggattagat acagcaacat 600
 attatagttc gaagcttctg gctggacgtt gtggctcacg actgtaatcc cagcattttg 660
 ggaggctgag gtgggaggat ctcttgaacc cagaagtcca agactagcct gggcaacatg 720
 gcaaaacctt gtctctacta aacatacaaa aattagtcag gcatgatgga gcagcctgt 780
 aatccagct actcaggaag ctgaggtggg aggatcgatt gagcccagga gtttgagatt 840
 gcagtgaacc atgattgaac cactatactc caacctgggc aacagagcaa gacctgtct 900
 caaaaaaaaa aaaaaaaaaa aaaaaaaaaa 928

<210> 25
 <211> 966
 <212> DNA
 <213> Homo sapiens

<400> 25
 ccacgcgtcc ggtatttttg tcaatgacgc cactgaaagg gatcaagtcc gtgattttac 60
 ctcagggtttt cctctgtgcc tacatggcag cgttcaacag catcaatgga aacagaagtt 120
 acacttgtaa gccactagaa agatcattac taatggcggg agccgttgct tcttcaactt 180
 tcttaggagt aatccctcag tttgtccaga tgaagtatgg cctgactggc ccttggatta 240
 aaagactctt acctgtgatc ttcctcgtgc aagccagtgg aatgaatgtc tacatgtccc 300
 gaagtcttga atccattaag gggattgcgg tcatggacaa ggaaggcaat gtccctgggtc 360
 attccagaat tgctgggaca aaggctgtta gagaaacgct agcatccaga atagtgtgt 420
 ttgggacctc agctctgatt cctgaagtct tcacctactt ttttaaaagg acccagtatt 480
 tcaggaaaaa cccagggtca ttgtggattt tgaaactgtc ttgtactgtc ctggcaatgg 540
 gactgatggg gccattttct tttagtatat ttccacagat tggacagata cagtactgta 600
 gtcttgaaga gaaaattcag tctccaacag aagaaacaga aatcttttat cacagagggg 660
 tgtaggcgtg agtttttaggt gaatttatgt gggtcctgct tgaaaacctt cccctctcca 720
 ggttcgggtt agagaacttt gccacaggtc ttctgggggt ttttaaggctg gctggagaag 780
 acagtgggag ggtgcccccg tctgacacc ctgggggttg tgaggggacg gttggagtgg 840
 ggatcggcct gcgaaaggat actgtgaaat cactaattaa ctaataaacc tgtctcaagt 900
 tgaggatttg aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 960
 aaaaaa 966

<210> 26
 <211> 1146
 <212> DNA
 <213> Homo sapiens

<400> 26
 gctgacctca taatccgccc gcctcggcct cccatagtgc tgggattaca ggcgtgagca 60
 ccgcgtccgg ctgtcagaat ttcttttaat gctacacata tataagcaaa taatgttttt 120
 aagaagctaa ccttgatgtt aagagtggca ggtgttctcc agtttttacc tctttcatat 180

12

gggaccaaag	tagctagttt	atggaacaca	tatgaaaatg	tggttatgcc	accaagtttt	240
actactacac	ttgtcttacc	acttttaagt	catgaattct	ataattattc	ataccctttt	300
gcctgtgate	agaagtaact	tttaaaatta	tcacttgact	ttggatggaa	ctagtataaa	360
ggcaggaatt	ttgtctttca	gtggagattt	attctgttca	aggttgaagt	ggacacatca	420
ttatcttggg	atggtacttc	tttattttaa	tagtccttta	aagttctttg	agtggcagag	480
aacgtttttg	gtttcagttc	gagaaggcta	ccaaatgggt	taagttcatt	tcatagttag	540
gagaaaaaga	ttttgagtag	tctaattgtc	tcagaaagga	ttaaaacgtt	gtatgtacca	600
agaaggcaga	atgamgaagg	atcacgttca	caaatgctgt	atgtttaaca	aaatacgttt	660
agatggtaat	atccaatagt	cttatcaagt	gctacaatct	ttttaaacag	ggaatggcca	720
aatccagtg	tattgaaaca	gcctgaagaa	tgcaatctta	atttgctgt	atgggaccca	780
agggttagtg	tattattttt	tcccctacca	attcacactg	tgcaataaca	agtaaaaatc	840
atctgcataa	actccaggga	gacttccagc	cttttactta	tgaatgggtc	tgtccgtatt	900
acactttctt	ttagataacc	tcaactataa	ttgtcctcaa	ctataattga	tgtgagaagc	960
ataattatgt	accctgtaaa	ccacatttaa	ttgtacatag	tttttaatac	agattttact	1020
aacatttttaa	tttattctat	atagaactac	cttgtaagtc	aaagtgtgt	gggcatttgt	1080
gctttaaaaa	aaataaaagg	atactaaaaa	aaaaaaaaaa	aaaaactcga	gactagttct	1140
ctctct						1146

<210> 27

<211> 802

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (337)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (359)

<223> n equals a,t,g, or c

<400> 27

ggtgggtgac	cagagagtcc	tgtctatcct	aggaggagaa	cattcagccc	aaatcccagc	60
cccatcatgc	acagatcaga	gccatttctg	aaaatgtcgc	tgctgattct	gcttttcctg	120
ggattggcag	aagcctgtac	tcctcgtgaa	gtcaacttgc	tgaaagggat	cataggtctc	180
atgagcagac	tgtcaccgga	tgagatccta	ggcttgctga	gcctccaagt	actgcatgaa	240
gaaacaagtg	gctgcaagga	ggaagttaaa	cccttctcag	gcaccacccc	atccaggaaa	300
ccactcccca	agagggaaga	acacgtggaa	yttcctngaa	atgcsctac	atgggtgrtng	360
acctacctct	tcgtatccta	caacaaaggg	gactgggttca	ctttttctc	ccaagtgtta	420
ctgccaytac	tgtaaacttg	aactggacat	cagggatgat	ccctgctgtt	ctttctagt	480
agcctgctcc	atctcagctt	agccttcaca	aggcctccat	ctcccaggca	ttctaaccct	540
tgaagaaagc	tctctgtccc	ctggactgcc	tgtgtggagg	gtaatgaact	gggtccttta	600
aggaatggca	cctgggtgcc	cagaggcatg	gccagaaggt	gtctgtgggg	gccatgcctt	660
agggggatgc	accaggggcg	gctgagagag	caactgcagg	agtttcccct	aaaatctctc	720
ctccagatcg	ttctcgaact	ttcccacta	cttcataat	aaaatgtata	cttggtgaaa	780
aaaaaaaaaa	aaaaaactcg	ag				802

<210> 28

<211> 1169

<212> DNA

<213> Homo sapiens

<400> 28

ggaatgtgag	tgcaaatttg	aatttccatg	tacacatgtc	cccgtgtgca	catatctgtc	60
------------	------------	------------	------------	------------	------------	----

tgtatgtgct	agtgttttcta	tgtaatgtga	ctagatgtga	atgtgttagg	gcattcacaa	120
cctgggacac	agagaaagt	aaatatttta	tggcacactg	gagtaaactg	aagagggtta	180
ggggtactag	agttgagtga	aaaggaattt	cttacatttt	cctcatatta	tacaattatg	240
ggaagaaaat	taaaatgcag	aatttttagg	gagttattaa	atattgaatt	tgtgtacaac	300
tttcaaata	aatcttttca	gttttttatg	acacacttga	gctcacttct	agaaacatgt	360
cttagtctgt	tttgtgctgc	tctaacagaa	tacttgagac	tgggtaattt	ttaacaagca	420
gagatttctt	tcttacagtt	ctggaggcta	ggaagtccaa	ggttgagggg	catgcatcta	480
gcaagggcct	ccttcctgcg	tcatcccata	gtgaaggcca	gaaaggcaag	agaacacgct	540
tttgcattgag	agagagaaa	agagagaaga	gaagggaagg	gaagagaaa	aagagaagag	600
aagagaagag	aagagaagaa	aagggaagcca	aactcatcta	tttatcagga	acccttctta	660
tgatagaaac	ccactcccat	gaaaacagga	ttaatctgtt	tatgaggaca	gagtcctcat	720
tacctcatca	sttcttaaag	gtctcacttc	tcagtactgc	tgcattgggg	attagggttc	780
caacacatga	acttcggagg	acacattcaa	gccgtagcat	tctyccttga	ctccmmaaat	840
ccatgtcctt	ctcatgtcta	aaatagatta	atyccatccc	aatwgcccca	aagtcttgac	900
tcgttccagc	accaactcaa	aattccaaag	tccagagtc	catttgaatc	agacaggaga	960
gactccagg	gcaattcatc	ctgaggcaaa	tttctctcca	gatgttagcc	tatgaaaata	1020
gcaaattact	ttcttccaaa	atacaatggt	gggacaggca	taggatagac	attcccattc	1080
caaaagggag	aaataagcaa	gaagaaagg	gtaactggtc	ccaagtaagt	ccaaaatcca	1140
acagaaaaaa	aaaaaaaaa	ggcgccgc				1169

<210> 29

<211> 1466

<212> DNA

<213> Homo sapiens

<400> 29

ggcacgagtg	cctcaaagac	tattattttg	gaggatctag	tgcaaatgtt	agtaatgtgg	60
atattgtgta	gtgtcccagg	atattaatgt	ttttagcctc	ttggctttta	ttctgtattg	120
ttgccccaaa	agatgatgct	cacttatctt	tcatccagtg	taaggatata	tggaaagaca	180
acagaaaagta	tagctgtttt	catttcaaaa	gtgatcagct	gcttgagcta	gcaagcaagg	240
cttgactag	cttccaggcg	cagtcacgca	gtttcacagc	aggcgcggtt	ccctcggagc	300
acccagagct	gccctgcggt	agtcagcagt	tgtgctgtgg	ctgcactgcc	aggctgggtg	360
gcargtggat	cggagccagc	agatgtggct	caggaagtgc	cttcttggcc	tctccttaat	420
ctctttcaga	stctgtgggc	cettgattgc	actgtgggtt	gtttcagact	ccagtattag	480
gagactgaac	cccttggtgg	ttttttggtg	tgtgtgtgct	gagmtgggtt	gaggacatgt	540
taagcaggty	gggtgcytcc	cctgggtttg	ctccgggtgg	tacctgtggt	gtggggtggt	600
tctgagtagt	tctggcccca	ctgctggagt	atctgccay	tcagtttgtg	agatggcagg	660
gcttcacct	ggtctggtgc	ctcattttct	tcttttagcag	tgggcttaga	accaatgcag	720
attcccaagt	taagtatttt	ttctgtagct	taattattac	aggcttctgg	tacctaaagg	780
ctttcttact	ttctgtttctg	aggggaagag	aagataatgt	tgtttctccg	ccccccccgg	840
agtggcccca	ggaccttgca	tggcatttgc	agcatttgc	gcgtgcttgg	gtttgcttta	900
ctagggtgaa	agtgttgac	ccccagcac	ccacaaagc	acctctgctc	accctccggt	960
gaggttctga	ctggccctgg	gacatcacst	gctccaggat	cctatgtggc	tcatcccagg	1020
agagatgtgg	gaggggaagg	gaaaaaaggc	ttacatttgc	tgagtgggat	tcatgtagat	1080
ctgagttccg	cattgattcc	taagctgcag	agcccttatg	ccttggtgtg	tttgtgaatg	1140
ttagtcggtc	ttaacctttt	tcaccgagtt	agcatttggt	gtctcaggag	gtcacagct	1200
cctgctctc	ctccagggga	gtgcgcctc	ctcctctgtc	ggtagctgtc	aggtgcccc	1260
ttcctctgca	gcagactgtc	ctgggtcctt	gcctggcctt	ccccttacac	gtgagcctgc	1320
agcttcattc	acagcccctg	tgtagaaaga	taggcactaa	aagcagctga	ctggcagccc	1380
tagaaacatg	aagggtttca	tttatagttt	cagtcctttt	ccttctttcg	agccttaatt	1440
taaaaaaaaa	aaaaaaaaa	ctcgta				1466

<210> 30

<211> 1226

<212> DNA

<213> Homo sapiens

<400> 30

14

ggcagtgc	gc	atgg	acta	aa	ata	ag	tat	tag	att	ct	gc	ctt	ca	ag	ctt	gct	acc	ag	at	60
g	c	g	a	a	t	a	a	t	a	a	a	a	a	a	a	a	a	a	a	120
ag	ct	ta	at	ta	tc	act	tg	cc	a	ct	c	act	c	act	gat	ag	gg	ttt	tg	180
tt	g	att	tt	att	gt	gg	tc	cc	tg	tg	ca	ct	ca	aa	cct	ct	ct	gc	tg	240
g	c	ca	ct	cc	ca	ag	cc	ta	tc	at	ca	cc	tc	ag	ct	cc	ac	ct	ag	300
t	ct	ca	ta	gg	ac	at	gc	aa	cc	tag	at	cc	ct	a	ac	at	gt	gc	ag	360
c	ct	at	ga	ta	tc	ta	at	gt	tg	cc	act	ca	ct	g	ac	ag	gg	gt	gc	420
a	gt	ga	tg	gg	a	at	gg	ct	gt	a	ag	ac	ag	at	g	at	gt	gt	tc	480
c	ct	gt	ct	gc	tg	cc	tg	gt	tc	ta	ac	ac	ag	gg	gt	gt	tc	gg	ga	540
a	tg	tc	ac	cat	at	gt	gc	cc	a	tt	at	gt	at	ca	gt	at	at	at	at	600
t	t	at	tt	tt	tt	tt	ta	cc	ac	tc	ga	act	gg	tt	tc	g	at	at	at	660
g	c	act	tt	gt	g	ag	ata	ct	ag	g	at	g	at	g	at	g	at	g	at	720
t	g	a	att	ct	tt	t	gg	act	gt	ta	ct	cc	ct	ga	g	aaa	a	att	ag	780
a	t	g	a	g	a	g	a	a	a	a	a	a	a	a	a	a	a	a	a	840
g	g	g	a	t	a	c	a	a	k	a	c	ag	g	c	a	w	t	t	g	900
c	ct	ct	g	a	g	a	c	act	g	a	g	c	g	a	g	a	c	g	a	960
c	ag	at	gt	gc	a	ag	ga	ag	ag	g	c	ct	tg	gt	a	c	ta	ta	a	1020
a	tt	ct	aa	ac	ct	act	ct	gt	ga	ta	att	ag	cc	ac	ct	tt	ga	at	ac	1080
t	t	at	gg	tg	a	at	g	cc	ata	a	tg	at	tg	a	tg	g	a	a	c	1140
g	t	g	a	ta	ac	at	ca	ta	tg	ga	gg	ct	ta	ag	t	at	tt	ct	g	1200
g	c	a	a	a	a	a	a	a	a	a	a	a	a	a	a	a	a	a	a	1226

<210> 31

<211> 1094

<212> DNA

<213> Homo sapiens

<400> 31

ggcagagccc	cagg	gtg	aga	ag	ttt	ggg	ag	ac	at	gt	tt	cca	ttt	tg	gt	ct	gc	60
a	t	t	t	ct	ct	t	at	g	t	at	g	at	g	at	g	at	g	120
t	a	t	t	ag	ag	c	g	t	t	cc	a	at	gt	cc	cc	a	ct	180
c	a	g	at	gg	t	at	g	g	a	a	t	g	ct	g	a	t	a	240
g	c	ct	cc	ca	ac	c	ct	ca	t	g	g	c	ct	g	g	c	ct	300
a	a	t	ag	cccc	a	c	ac	ct	g	cc	ag	at	gt	ct	tc	ga	a	360
a	ag	tc	ct	tag	a	a	t	ag	ag	ca	t	at	ag	ca	a	a	a	420
c	t	t	t	tt	gg	ga	tg	act	aa	ag	t	at	a	a	a	a	a	480
a	c	a	t	aa	act	t	t	g	a	a	ta	act	ct	ta	ta	ct	g	540
a	a	t	a	ca	at	at	t	tt	ct	a	ag	t	g	tt	cc	a	a	600
t	ct	gt	tt	cc	ac	aaa	ac	ca	a	a	a	a	a	a	a	a	a	660
g	tt	g	a	c	att	tt	ct	g	ac	ca	ac	ac	g	g	g	g	g	720
t	ct	g	gg	gg	ac	ct	ct	gt	gc	cc	ca	cc	ca	cc	ca	cc	ca	780
t	t	c	ac	gc	ata	c	act	gg	gt	g	g	g	g	g	g	g	g	840
g	c	a	c	a	c	act	c	ag	at	c	a	c	a	g	a	a	a	900
g	t	a	act	ag	gt	ta	tt	ca	att	tc	ag	ac	gt	aa	at	tt	ct	960
t	t	g	at	ct	gt	g	a	a	t	tt	ct	ct	ct	ct	ct	ct	ct	1020
a	a	ag	tt	ct	ct	g	a	a	gg	ct	ct	ac	ag	t	tt	ct	ct	1080
a	a	a	a	a	a	a	a	a	a	a	a	a	a	a	a	a	a	1094

<210> 32

<211> 1037

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (6)

15

<223> n equals a,t,g, or c

<400> 32

ggcggnccca	tgaagactg	cgagtacagc	cagatcagca	cccacagctc	ctcccccatg	60
gagtcgcccc	acaagaagaa	gaaaatcgcg	gcccggagga	aatgggaggt	gttccccgga	120
agaaacaagt	tcttctgtaa	cgggaggatc	atgatggccc	ggcagacggg	cgtctttctac	180
ctgacgctcg	tcctcatcct	ggtcactagc	ggactcttct	tcgccttcga	ctgtccgtac	240
ctggcggtga	aaatcacccc	tgccatccct	gcagtcgctg	gcatacctgtt	cttctttgtg	300
atggggaccc	tgctccgcac	cagcttcagc	gaccccgagg	tcctcccacg	agccacacct	360
gatgaagccg	ccgatctgga	aaggcaaata	ggtaacactg	aaagtctgcc	catggcctct	420
ggtcacttcc	cgcttggg	cagctacagt	ggggaaggca	ggccgagggc	tktgaggag	480
gagctgascg	ctgggaagga	aggaggccag	aagtcagcgt	tccttagctc	gctgggtggg	540
caggatgagc	tgaagaagag	gtgtgatata	aggctggagg	gacaggatc	ctggaggcag	600
gactgcaggc	ccacttgagc	aaagcatcag	tgtgagctgt	gcttctgatg	tttctttgaa	660
acccaagtgt	ttgattccac	tctactaagc	agcctccacc	ccgagagatt	acagttgtag	720
gttgctgccc	ttactccctt	ttgcaatcct	tagagcatga	ccagtttcca	tgatataaat	780
ctaggaaagt	tacatcttag	gcagcttctt	tgtttatcca	ggccaggatt	gagaatttcc	840
tttatttagg	tataataaac	atgtatctgt	gatatgtatt	gagatgaata	gctttatttt	900
tccttagata	ttaaaaccta	tactaaagtt	tattacaacc	cattttgaag	atattaaaac	960
agatccta	cccttacaca	ataaaactttt	acagtttttt	tttttaaaaa	aaaaaaaaaa	1020
aaaaaaaaaa	actcgag					1037

<210> 33

<211> 1376

<212> DNA

<213> Homo sapiens

<400> 33

ggccaccctt	ccccaccaca	gagactgggc	agctgtgtct	ggtggactct	gcgccccgcc	60
ccctgcagcc	gtacttgccg	ctcatgcggt	tggacaagcc	cattggaacc	tggtctctgt	120
atttaccatg	tacctggagc	attggtttgg	cagctgaacc	aggttgtttt	ccagattgggt	180
acatgctctc	cctctttggc	actggagcta	ttctgatgcy	tgagagcaggc	tgtactatta	240
atgacatgtg	ggaccaggac	tatgataaaa	aggttacaag	aacagccaat	cgtccaatag	300
ccgctggaga	catttcaact	tttcagtcct	ttgtttttct	tgggggacag	ctaaccctgg	360
cactgggtgt	tcttctgtgt	ctaaattact	acagtatagc	tctgggagca	ggatccttac	420
ttcttgtcat	cacctaccaca	ctaataaaaa	gaatttcata	ctggcctcaa	ctagccttgg	480
gcttgacatt	taattgggga	gcgttacttg	gatggctctg	tatcaagggt	tcctgtgatc	540
catctgtttg	cctgcctctt	tatttttctg	gagttatgtg	gacactaata	tatgacacta	600
tttatgcccc	tcaggacaaa	agagatgatg	ttttgattgg	tcttaagtca	acggctctgc	660
ggttcggaga	aaataccaag	ccgtggctca	gcggcttcag	tggtgcaatg	ctgggggcac	720
tgagcytagt	gggtgtgaac	agtggacaga	ctgctccyta	ctacgctgcc	ctgggtgctg	780
taggagccca	tctgactcac	cagatttaca	ctctagacat	ccacagacct	gaggattggt	840
ggaataaatt	tatctccaac	cgaacactgg	gactaatagt	ttttttaggg	attgtccttg	900
ggaatttgtg	gaaagaaaag	aagacagaca	aaacaaagaa	gggtatagag	aataaaaatag	960
aaaattaatg	aatgaaat	atctaggaat	tttttaaaaca	ttttttacaa	aatataatta	1020
gatttgaata	caaaatctga	tacaatatgt	taaagaatta	agaacctgaa	gatgaagatt	1080
tagagcatat	ttacctggat	tttacttatt	tgctagcaaa	attccccctt	gtcacagaaa	1140
ccagggactc	ttcaggattt	gagatggcct	tgagtatttt	agttgatata	ttcttctgcc	1200
cattataatt	ctcacctgaa	gttatgggga	ttgcacgggt	tttggcactt	tagaaaaagc	1260
ctgatgtggg	tcttacataa	atgaatgtct	gtataagaaa	atggactctt	tttttttaggg	1320
aaaaataaaa	gcaactatgg	gaaaaaaaaa	aaaaaaaaaa	aaaaaagggc	ggccgc	1376

<210> 34

<211> 1220

<212> DNA
 <213> Homo sapiens

16

<220>
 <221> SITE
 <222> (803)
 <223> n equals a,t,g, or c

<400> 34
 ggagaagtgc ctaacatccc atgtgtgcta aggtggcaga tcctgagctc tccagctgcc 60
 ctcaactgtgg cctcacggca caaccagggc cagagtcagg aaacatcagt cactcgctta 120
 gggaaggaggag tccgagaacc ctcttcgtgg actcaacttc ccaggtctct gtccctgctg 180
 cagaatgccc aggccacaga gaagggaccc ccttttcagg agcttccacg tcacaggctt 240
 ttatgggggcc tttcttgctt gtgtttctgt ttcccatcct gaggggtgtgt ggaataatac 300
 gagagcctac ccaggactgg agcgtgttat tagaaagagc acgcttgact gctcccgggc 360
 agccacctgc cctttttccc ttggaaagtg gacctatggc cactgcacaa aacacttcc 420
 aataccaggc tgctggtgac tacattttgt ggtttcttg ttagttgata tttatagttc 480
 agatctcatc actctgccaa gtgcaataaa cattttgttt taactggaat ataactttta 540
 ttattgttat tatttttctt ttaaagtct taaaagtgc cagcttgctc aagtggagtc 600
 agaaacccta aatagtctaa atgccaaagtc agtgcgctct gtagatcagt gtacgattag 660
 aagacaggct ctcatattt actgcttgta tttatataat ggaagttagg taacataagc 720
 ccctctgcag gccagcttag atgttgatga tgcttaaatg gtaaaatctg tgagaatgtt 780
 caacaatgcc ctgtatttgt acngcctaaa actgaataat acatatgtga catgaataaa 840
 taggtcagtg atttttatga catattaatt tcctttacct aagttcctgg gaaatatagc 900
 tgaacagtta cagagaatga aaacaaacac tggcatctta caaacacctg ctttgattat 960
 ttagtctaata taggttaaat ttaactattt ccattttaac aaattgaaaa caattcttat 1020
 ctgttccctcg gtcatgcact ttatcagtg tggatgcaga tgtgtttaat atgacacagt 1080
 tgggtgctgtt ttctcagctc tgggtgggaa ataaaaagga attgagaaca tattttttaga 1140
 gtcttctggg gactttgtat ttctgggcct atagctatga aattttaagg aggaggaaaa 1200
 aaaaaaaaaa aaaactcgta 1220

<210> 35
 <211> 1346
 <212> DNA
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (537)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (880)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (1115)
 <223> n equals a,t,g, or c

<400> 35
 ggcagaggct tgtgaagggt aaagttaaaa cccctctgct tagccctgc ctccagcctc 60
 tgccaggagt aatgtgctcc catagtactc tgatccactt gtatttggtg cttccttttt 120
 tttttctttt ctttcttcc tcctttcctt tcccttcty ttcctsttcc tccattcttc 180
 cctccctcgc tcttcttcca ttcttccctc cctccctatt cctccattct tccctcctc 240
 cctccctctc acatccttta ggactcagca tcacctctc taggcagtct ttctggayt 300

accaccaytt	atgcacaaaa	cacctaagca	ytaccttatg	tggcctcatt	tatcactgct	360
taaatatttt	ttawacacgt	gctgtgatgt	ggcacatgca	ggtgtcattc	ttgagkatcc	420
actgggttatt	gccttgaggg	gatgacaact	gcccggtagg	gtwacctggg	gtgactgcac	480
ctaaaacagc	aataccaaaag	ggcccattgc	cagttctgtc	actgaccagc	tggggcntct	540
gagtatatcc	cttaaccact	ttggacctta	atttggcatc	tgtcaaataga	gatggtggaa	600
cttgaggaac	tctaaggccc	ctactgtgca	ggtcttatta	atgattacaa	cagcagcagc	660
agccagtgtt	tactgaggac	ttacaaagca	ccaagcactt	tgcctatcct	aatccttaca	720
tcaactctac	gaagttagta	tggttactat	ccctatttta	cagatgagga	gactaaggct	780
aagagagggt	atatgacttg	accacaaggt	cataataaag	aaacagattt	gaatccaggc	840
attctgactt	tactgttctt	agccacataa	tgggcacasn	ttygacacac	rgttttgtgt	900
actgtttggg	ggtcactcac	agactccatc	ccagactctg	catgaaccat	ccctgttcta	960
cattttttaag	gctcaaactg	gagtctgggt	gaaacctggg	gacagaagac	tgctatagtc	1020
acaattatta	gagggaaatg	ggtgaggacc	agtggccagc	tctgttcattg	aacctttgac	1080
aattctcaca	gagagtcttg	ctttggacag	agacnactta	cgttgctgtt	ttcagttacc	1140
ctcttttagga	ggggagagta	ggcctgagtc	atgcttcaga	cacagattaa	aatcagattt	1200
ggtaccagggt	gcagtgggtc	acgcctgtaa	tcccagcact	ttgggagggt	gagttaggag	1260
tatcacttga	ggccagaagt	ttgagagcag	cctgggcgac	atagtgagac	atcctctctc	1320
tttaaaaaaa	aaaaaaaaaa	actcga				1346

<210> 36

<211> 1026

<212> DNA

<213> Homo sapiens

<400> 36

ggcagcagcc	tctgtcacca	aacagtatga	agtatctaaa	caatgtttta	aaacaaatta	60
tttgtagcct	tcactgttgg	ctgtctccat	ggcaggtgag	ctgcgacctt	ttgtcagcag	120
cctagctgag	atggctgctg	atgcctgcag	gtataagtga	ctgtcaattt	tccttactca	180
tttatcttgc	tgtctcatgt	ttaaactaga	agaagttgtt	catctcaagt	gtcctcaatg	240
tcaatctatc	atgtttgcct	aattttgctc	ttgtacatca	catctcacag	tcaccagaac	300
atgagtagct	gtctccagggt	gcctctgtct	ctgttatctt	gcccactgaa	gggagagcac	360
ttaagccaat	ttgcaggaga	ccacagtttg	ccagagggtca	gagacagaaa	tcaccactgc	420
attttgttta	aagaatcaca	tcagaaaaga	aaataaagga	cagggaggga	aaagaaggga	480
aacctataaa	acacgcgggt	cctaaagttt	gattccaaga	tttatgacag	aatcaggcaa	540
aactaaatta	aaataatata	tgtgaaaact	ggacaacctg	aacataagtt	gatttttcca	600
gagaccaaag	aacaaatcat	tgcacaaaca	catacctttt	caaactgaaa	atgattccag	660
agttaacttc	atggacctaa	atatgaatat	taacatctca	caaatactat	ttgtaatttt	720
atccttgagc	agtatagtgg	agagggtgact	tctaagcaaa	ttattagtag	gaagtaagggt	780
taagtgggca	tattccattt	ttacatcctt	tcttctccat	ttatttactt	cctatatggt	840
taggggcttc	caccaccctt	atcctctgaa	ataacactag	agcttttgcc	atttcctcca	900
cccaaagctt	ctcagatggt	ggaccagcaa	ggcgatcaag	tttgttggtt	gtttgtttat	960
ttgtttgttt	gtttgtttga	gactgggtct	cactgtcacc	cagggttgag	tcagagtggc	1020
tgcgta						1026

<210> 37

<211> 832

<212> DNA

<213> Homo sapiens

<400> 37

gctataacca	accatgctag	caaacttcac	actgttcatt	ctgactctaa	tatcttttct	60
gctgttagtc	tgttctccgt	gtaaacatct	caagatgatg	cagctccatg	gcaaaggatc	120
tcaagatctc	agcaccaagg	tccatataaa	acctttgcaa	actgttatct	ccttccttat	180
gttatttgcc	atttactttc	tgtgtataat	cacatcaact	tggaaatccta	ggacacagca	240
gagcaatctt	gtattcctgc	tttaccaaac	tcttgcaatc	atgtatcctt	cattccactc	300
attcatcctg	attatgagaa	gtaggaagct	aaaacagacc	tctctttcag	ttttgtgtca	360

18

gggtgacgtgc	tgggtgaaat	aacagaaacc	ctcaactcca	tagattcaca	aggggagcat	420
gggtgtgtctt	ttagcagaaa	agaaactgat	gggtgtctaga	acgttttata	tttctgtcag	480
tttgttgtag	tgtatgtatt	tgagtaattt	caaaacagat	tcctaggata	gtcttttata	540
tatatataat	atatataaaa	ttcatatata	tataaaatac	gtatgggtgt	atatgtgtgc	600
atgtgtgtga	ataataacat	tgaccataaa	ttatgaagcc	tagtatattt	catatatata	660
agtatgtgta	ttttatgata	gctaattgta	tgatatttca	tttgaagaat	ttatctctct	720
ttgtaattaa	gaaattacag	catttatcag	aaaatcattg	ctgttttcca	ttgtaatttg	780
taccacatac	atgtacttaa	ctatcaaaaa	aaaaaaaaaa	aaaagggcgg	cc	832

<210> 38

<211> 706

<212> DNA

<213> Homo sapiens

<400> 38

ggcagaggtg	acaagccccg	ccaagacaga	cctgcaagtc	ttcgtctcaa	gggacctccc	60
tcatgccagg	cccctgcctc	tcacagcagc	accctttcct	ctcattgtcc	ctgttccctt	120
tttgctgtg	gatctgtttg	gccagggtcc	ctgggggtcag	gaatatttgc	aagactcagc	180
cagctccttc	ccagcccagc	ctcttggggc	tgggactttc	tcaccctgcg	gcaggcacia	240
cagatgctgg	gacccagctc	ctgcccaggt	cacagcacaa	gtgcacatca	gcactatggg	300
gcctatgtcc	tgcccagaga	cctctgctcc	ttcctgctca	catccacagt	tcagggcacg	360
gcgccctca	agaactccag	agtcacctgt	ctcatcggtc	cccagcaagt	gcctctttgt	420
ctatgatgtc	ccccttctct	gaggcctgga	cccaccatc	tttgtccctg	gggcctgtct	480
ccagccactg	aggcccgcct	tggccagggg	agaaggagct	gccgtgcgtc	ttccctgtgc	540
cccgtctccc	tgcttggttc	tcccctccct	tccttgcccg	gctgccatgg	ccaggagcta	600
agtgcctttt	tgtgtgcaac	cacttaccct	ttctctgaaa	aacctgttct	caggaaggat	660
ctgataaact	catttactct	yaaaaaaaaa	aaaaaaaaaa	aaaaaa		706

<210> 39

<211> 1347

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (83)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (334)

<223> n equals a,t,g, or c

<400> 39

gggcagccct	caggccctcc	ggcagcctgg	ccggggcccg	gtggccatgg	cagcactggg	60
gtggctgctg	gcgggagcac	atngtcaagc	ctcaacaagt	ggatcttcac	agtgcacggc	120
tttggggcgg	ccctgctgct	gtcgccctg	cacatgctgg	tggcagccct	ggcatgccac	180
cggggggcac	ggcgccccat	gccaggcggc	actcgctgcc	gagtcctact	gctcagctct	240
acctttggca	cgteccatgg	ctgcggcaac	gtgggcctaa	ggctgtgccc	ctggacctgg	300
cacaactggg	tactaccacc	acacctctgt	tcancctggc	cctgtcgggc	ctgctgctgg	360
gcccgcgcca	ccaccactt	cagttggccg	ccatgggtcc	gctctgcctg	ggggccgcct	420
gcagcctggc	tggagagttc	cggacacccc	ctaccgggct	tggcttccct	ctcgcagcca	480
cctgcctccg	cggactcaag	tcggttcagc	aaagtgcctt	gctgcaggag	gagaggctgg	540
acggggtgac	cctgctttac	gccacctcgc	tgcccagctt	ctgcctgctg	gcgggtgcag	600
ccctgggtgct	ggaggctggc	gttgcccccac	cgcccactgc	tggcgactct	cgcctctggg	660
cctgcacctc	gctcagctgc	ctcctgtctg	ttctctataa	cctggccagc	ttctccctgc	720

19

tggccctcac	ctctgccctc	accgtccaag	tcttgggcaa	cctcaccgtg	gtgggcaacc	780
tcatectgtc	cgggtgttg	tttggcagcc	gcctcagtgc	cctcagctac	gtgggcatcg	840
cactcactct	ttcaggaatg	ttcctttacc	acaactgcga	rttcgtggcc	tcctgggctg	900
cccgtcgggg	gctgtggcgg	agggaccagc	ccagcaaggg	tctttgagac	ctgggggatc	960
tcaggagcca	cctgggatgg	ccctggcctg	aatccagcct	ccgctgtggc	catagaagga	1020
atggagaaca	gggctgggca	tggtsgtca	cgcctataat	cccagcactt	ccagagtccg	1080
aggtgggtgg	atcacctgag	gccaggagtt	cgagaccagc	ctggctagca	tggcaaaacc	1140
tcattctctac	taaaaataga	aaaattagct	gggcatggtg	gcgcgtgcct	atagtcctag	1200
ctacatggga	ggctgaggtg	ggaggatcac	ttgagccctg	gagatcgagg	ctgcagtaag	1260
ccaagatcgc	atgctactgc	actccagcct	gggagacaga	gcgagacgct	gtctcaatta	1320
aaaaaaaaaa	aaaaaacg	cacgtag				1347

<210> 40

<211> 1467

<212> DNA

<213> Homo sapiens

<400> 40

gaattcggca	cgagcagagc	aagactccaa	ctcaaaaaaa	ataaaaagaa	agaaagaaac	60
atccctggca	ccctgctctc	catcatgtct	gtgtgtttac	cgctgcacct	ccctttcctg	120
atgctagcaa	aggtagccac	cagcttttgc	agatggcagc	tcacactatt	tgtgtccact	180
ttttacaaag	atgccctggt	tcatactgtc	aatgacagaa	atcaggaagc	agaattggaa	240
gcactcaaaa	agtctgtttg	acagtgcaca	tcccctcatg	ccctggaata	tcacacagag	300
agctttgcag	tgagctggaa	agtcctccag	actcaaggcc	gggcctgggs	ccacaacaga	360
ctttgacagc	cctgcagtyc	cgggctcacc	agggaargga	gccaggagtt	cagccccgc	420
agcggcccc	gctcacagca	caggcaaaag	ccacagccac	acaggatggg	gcctccctac	480
ttcagcccaa	ttgtgatttc	ctgccttgca	aatcatctct	cttgaagctc	ttcagctcct	540
cttcattttct	tacaccagca	ccagcagctg	gccttcttct	tgagccaatt	cccagaaagc	600
aatargctga	gtgtcacttt	ttcagggttg	ctgagagctg	cccatggggg	gaggcgggac	660
aatagcagca	gcrgggttac	ccagcaamsc	aactccctcc	caagcaccag	ccctgtggca	720
gggtggacttc	aaagttacct	ccctgaggaa	agcacaaatc	atgtgctgga	cctttgacat	780
cctccccgag	ggtcacacac	tgacaattac	actctgggta	taggatccac	cccgtgggga	840
ctggctaggc	accctgcaga	atcccaagag	ggccctgccg	acagctagga	cccgccaagc	900
ccattagacc	tctagcacc	ttccactgtc	cctgtttcct	ttgggttttg	ttgtgttttt	960
tgtttttag	agataagtat	cacttggtcg	cccaggctgg	agtgcagtgg	tacaatcatg	1020
gtacactgca	acctcaaact	cctgggctca	agtgatcctc	cacctcggcc	tcccaaaatg	1080
ctgagattac	aggcaccatg	cctgggtcat	attttttttt	ttttgagatg	atgtttcact	1140
ctcgttgccc	aggctggagt	acaatggcgc	gatctcggct	cactgcaacc	tctgcctccc	1200
gggttcaagc	gattcgctg	cctcagcctc	ccaagtagct	gggattacag	gcacccacca	1260
ccatgcctgg	ttattttttg	tatttttagt	agagacatag	tttcaccatg	ttggctgggc	1320
tggctctgaa	ctcctgacct	caggtgatct	acccaccctt	tagcccccaa	agtgtcggga	1380
ttacaggtgt	gagccactgt	gcccagccaa	tttaaaaaaa	tttttttaag	agatagggtc	1440
tcactatgtt	gcccaggctg	gtctcga				1467

<210> 41

<211> 914

<212> DNA

<213> Homo sapiens

<400> 41

ggcacgagtg	aaaatctact	ctatcagctg	ctgtgggttg	caccattctc	aggaccctcg	60
ccatgaaaag	ccttatgctg	ctcaccctgt	ctgttctgct	ctgctgggtc	tcagctgaca	120
ttcgctgtca	ctcctgctac	aaggctccctg	tgtgtggctg	tgtggaccgg	cagtcctgcc	180
gcctggagcc	aggacagcaa	tgcttgacaa	cacatgcata	ccttggttaag	atgtgggttt	240
tctccaatct	gcgctgtggc	acaccagaag	agccctgtca	ggaggccttc	aaccaaacca	300
accgtaagct	gggtctgaca	tataaacacca	cctgctgcaa	caaggacaac	tgcaacacgc	360

26

caggacccccg	gcccactcca	gccctgggcc	ttgtcttctt	tacctctctg	gctggccttg	420
gcctctggct	gctgcaactga	gactcattcc	attggctgcc	cctctctcca	cctgccttgg	480
cctgagcctc	tctccctgtg	tctctgtatc	ccctggcttt	acagaatcgt	ctctccctag	540
ctcccatttc	tttaattaaa	cactgttccg	agtggctctc	tcattccatcc	ttcccacctc	600
acacccttca	ctctcctttt	tctgggtccc	ttcccacttc	cttccaggac	ctccattggc	660
tcctagaagg	gctccccact	ttgtcttcta	tactctgctg	tcccctactt	gaggagggat	720
tgggatctgg	gcctgaaatg	gggtcttctg	gttggtccca	gtgaaggctc	ccacaaggac	780
ctgatgacct	cactgtacag	agctgactcc	ccaaatccag	gctcccatat	gtaccccatc	840
ccccatactc	acctctttcc	attttgagta	ataaatgtct	gagtctgaaa	aaaaaaaaaa	900
aaaaaaaaaa	aaaa					914

<210> 42
 <211> 1131
 <212> DNA
 <213> Homo sapiens

<400> 42

ggcacgagga	ttcttctttt	atatactgat	tacaagactg	acacctatca	agtatgatgt	60
gaatctgatt	ctgacagctg	tcactggaag	cgctgggtgga	atgttcttgg	tagctgtgtg	120
gtggcgattt	ggaatcctct	cgatctgcat	gctctgtgtt	ggactagtgc	tggggttcct	180
catctcgtca	gtgactttct	ttactccact	gggaaaccta	aagatttttc	atgatgatgg	240
tgtattctgg	gtcactttct	cttgcatagc	tatcttcatt	ccagtagttt	tcattgggctg	300
cctaagaata	ctgaacatac	tgacttgtgg	ragtcattgg	gtcctattc	ggtgggtttta	360
agccattgac	agttactggg	ccacaagcct	ttcctacatc	actttgaacg	tactcaagag	420
agcgcacaac	aaggratttc	cacagagctt	tcacaaatgt	gccttttcaa	actaatgact	480
tcattatcct	ggcagtatgg	ggcatgctgg	ctgtaagtgg	aattacgtta	cagattcgaa	540
gagagagagg	acgaccgttc	ttccctcccc	acccatacaa	gttatggaag	caagagagag	600
agcgcagagt	gacaaacatt	ctggacccta	gctaccacat	tcctccattg	agagagaggg	660
tctatggccg	attaaccacg	attaaagggc	tcttccagaa	ggagcagcca	gctggagaga	720
gaacgccttt	gcttctgtag	atgcccaggg	gcttgggtcag	tgtgcctcag	ctttggagtt	780
catgcctgga	gtggttcaac	agtctctggg	gcaagtctaa	taagagatca	ggcatatata	840
tctgttcttt	gcataatatt	atgggtgccct	tattgatata	tggttaaggg	gtactagggg	900
attaggatga	ttgtaagaga	atgagaaaaga	tgaccaaag	gttgggtgga	gggaggcttt	960
ttcttatttc	caaatacttg	agaaattacc	ttttgggtta	caaattctatg	atcaacttat	1020
tccattaaat	agatacatta	aaaaaattaa	aaactgaatt	cttctgtcag	aaaaaaaaaa	1080
agaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	a	1131

<210> 43
 <211> 1333
 <212> DNA
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (411)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (1264)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (1319)
 <223> n equals a,t,g, or c

<400> 43

21

cctggaacac	ttcaacaacc	agtatccagc	cgagaggtg	gtgaactttg	gcacctgggt	60
cctcttcagc	ttccccatat	ccctcatcat	gctgggtggc	agctgggtct	ggatgcactg	120
gctgttcctg	ggctgcaatt	ttaaagagac	ctgctctctg	agcaagaaga	agaagaccaa	180
aagggaacag	ttgtcagaga	agagsmtcca	agaagaatat	gaaaaactgg	gagacattag	240
ctaccagaa	atggtgactg	gwtttttctt	catcctgatg	accgtactgt	ggtttamccg	300
ggagcctggc	tttgtccctg	gctgggattc	tttctttgaa	aagaaaggct	accgtactga	360
tgccacagtc	tctgtcttcc	ttggcttcct	cctcttccct	attccagcga	nagaagccct	420
gctttgggaa	aaagaatgat	ggagagaacc	aggagcactc	actkgggacc	gagcccatca	480
tcacgtggaa	ggacttccag	aagaccatgc	cctgggagat	tgtcattctg	gttgggggag	540
gctatgctct	ggcttctggg	agcaagagct	ctggcctctc	tacatggatt	gggaaccaga	600
tggtgtccct	gagcagcctc	ccaccgtggg	ctgtcaccct	gctggcatgc	atcctcgtgt	660
ccattgtcac	tgagttttgtg	agcaacccag	caaccatcac	catcttctctg	cccacctcgt	720
gcagcctgtc	tgaaacgctg	cacattaacc	ccctctacac	cctgatccca	gtcaccatgt	780
gcattctcctt	tgcagtgatg	ctgcctgtgg	gcaatccccc	taatgccatc	gtcttcagct	840
atgggcactg	ccagatcaaaa	gatatggtga	aagctggcct	gggagtcaac	gttattggac	900
tggtgatagt	aatggtggcc	atcaacacct	ggggagttag	cctcttccac	ctggacactt	960
accagcatg	ggcgaggggtc	agcaacatca	ctgatcaagc	ctaagcccaa	gtgtacaaa	1020
tggcccaacc	acaggagctg	ccagtatcca	gcagtatctg	gaccacaggc	aaagaaaacc	1080
actaggacca	ccaggagcac	acaacccag	acccacgccc	gagggcattc	ctccaccaga	1140
agattccgcc	acctcaagtg	aactgcagga	atcctccaac	aaccacaaac	acatgcttcg	1200
ctgttagtgt	cttcttctctg	ccctcagcac	cacagctcaa	gaaaacctaa	agtttcaata	1260
caanccatag	gtcacacaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaana	1320
aaaaaaaaaa	aaa					1333

<210> 44

<211> 1004

<212> DNA

<213> Homo sapiens

<400> 44

gcttcaactgg	agttctaagt	tttctcccct	ctgttttgaa	tgagtcagct	ctgcttctca	60
ctactgcttt	cttccacatg	ccacggaggg	gttgccagcc	tcttgacctc	agaccttagc	120
tctcagtcct	atcgtttctc	catctgcact	aatgtgaatc	actctaagta	ttctagtctc	180
tgatgtgttt	tgaaggcaga	agcagtcaga	gggcactgct	caccaggctg	ggctgggcag	240
gcagatcaca	cggaagccct	gccctgtcac	agttgttaat	actgcaggga	gatgggtggg	300
agacactatg	gggaacttga	ggagtcatgg	ttcacaatgt	acttctaaac	cactgtgagt	360
ttttttgtct	ttgtcttttg	aatataatac	tttattgtctg	ggggataatg	agtatttact	420
ttaaaaaaca	gatgcatttc	taagtccctc	tgttttgtct	tgacttccag	ctccccaaca	480
tactcacatt	ccactactta	ttctctatct	taactttact	gcttctttta	ctttttttta	540
gttttacttt	tattttttat	ttttttgaga	cagartcttg	ctctgtcaca	caggctggag	600
tgcaatgacg	cgatttttggc	tcactgcaag	ctccgcctcc	caggttcatg	tcattctcct	660
gcctcagcct	cccaagtagc	tgggactaca	ggtgcccgcc	accacgccct	gctaattttt	720
tgtattttta	gccgargcag	gtggatcacc	tgargtcagc	agtttgaaac	cagcctggcc	780
aacatggcga	aaccccatct	ctactaaaaa	tacaaaatta	gcagggcgtg	gtggtgcact	840
cctgtaatcc	tagctacttg	ggaggctgag	acaggagaat	cacttgaacc	caggagccag	900
aagtcgcagt	gagccgtgat	catgccattg	cacccagccc	tgggcaaaaa	gagcgaaact	960
ccatctcaaa	aaataaaaac	aaaaaaaaaa	aaaaaaaaact	cgag		1004

<210> 45

<211> 1494

<212> DNA

<213> Homo sapiens

<400> 45

>>

gtcgacccac	gcgctccggcg	gcggcaggcg	cgggcgaggg	ccacggggag	aggagacgca	60
gccccgcggg	tggcacgctc	ggccggggccc	cggccccgcg	tcaacggggcg	cgatgctctt	120
ctcgctccgg	gagctgggtg	agtggctagg	cttcgccacc	ttcgagatct	tcgtgcacct	180
gctggccctg	ttggtgttct	ctgtgctgct	ggcactgcgt	gtggatggcc	tgggtccggg	240
cctctcctgg	tggaaactgt	tcgtgccttt	cttcgccgct	gacgggctca	gcacctactt	300
caccaccatc	gtgtccgtgc	gcctcttcca	ggatggagag	aagcggctgg	cgggtgctccg	360
cyttttctgg	gtacttacgg	tcctgagtct	caagttcgtc	ttcgagatgc	tgttgtgcca	420
gaagctggcg	gagcagactc	gggagctctg	gttcggcctc	attacgtccc	cgctcttcat	480
tctcctgcag	ctgctcatga	tccgcgcctg	tcgggtcaac	tagcctcacc	gaggtgccgg	540
agagggagcg	ctggacaact	agaatgttga	cctcgagccg	aggccctact	tgcagcgcac	600
cggaggagag	gctctctagt	ctgaaggcac	cgcgggcttg	cgccgagctg	agtgcggggt	660
ttccctattc	caatcctgtt	tgaatgggtt	tcttcagcag	ggcttaaaag	agcagccttc	720
atcctgaaaa	tgtatttcct	tttgtttaat	gctttgagta	gataatcctg	aattgaggtc	780
atgaggaggc	ccccaggcc	agacagtcct	gaacccctct	gacacttgga	aactgaatat	840
aagtaaaatg	tccaggtgga	ctctgagtat	ttcctgtgga	tcctgggaaa	gtactgttgc	900
acaaaggctg	caaagctgga	ctcaggaatg	tcctccaacc	agcagcgctg	acctaagagc	960
tccctgtgcc	gtctatccag	accagacttc	ggtagatgcc	tttgtagat	ctatcacatg	1020
taaacgagct	tgtatctcct	tccctgtgcc	acgagagaga	ttggcttttt	attccagtct	1080
aggcagagac	agaagaatgt	tgaataagag	cacgattaga	gtcctgtctg	gttatctggt	1140
gccaagaaa	agaactctgc	tgtccaggca	ctgcttggct	tactatccca	gcaaagactg	1200
cagttttgtg	gacttttgac	caccttgggc	tggcactctt	agcacacctg	agacagattt	1260
aagcctccct	aagagactga	agagaggaac	agggtgtcaga	tactcatagg	cactgagatc	1320
tacaaatggg	aagcttgtga	gtggcccac	tttgttggcc	tacgaacttt	ggtttgatgc	1380
cagtcagggtg	ccacatgaga	acctttgtctg	agatgcaaat	aaagtaagag	aatgttttcc	1440
tgaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaagggc	ggcc	1494

<210> 46

<211> 1166

<212> DNA

<213> Homo sapiens

<400> 46

gagacctgcc	catcgaatgg	aattgagctt	cgtcaggcgc	ctacttctct	ttacattctt	60
cttctccaca	ttcagcccac	ccccaccac	cccattgctt	gaaggctcta	tgtcctgcct	120
gccttctcca	ytaayyaaga	acacggcggg	ctcccaaacc	aaatccctaa	gggagatagg	180
aacgggaata	tcagacactc	acgtatcacc	ttccctgct	caagctccac	tttgcagccg	240
gagtcaccac	tgggattcat	cagatccaaa	ctccatggac	tagttaggcc	tggaaaagac	300
cttagaggta	ggaggctctt	acagcttagc	aaacattctt	tgagtacctg	ctatatgttg	360
agggtgggaga	cttataaaca	agtaagttag	actgcagtgt	gaaaatgaca	gtgtagatga	420
taaagctgaa	gcttaagcga	tgttgacaca	tagacttctg	ttagctctag	ggctttttcc	480
tgccttaact	gacagacttt	gacaaatctt	ctaggactgt	gtgagggaat	aatatagatg	540
atcgaaaagg	gaattctttt	aagacttggt	aaagaagcta	cattggagaa	ctgaccttcc	600
catgtactga	ggacggcctg	tgtgtgtgta	tgtgtgcaca	tgggcatgtg	cacacatgtg	660
tgtaggaacc	tgtgtacatt	tcttttatca	ttctcctgat	gcctttgtta	gtgtttcttg	720
ctccctgcct	gttcttaata	aaatccacag	tttatcttgt	agcttracct	ttcttgaggc	780
catgctacat	ttcaattttg	gtataataga	aagatgttgt	caaagctgat	gggggctagg	840
tcctaagctc	attgaagact	ggaggetgtg	ccttgttttc	ctctacatcc	cctaagcctt	900
ttgtacagtg	tgtggcacct	agaggggtgt	tactgtcagc	actgaattgc	gccagtgcc	960
caccagggac	ctgcacctct	catcaggaat	gactttggga	ttgccagctg	ttctatctgc	1020
tcctctcagc	atgaacacta	agcagagtgc	cttgttttta	ctgcacttgg	gagccatate	1080
ctgttgccct	tttggatgat	tcgggggaaa	ttccttgtct	gtggttccgt	atgtgttgca	1140
ggctgagggt	tggcttgctc	cctcga				1166

<210> 47

<211> 1536

<212> DNA

<213> Homo sapiens

33

<400> 47

ccacgcgtcc	ggggcagaaa	ggaaccgggt	tgtcttgggc	cgggcagggc	gggtggtgac	60
tctcaaaagg	aaataggatc	atggcagcag	atgatgacaa	tggatgatgga	acaagtttat	120
ttgatgtctt	ttctgcttct	cctcttaaga	acaatgatga	aggctcactg	gacatatacg	180
ctgggttgga	cagtgtctgt	tctgacagcg	cttccaaatc	ctgtgtacca	tcaagaaatt	240
gtttggactt	atatgaagag	atcctgactg	aagaaggaaac	tgcaaaggag	gcaacatata	300
atgatttgca	agtagaatat	ggaaaatgtc	aactacaaat	gaaagagctg	atgaaaaagt	360
ttaaagaaat	acagacacag	aatttcagct	taataaacga	aaaccagtct	cttaagaaga	420
atatttcagc	acttatcaaa	actgccagag	tggaaataaa	ccgcaaggat	gaagaaataa	480
gtaatcttca	ccaaaagatt	gtcctgagtt	tccacatttt	cgaataatc	ataaaaactgc	540
aaggacattt	gatacagtta	aaacaaaaga	tcttaaactc	agatctccac	atttggtatga	600
ttgttcaaa	actgatcacc	agagctaaaa	gtgatgtttc	taaagatgta	catcatagca	660
cttcaactgc	aaatctggaa	aaggaaggaa	aaccacattc	tgataaaagg	agtacttcac	720
atttacctac	atctgttgag	aaacactgca	ctaattggtgt	ttggtcacgt	tctcattatc	780
aggttggcga	gggtagctca	aatgaggata	gtagaagagg	aagaaaagat	attagacata	840
gccagtttaa	cagaggaact	gaaagagtac	gaaaagactt	aagtactggc	tgtggtgatg	900
gtgaaccaag	gatattggag	gctagtcaaa	ggctacaagg	gacatcctga	gaaatatggt	960
aaaggtgaac	caaagactga	aagcaaaaagt	tcgaagttaa	aaagtaactc	agattctgac	1020
tataaagggtg	aacgcattaa	ctcttcttgg	gagaaagaga	cccctggaga	aaggctcacac	1080
agtcgagtag	actctcaaa	tgacaaaaaa	ctagaaagac	aaagtgaag	atcacaaaat	1140
ataaatagga	aagaagttaa	atcacaaagc	aaagaagaaa	gaaaagtga	tcaaaaacct	1200
aaatcagtag	taaaggacca	agatcactgg	agaagatctg	aacgagcatc	acttcctcat	1260
tccaagaaatg	aaataacatt	ttctcataat	tcaagtaaatt	accatctaga	agagagaaga	1320
ggatgggaag	attgtaaaag	agacaagagt	gtaaacagtc	atagttttca	agatggaaga	1380
tgtccatctt	ctctttcaaa	cagtagaact	cacaaaaaca	ttgactctaa	ggaagttgat	1440
gccatgcac	agtgggaaaa	tacaccttta	aaagcagaaa	gacatagaac	tgaagataag	1500
aggaaaagag	aacaagaaag	caaaaaaaaa	aaaaaa			1536

<210> 48

<211> 1038

<212> DNA

<213> Homo sapiens

<400> 48

ccacgcgtcc	gagacattta	aactagattc	ccagtcctct	ccttcaaaag	cttgggtcttt	60
gtttttccta	tagggaaaaa	agtcaaaaata	agttccaaaa	actatcctca	aagtagtatt	120
gtgctttag	taaatgaagg	ttggatggat	ggatactgac	aatgggtggca	ggcatttcaa	180
gcctttttaa	ttagtacttt	ttgtcgtctt	gcttattaaa	atthttgttaa	tttagcaaaa	240
gaccaattgt	tgtgataaac	tggtgttttt	tggatgcttc	aagcacacgt	taaccaattt	300
tttaattccc	cttttggttc	ctcccattgt	tctaaaatag	gactttcata	ttattaaaac	360
ctcaaaagat	gatccacca	ggatgaacaa	agatcaccaa	ggggaaagaa	aacatttttt	420
atctttacag	aaaacatgtt	aagattatat	atagatgtat	tctttacatt	ggatattgta	480
ttagagtctt	ccttacaaga	aatgaaatag	tttttagcac	tcttagcatt	agagttccta	540
gattgggtgt	gatagctaca	gttttaaaat	gtataacctg	aaaatgaagg	ttaattttgc	600
attgtaagag	cacatttgat	ctatgtaaaa	agtgtccatt	tgggtgtattt	ttttaaaaaa	660
gagaaagcac	tttcatatta	agtagcatgt	gtatgaattt	agattttcat	atttgttgtg	720
tctgtattca	gtgaagttaa	ttgagcattt	aaatgtttgt	tgatggcaac	attaactatt	780
aaattaaagc	accttatact	ctgctgctta	acttgcttgt	aattgcacct	ttgttacctg	840
cacattttca	tatagaatat	tgttgtaaca	ttgcttcatg	tgggtctgga	tgggaagatta	900
gtgggcctac	aggatcattt	atttatattg	tttatattac	aataatatat	tgtagatcag	960
ttgtaagttc	atttctttac	aaataaaaagc	ctcttccatt	tgaaaaaaaa	aaaaaaaaaa	1020
aaaaaaaaaa	aaaaaaaaaa					1038

<210> 49

<211> 1176
 <212> DNA
 <213> Homo sapiens

H

<400> 49
 ccacgcgtcc gaggacaaaa aacaggcctt ttaaaattct gttcagaacc agaagatgat 60
 caagaaaggg aaatgcccac tagcagcttc tactgtaagg ttagagctga cagagaaaga 120
 actccttaac tactatttgc ttcattctct acaaaggaaa ctagagaagt gggttgatgt 180
 aatagaaaga acatgtgttt gtggggccag gcaaacctgg gtttaattct gtttcaacac 240
 tgcttaacaa aatttatggg aggctattgt tttggattgg gtcctgcac caggccccctc 300
 cgggaccaa ccaaaatgga gtcactcata ctaaaactcc aggtcactga accaaaaacta 360
 agttgtttta tctgaccttc caagaaatca ggaggagaa aacaaccaa tctccaaaca 420
 ggccagtttt aatcagcgtg ataaggaagt cctctctttt ttaaccctat aaagaaagta 480
 actttttgaa atgatcaata cactttgtat tccttagttc tgctttcttt agcccccttc 540
 tgcctataaa gccacttcc tctgctcaac ttactgaagc agtattccat tttatagaat 600
 gagatgctgc ccaattctgg aatcactaat aaaagccaat tagatcttta catttggtga 660
 aattttgtct ttgacaacat tactagttaa attattctgg gtatcctttt ccccatctgt 720
 aaaatggaca tagcgatata cttcccatca gatttttctc attaatagaa gtaatacatt 780
 caaaacactg agccaggtc acaccagtga gcaacttggt aatattactc agaagcatat 840
 tatagatatt gacagaaagc aatactgttc ctaattaggg gaagaaattc taagagagta 900
 cctaagagtt tgaatttaga ttataacatt tgctctgagt attttacatt acagcctttg 960
 gggggaaaaag tacaatatgag atctgagaac agtggtactc atctttgagg aattatggaa 1020
 aacgtaatag aacactaaac atgggaaaaac atcggccttc aggttgaaaa gtggaaatct 1080
 caatccctga atttttttt ttttttacta agtaactttt ttgccattg gtgtcattta 1140
 accaaaagaa gaagaaattc caaaaaaaaa aaaaaa 1176

<210> 50
 <211> 731
 <212> DNA
 <213> Homo sapiens

<400> 50
 ccacgcgtcc gagacgccac ctgggcggac agccaggagc tctccatggc caggctgcct 60
 gtgtgcatgt tccctgtctg gtgccccttt gccgcctcc tgcaaacctc acagggtccc 120
 cacacaacag tgccctccag aagcagcccc tcggaggcag aggaaggaaa atggggatgg 180
 ctggggctct ctccatcctc cttttctcct tgccctcgca tggtggcct tcccctccaa 240
 aacctccatt cccctgctgc cagccccctt gccatagcct gatattgggg aggaggaagg 300
 ggcgatttga gggagaaggg gagaaagctt atggctgggt ctggtttctt cccttcccag 360
 aggggtcttac tgttcagggt tggccccagg gcaggcaggg gccacactat gcctgcgcc 420
 tggtaaagggt gacccctgcc atttaccagc agccctggca tgttccctgc ccacaggaat 480
 agaatggagg gagctccaga aactttccat cccaaaggca gtctccgtgg ttgaagcaga 540
 ctggattttt gctctgcccc tgaccccttg tccctctttg agggagggga gctatgctag 600
 gactccaacc tcagggaactc gggtaggctg cgctagcttc ttttgatact gaaaactttt 660
 aaggtagggg ggtggcaagg gatgtgctta ataaatcaat tccaagcctc aaaaaaaaaa 720
 aaaaaaaaaa a 731

<210> 51
 <211> 1437
 <212> DNA
 <213> Homo sapiens

<400> 51
 cggccgagcg cggaactgag agctctcagc gggagccgag acggtgcagg gccggagaag 60
 caccctcact cccagcctgc gcccgatgc tgcgcgttct gtgcctcctg cggccctgga 120
 ggccccctcg ggccgcggc tgcgcttcag acggggcggc cgggggctca gagatccaag 180
 tgcgcgccct ggccgggtcg gaccaaggga tctagagat tctgatgaac agaccttctg 240

25

cccgcaatgc	cttggggaat	gtcttcgtca	gtgagctgct	ggaaactctg	gccagctgc	300
gggaggaccg	gcaagtgcgt	gtcctgctct	tcagaagtgg	agtgaagggc	gtgttctgtg	360
caggtgcaga	cctgaaggag	cggaacaga	tgagtgaagc	agaggtggg	gtgtttgtcc	420
agcgactccg	gggcctgatg	aatgacatcg	cagccttccc	tgcacccacc	attgcggcta	480
tggatgggtt	tgcccttggg	ggaggcctag	agcttgccct	ggcctgtgac	ctccgagtgg	540
cagcttcctc	ggcagtcacg	ggactgattg	agaccacg	agggctcctc	ccgggggcag	600
gagggactca	raggctgccc	cgctgtctgg	gggtggccct	ggcgaaggag	ctcatcttca	660
cgggccgacg	actgagtggg	actgaggccc	acgtactggg	gctggtgaat	cacgctgtgg	720
cccagaacga	ggagggggac	gccgcctacc	agcgggcacg	agcactggcc	caggagatcc	780
tgccccaggc	ccccattgcc	gtgcggctgg	gcaaagtagc	cattgaccga	ggaacggagg	840
tggacattgc	atctgggatg	gccattgaag	ggatgtgcta	tgcccagaat	attccaaccc	900
gggaccggct	agagggcatg	gcagccttca	gggagaagcg	gactcccaaa	tttgttggca	960
aatgaccccc	attttaacct	tcagcatggg	agatgcatgc	cctgaagagc	aggatccaga	1020
aggaagattt	gtggccagat	tgcccttcac	atttcacctc	tccagacttc	catttcttca	1080
caaggatgat	gatggaaata	aaatgactgg	cgtgatgcct	ggaaccaagg	tgctgaccc	1140
accacctaact	gctaccttcc	ttagcttcac	cctggctaga	aataatcacg	agggttgggt	1200
ttgcttttga	aaatgcctgt	ctctctactt	gaatgataaa	gaattaaatt	agatctctct	1260
gagtccttgg	atcattggct	ctcagccctc	gacctctctc	agttatcagg	cactcattag	1320
agatgtcaga	agattttaag	atacccttag	tttcttccct	tggaacaaca	gaggtaataa	1380
ataaactctg	gacatcggtt	gaaccagtgt	caggggtcag	actgcagatc	ccagtct	1437

<210> 52

<211> 1369

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (3)

<223> n equals a,t,g, or c

<400> 52

agncagacgt	agaacgtagt	ggatccccag	gggtgcagga	attcggcacg	agatttgata	60
cccagtgcca	tattgtccct	aagaaagggt	gcacaaat	acactccac	caagagtgtg	120
ggagagccat	ttgccacata	gactcaccag	tattttttt	ttaattttt	atccatttgg	180
aagttatttg	ggaacttggg	tgtttttccc	caaaagcaaa	ggcaattgcc	tcaacaccag	240
ttatcaaagg	atccctacag	atctattttc	cctgtagatc	tgaaatgctg	tcttcattgt	300
atcttttgg	gatgcccccg	tacaaattta	caggtgagct	ccatcctcct	gtagcagcca	360
cctgcctaact	gacagtccta	ctcgggtgtc	tgataggtgt	ctcaagtgat	ggatggatat	420
gacagtagag	tccttgattt	actgtcccat	ggcccttgc	catcttctct	ttcttgattg	480
atggtgccat	catctacce	gttactttga	ccaaagcaat	gggagtcac	ctggattcct	540
tkcttlytct	ccacttctaa	gccatcagta	tactggcact	gtgttaaagc	catatttcaa	600
accagaccac	ttgtcaccat	tcctgtcact	tctacctctt	ttatcccaac	atcacctctt	660
gggtagacca	ttgcaaactg	gaagtgcatt	tccagaatat	tcttgttgg	atagaaccct	720
ctgtctacat	agtagctaga	gcagtttttt	tttttttttag	atgttaaaca	gatacatcgc	780
tctgctaact	aaaacccttt	aaagtgtttt	ccatctcaat	tagaatagaa	ttcatagtcc	840
tcaccagcca	ctgcaagggt	tatataatct	agcccttgcc	tatcttccct	gcctcttctc	900
tgttacccac	aacctgcttt	ttgtttcatg	atgtgtgaac	tcatttcaac	cttaggggtcc	960
tgctcttgcc	tcttcccttg	cctagaatgc	ttttcccttg	tcctaaatca	tctgtgttat	1020
gctagttttt	argtctcaac	tcatgtcacc	ccgttgscct	ttatctcatt	gtctggcttt	1080
atcttctcta	aaacacttgg	cactatgtar	atgttctatt	tatttactta	ttttaagggt	1140
agaatcttta	tctgttttgc	ttggtgccaa	ttattcaaca	tggtgaatag	tgccctggcac	1200
ctagcaggca	ctggagccta	tttctggaat	ttcatgttgc	accattgccc	tctctgtttg	1260
ttctccatta	ctaaattcct	ttcaagccaa	ccccatggcc	tccatgaact	tttcaaaaaa	1320
aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaactcga		1369

<210> 53
 <211> 1037
 <212> DNA
 <213> Homo sapiens

<400> 53
 ggcacgagct cgtgccraat tcggcacgag ggtcatagtc cacagaggta aaagttaaca 60
 attctgatgc tcttgatgt gcataccaga ggctctaggg aagaattccc tctttctttc 120
 ttccaccttc ttgtggctgc tggcattctt tggcttggtg tcacatcact cctatcttga 180
 aggccagcat cttcaaactt gtttcttctt cacatagcct tctgtgtgtg cagtgcctc 240
 tacctctctc ttataaagac atttgtgatt aaatggaggg tttaggataa tctcgtcaag 300
 atccttaact taatcacaac tgcaaaaacc tctttcccaa ataaggtaac attcacaggt 360
 tccagggtat aggacctatt atctttggta agtattatc agcctaccac aatagctaaa 420
 acaattctga aaaagaagaa taaagtgaga gaaatcagtt tatctgatt cgatacttat 480
 tgtatagcta tggtaataa ggctgcatgg tattaagaa aggacatata tgaatgaaac 540
 agaatagagg acccagaaat agaccacac aaaggagccc aaattatctt taaccaaggt 600
 agaagacaat ttattggagg aaagacagcc ttttcaacaa atggtactat aacaattaga 660
 tatccatagg caaaaaaaaa aaaaagaatc ttgatctaag gtcacacct tatataaaat 720
 aatattaaac tcatggccag gcacagtgac tcatgcctat aatcccaata cactgggagg 780
 ctgaggcaag agtatcact gaggccaggg gttcaagact agcctgggca acacagtga 840
 actctatctc tacaaaaaaaa ttataaacta gctgggcatg gtggcacatg cctgtagtca 900
 caactactca cgaggctgag aagatcactt aagctgagtt gttcaaggtt ctaatgagct 960
 acaatcgtgc cactgcactc cagcctaggt gacagacaaa gaccccatct caaaaaaaaa 1020
 aaaaaaaaaa actcgta 1037

<210> 54
 <211> 1373
 <212> DNA
 <213> Homo sapiens

<400> 54
 ggcacgagag caaagagaag tgtccaccag gcccgtgca ccaacgctgc gtctttaaca 60
 gctctggggc tgggcgcgtc atggctacga gaaagcgcac gaccacttcc ctggttggtt 120
 tgggttggtg tgtgtgtcag ggccagggg tttctgcttt cactcaagtt aatttatctt 180
 ccttttctt ggtaattgtg aaaaaacaaa ataaacctc ctgtgagcct tttggaactt 240
 ctggaaaagt ccctttgctg tgarcsgctg actctgagaa gagctttgag cagggttgga 300
 aaccattttt ctgcaacctt ttctttcttg gggatgtct ggggtgcacac aggcctccca 360
 caaggcaaaag gctgtccctg gatggttggc aaaatgcgcc acaccagagt ggggttggtg 420
 tggcaggagg catgaraaaa ccttgctgat ggcaggggag gacggcgaca ccacgatggg 480
 aacaaaatcc tctccttac ytctaattac aaagaggaaa aagtcactga aaaaaaagt 540
 ttaaaatgtc ttaataaag agtcataat aatccaaagc taccaaaggc caagtgttta 600
 gggggaagtt tctggtggtt aaccctactt cagggggatt taaagtggtt gtggtgagga 660
 tttggttcca ggtatgcgtc ctgccaacct ggggtgggtg tccctttggt ggagcctctt 720
 gaaaaatgar ggartggctg ggtgcagtgg ctcatacctg taatccasc actttgggag 780
 gccgaggcgg ccagatcacc tgaggccagg agttcaaaac cagcctggcc aacatggtga 840
 aacttcatct ytacaaatat acaaaaatta gctaggcatg atggcaggct cctgtaatcc 900
 cagctacttg ggacgctgaa scaggagaat ctctgaacc caggagtagg aggttscagt 960
 gagctgagat tgtgccactg cattcccgctc tgggcgacag agcaagactc catctcaaaa 1020
 aaaaaaaaaa aaaaaaagag ggagtggctg ggtgcagtgg ctcatgcctg tatcccagca 1080
 tattgggagg ccaaggaggg aggtatgctt gagcccagga gttccagatc agcctgggca 1140
 atctsscaaa acccgggcag tacaaaatcc ataaaaatta gcggggcatg gtggtgcgcg 1200
 cctgtagttc cagctacttg ggaggctaag gttggagagt tacttgagcc taggaggttg 1260
 aggcctgcagt tagccatgat tgtacctctg tacgccagcc tgggtgacaa agcaagagcc 1320
 tgtctcaaaa ccaaaaccaa acaaaaaaac aaaacaaaaa aaaaaaactc gta 1373

<210> 55

<211> 1347

<212> DNA

<213> Homo sapiens

17

<400> 55

```

ggcacgagct ggtctatagt gaagtacttg gccattccaa aggctatgga tttgtggaat      60
acatgaaaaa ggacttttgc gcaaaggcta gactggagct attgggtaga cagttgggag      120
catcagcact ctttgcacaa tggatggatg ttaatctatt ggcttcagag ctcatcatt      180
ctaagtgcct ttgtattgat aaactcccca gtgactacag ggattcagaa gagctgttgc      240
aaattttttc cagtgtccat aaacctgtgt tttgccagct tgcacaggat gaaggtagtt      300
acgttggtgg ctttgcagtg gttgaatata gcaactgcga gcagctgaag aggtccagca      360
gcagcagacg gtatgaccat caaggcagca aagtcagggt ttcttctgtg gtcctggag      420
cgccagggcg aagtacatta gcagcattga tagcggtcca acgtgtgatg cacagtaatc      480
aaaagggcct acttccagag ccaaattccag tacaattatg gaaaagttaa aacaaccctg      540
ccatgttgca agttcttcta cagccccagt tatgtggacg agctgttaaa ccaggtagg      600
accatgtatg tatactgatt aattaaagaa tgctagcatg aagttaattt tttcacatgt      660
gaaatatgga aaaatacatt gatttgtgaa aaatatattt aaattagtat aaaatattta      720
atttttctag gacctgttaa tgaattttta caataatctg aggtgatata ttttgactat      780
ttttgacata tacaacctgc ttatgtggta ttttttacta ttaataaagc taaatatcaa      840
accttctgtt tagctcttag ttgacatgaa tttatagacc aaggtaaagg ttaacacact      900
acattgttat aacctatatt aacaaaagaat taatactctc tatgtaatat ttttagcaga      960
attattttgt tgaaaagtgc caagtgtttg tttctctctt gttcttcccc tttttgttgt      1020
aaaattgttt cactttgtag caaatgatga aaacattatt attttctaag tgttatgcaa      1080
atctttataa tatcagtata cattaaatat ctacctattt agtattcttt ctctagttag      1140
agcttacctt ctgtgcattc tgaaatgtac aactttttat gtacaaaaat gtctgtttta      1200
gcattatgag gaaatgaatg cctatacagt ggtctcccct tatctatgtt cttgtgttcc      1260
acagtttagt tacctgcagt ctgaaaaatat taagtggaaa attccaaaaa taaactactt      1320
ataggtttta aaaaaaaaaa aaaaaaaa

```

<210> 56

<211> 822

<212> DNA

<213> Homo sapiens

<400> 56

```

ccgggttgac ccacgcgtcc gcggacgcgt gggcaaatat tggtaatgct gggaaaaggg      60
agttcagaat gccaaaacgt ttctgggtttt atttgtcttg ggtgaggacc cagaggggtg      120
ggagatggag gtgtgagcag catggtctgt tgtgggtttt tcttgttgtg gagtagagtt      180
agatcataca tgaagctctc tgggcatagg tggagtagca gctgtccaca ccattgctat      240
tcaaagtgtg gtttgcacac cagtaatgga aaatcatctg tgcacactgt ttagttaaac      300
tgatactttt tttttcatag caagatttct taatgaagga agtaatgtat tgatttacat      360
tctgactcat tgtctttatc ttgtctttga tcagtttgta gactggcact ggtccacact      420
ttgaataaca ctattcttca ttctactttc catgtaccgc gatgccaggc aaacaggggag      480
ttttacgctg ggtggagaac ggaacattct gctgactcct tgaaagggct tatctcacca      540
ggcatggtag ctacgactg taatcccagc tctttgggag gctgaggtgg gaggattgct      600
tgagctcagg agtttgagac cagcctgggt aacataggga taccttgtcc ctacttaaaa      660
aaaaaaaaaa aaaaattagc tgggtgtggt tgtgcacacc tgtagtcca gctattccar      720
aggctgaggc aggaggatag gttgagcatg ggargttgag gctgcartgt gccttgatgg      780
cgccactgca ctccatctg gttgacaaaa aaaaaaaaaa aa

```

<210> 57

<211> 536

<212> DNA

<213> Homo sapiens

<220>

<221> SITE
 <222> (536)
 <223> n equals a,t,g, or c

18

<400> 57
 ggacgagtgg ggagctggaa ggaggatgga gtgggaagat aatcttccct tggagttcag 60
 ctgtcccgtg accaaactcc tctctgtccc cagctggact cctctagatg ctcagatgct 120
 ccttctcttc tttccttctc tgtcacacca ttcttctgtt ccttggctct tctgctcctc 180
 tccttgtgga gscawagggt tggggtttat atgagtacag gatagggtgac atgggtggatc 240
 aaaaggcaac attttgtgtg caaaaacagg aatgcctgtt cccattaggg tcatgggttk 300
 ccagggttga ggggtggggc tttgctaggg aaccaccctc ttctaccag tattttcctg 360
 tctcctgtct gtatcaatag gtacacaata twtattaaat taatkaatga ctatacatta 420
 tgaaatggga aatgcaagggt ataaaggaga attgctgtcc ttgaaaagaa atttagtttg 480
 tttttttgtt gagatggagt cttgctctaa gctagagtgc agaattgtaat caaggn 536

<210> 58
 <211> 1262
 <212> DNA
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (12)
 <223> n equals a,t,g, or c

<400> 58
 atcctgcagt tnttctacgc aaataaaatt ctgtaagcgc agagcagcag ggaaatgggt 60
 ctgtatgagt gcattgctga gccatcatgt tcccctgttt tatctcactg gatgcctctt 120
 ctcactgctt gcctcttggg attgtaatgg gaaagagggt gctgggagag caatcaaagg 180
 caaaaataat acatggaatt gtatgatttt atctaaagta aaattctaga ctgctttcac 240
 ttatttcatt tcctcccat acaaatttga tgagcaggta cacactttct ttaaaacatt 300
 ccaagtgtat atagattcag tgcttatttc tcagcttttt ctttcttaag ttcatctctg 360
 tcacctagct tttattttta atactcaatt tctgaggctt aggaaatact tgttacctta 420
 agcgtttttt tgtttttttt ttgttgttgt tttgtttttt tttagtgtat ttgctattga 480
 tactttgcta ttgatacctt tacttgccaa gatattttta aagttggcca agataccaag 540
 aagggtggct gcaggggcat ctttggttc tcataatttt aaacagtcac tgctttcaag 600
 atcatcctaa acaagaatta atagagagaa aaaaatctat aaagcatgtg ttgaaaatcc 660
 tcaggctcat gataacttac gggataacaa tagtaggtaa caccgagtgc ttacaattca 720
 gtgggcacgg tcaaagtgtc ttttgtaagt tctatttaac acttgcgaca accctgagat 780
 agtaagtatt attattgttc tcattttata gatgagaaag ctaacacata gggaaattag 840
 gtaactttcc ccagataact cagtaaacag tggagctgga attcaaacat gtacggctctg 900
 acttcttgag cccatactca ataactcatgc catcctgcct ttccatggaa taactgatgg 960
 atggttagaa atgttaaaaa caaagcaaat gcaaacatct ctgcttctgc tcctctacta 1020
 agcccagtta acttttgatt ttatatgtac taagtataca tttttattaa tttatgtgag 1080
 ctgagtgttt ttattaatta ctttttggct attaatgtca attatttttg tgatcaatcc 1140
 tagttatcca aggagtcaca tcaactgtaa aaattacaag aaaaatctct aagtcctctc 1200
 caccctccct ctaaccagat tgcaccccaa gcttctatag aaactctggt accactcgtg 1260
 cc 1262

<210> 59
 <211> 1269
 <212> DNA
 <213> Homo sapiens

<400> 59
 gaattcggca cgagarataa cgagatgcta ctagtctggg taccagctgt acgaggaagc 60

atacagctaa	gtaggtcagt	gattaccaat	cagtgtggtg	agtacatgaa	acagcaggca	120
aggtttggga	cagccctgtt	tgaaatttgg	aagtgaaaat	catggtgcac	aaagccatcc	180
ttgctctgct	tccctgggga	ttctctgctg	atgaattact	ggcttcccta	atgatgktc	240
ttacagagaa	gtatcagaac	tgcagttcta	ccacagacat	astgaatcaa	caactcagga	300
gcttgggcca	gaactttatg	tttcaacaaa	atctccagtt	gattctgatg	tagcctaaag	360
tttaagaacc	acattgctat	agagcataaa	ttatttgagg	gtagtgctca	cggattattt	420
aaactgatat	ttctagtgtc	cagtgcctga	cctaaagtaa	gaattccaga	catgtttatg	480
aaagagtga	gagggcaca	gattttgctt	ttgaaccttt	tgagtttctg	tatgactcct	540
ttgggtcatc	taggaagatg	ggattatgaa	gctgtcaggg	tccatatwac	aacsaaaagt	600
aaatgtccag	gaaagacctg	tgactgaaaa	agtcagtcta	ttcacccttc	aagtccact	660
ccaagtcata	cctgctttgc	agatctttct	aacttaaacc	cckggtccta	tgatcctcct	720
gtctgcgtcc	attctttatt	atttaagtga	ccatataccta	actgaatatt	gaccatcatt	780
actcattttt	ataccttcaa	cagctggcac	actttttaga	atagtgttaa	tgcagaacaa	840
atctttcctc	attaataagg	atttgtggca	tacctatgtt	aatgttgaat	taaatcagtg	900
ttgaaattat	tagcaaagct	attcctatca	gacttagtcc	tgaagacaag	ccaggctctg	960
gtaattttac	accttacatt	attttcaaac	agatcagggt	acagtccarg	agccagggtt	1020
tattgggcac	cttctctctc	ttttaacatt	attgtttgtt	ctctccctgc	tgtctcttga	1080
ccagggaat	gtcttcaata	tcattttatt	tgagatcttg	ctcgaaactg	tcattggaga	1140
tgttagctgt	taatgaatgc	aaaaagttga	taagttttag	ctttcttttt	ttgatatgtt	1200
ggcttttagaa	agtcaaaatt	ctaagactaa	aagaccttca	agagagaaat	taataaaaaa	1260
aaactcgta						1269

<210> 60

<211> 1829

<212> DNA

<213> Homo sapiens

<400> 60

ggcacgagtg	gccctgatag	ataacggaag	tgaggaagga	tgacctgaaa	gtgggtcagga	60
ttatgatgac	gtcgagcctt	ggcctttcct	tcctccttaa	cttaatcctg	ggtatgaaat	120
tcacctatct	gattcctcaa	aataaatata	tacaactctt	cactaccatc	ctcagtttct	180
tctcaggagt	cctctctctc	ctagagtgc	agttgtctac	cagtagctgt	acctgcctga	240
acatccataa	atctgacaac	gaatgtaagg	aatctgagaa	ttctatcgaa	gatatttcat	300
taccagaacg	cactgcaatg	cctcgtagca	ttgtccgtgc	acacactgtg	aattccctaa	360
acaaaaaagt	ccaaacacgt	cacgtaacct	gggctctgtg	atgttggatc	tatttcttgc	420
agtatatgct	catctttatg	gaaaaagctt	tgtgggtgtg	tgctgtgtct	ccaacctagt	480
tgtctctatt	tgggaattatg	gttgggggtt	gtaaaaagat	ctggaagatg	gtttttttaa	540
aaatcctggc	ctgctgaatg	aatagtttct	cctgcaacat	tgtgggtaat	ataataaata	600
ttatcatata	aataataactt	cctgtattgt	taagtctata	catcacaaac	acttatgata	660
tataacagtt	gtatagtga	atatttcaaa	catacataag	aaatacaaaa	tgatgatcaa	720
acacctatgt	acacatcaac	ccaactttat	aaaattttta	cattatttcc	tggagaagtt	780
tttccgttaa	atttatagat	ataactgatg	ttcctctaaa	aacttctoca	atttcatttc	840
catctcttgt	ttcctatcat	aaaacacttt	taaggatgcg	ttgctgatga	ttctacattc	900
atgttgacat	tttcaactaca	aattttatata	tctataaaaa	ttatatagac	cagcttttca	960
aatgtttgaa	agttgggtcaa	tattacactg	tagagatcat	gttgatttta	ttttcataaa	1020
tgctgaaata	aatcttctaa	ttatttttaga	attcctgaat	ctatctatgt	aagtaaaact	1080
tggctcttca	tgcctttttg	tggttcatac	tttgtggggc	taagtaatat	tccatcgtaa	1140
gactgtacca	cagttgggtt	atctatttca	cctactgaag	gatattcttg	tgttccaagt	1200
tggggcagtt	atgaataaaa	ctgttataaa	caccgggtgt	gcaagggttt	ggtgtgaaca	1260
ttaagctttc	actttttgtg	gggtaaatac	ccaggaacat	ggattgccgg	ggttatatgg	1320
agaagaatgg	gttttagttg	taaaaaactt	ccaaactgtc	ttccaaagtg	gctataccat	1380
tttgcatctt	accagcaata	tatgagagtt	cctgccgtac	cacatccttg	tcagcatttg	1440
atgttgtcag	tattgtgaat	tttgaccact	ttaatagatg	tgaatgtgtc	ttattgttta	1500
aatttgcaat	tctctagtga	tttataatgt	gaaaagcatc	ttttcatatg	gttacttggc	1560
atctatgagg	tatctgttcc	attcttttgc	cctgtttgtg	atcagccttt	tcattttttt	1620
ggtgagtggt	taaatgttat	ttgtatatatt	tggataacag	tccttttatca	gatatgtctt	1680
gcaaataactt	tctcccagcc	cgtgggtttgt	attttcatte	ttgacaacgt	cttttgtagc	1740

3>

acactggttt ttaatttttaa taaagtccag cttattaatt attaaaaaaa aaaaaaaaaa 1800
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1829

<210> 61

<211> 1112

<212> DNA

<213> Homo sapiens

<400> 61

ggatagcagg ggctcagtc ttttttttgg atgaatgaac aaaccctgaa gatgctacat 60
 ctgactctat atcttcattt tatccttttc gtatttccta ttacctctaa tttctcttcc 120
 ctgcacccat ttctatttat ttcattccag tttacctcct gctgccagat taattttcct 180
 aatgcacagg ctctatcata tcatgagttt ctcatgcta catatgacta atttgccaat 240
 atttttgcac atcagaatgt gtatcacttt gaggctgggt ctgtgtttgt tttagtttag 300
 gaaaagctgt tcagattgtc tgtaaatccg tatggggatc tttgcatagg attttaaagc 360
 agccacacat cttgtacaaa atgtataaga ttaattttct atgttaggac catttgtttt 420
 caccaattcc atagagctcc aatgtgtaaa agaagacact gatctaactc ttgtgttaaa 480
 tatttagtaa ctcatttatc tggaagaaaag caaaacaaaa caaaaatata aggaataaaa 540
 atcactggga gtgcttttca ttcactgaat aatgagtttt gcaaggagca cgtggatggt 600
 gacattatat cttttacatc tttattttct gtttcttttt tgactcctta tcagtgaatt 660
 tatcttattt tatactttta ctttctattt ctttcttgac tctttgttgg tgaattggta 720
 gcaagagact tactgtctga tcagaacttt gaatcttctt gcctctcttt ctttgagggt 780
 gacagggata aagataatta agatagcgct tgggtgtgat gacactggaa gacaggctgg 840
 gtcagggcct gtagtarara cttccccctt ctattgaatg ttaatctgaa agtgaatctg 900
 aaagcagatg gtcatgaact acccagggtc tccattaagc ccatgaagtt tattttaaaa 960
 ctcttaaaat agattgagat tcaaattgag attcatgtct attttttaaa cattgtgtct 1020
 taacaaagta gatgttcagt catacagtta ggcaaatgtt ctaaggaaaag atgtttmcca 1080
 tgctaagtta aaaaaaaaaa aaaaaaactc ga 1112

<210> 62

<211> 1674

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (734)

<223> n equals a,t,g, or c

<400> 62

gagaagtcct agtggctctc tgctggtgct gaggcaggaa gggccagaag ctggtgagct 60
 tcctgaacc ctggtgtgcc tgggagtcag ttccgaagaa ggcagctgtc ccagagatgt 120
 gacaggacct ggctgttgtt tctctctgac cctaacagga tttatgccct gcatccgagg 180
 tgtttttcac tgttttatc ttattattct tattctcttg gcgtcacatg cgttttagtgg 240
 atctggaaat caaaggctga aggaagcgct gacattgac gtatctgtta atgtggatat 300
 tgccagacac aggcccttct tggagcgtat acatgttaag aagggaaca cttagcccc 360
 aaaacagtac ccagtggtg cgtttacatg cagcagaaat agctggatgg agagagattt 420
 tgagccaggg tctgccaatg ccacttcagc ctcarggctg agctgggcct ggagaggaaa 480
 gctgctgagc caccgttgga gcaaatgact gaggacctct gctgccctta cttctgccc 540
 agagctgagc tgacctgatt gtgctttggc atcaccacc actctggccc caacatctag 600
 ccttgccaag ttcaggtgct agtcatgacc atgtagagct caccgtgtac ccaaagactg 660
 tggcagcttc ctggctgctg gagctttcca gggccccctat cttctgttta ttcactcasc 720
 cagcattaac tgancatgga ctggataggg ttgctgccct tgtggaatgc accatctgga 780
 rarararagg aatgtggtat cgagacgct acagttgctg ccagtttagg gatagaagta 840
 tcagggatgt cttcccagaa gcagtgtatg tccaacaaca tctcaggcac ttggctgtgg 900
 ccacatatcg gtgcagaagg agaagcccat gcaaggcacc cacagttgag gaagcagagg 960

31

gtggcaagcc	acgagcagtg	ccctcaggca	caggattcca	gaagcatggg	caagagcctg	1020
gaggggtccac	atccccacat	tgggtctggg	gacatttgca	actccttgtc	ttgagtgtca	1080
acaacaggca	gctctttgtt	cagggcaggg	cagggatattt	ggagatgact	ggtttgccct	1140
gtccaaagct	gctgctcaca	ctgttgagag	gcttgacgcc	tggggtggga	catgggctgt	1200
gtgcatatcg	gcgaggtgt	ctggcctggc	ggctggacam	agcctcctag	tggagatgat	1260
atttggcctg	tgaactgaat	caggtgaagg	ataggagaag	aaamcascac	atttggagac	1320
ttagtggcca	gagcatgtct	tggcagagaa	actggaagga	ggctggaatg	gctgcaccag	1380
gcagaaggaa	gactcctaaa	gtggggatga	cgggtaacct	gggtcacatg	tgcgagatct	1440
tcttgtagcc	tgggggagga	gcttacactg	tcctgagagc	agtggggagc	ctcaggagcc	1500
tgcctccaga	gtatctaggc	attagttaca	tgccatgtta	cagatgagat	tgcccaggca	1560
cagtggctca	tgcctaaaaat	cccagcactt	tgggagacca	cagtgggaag	atcacttgaa	1620
gccaggagtt	ccagaccagc	ctgtctctac	aaaaaaaaaa	aaaaaaaaact	cgtg	1674

<210> 63

<211> 1045

<212> DNA

<213> Homo sapiens

<400> 63

tgcaccacg	cgtccgagat	gcacgaactg	attaattcat	ttgttctagg	gctctgagga	60
gtcgtctact	taaccttttg	ggttgctggg	cttacctatg	ttctcacgcc	tccattttct	120
caccactca	ctcagccttc	tccattttacc	ctcccaagtc	tttgccgagg	tacactcatc	180
ctgcgtatca	tactgccat	gtcctgatac	cccagctctg	ccatattgcc	cttctttttt	240
gcggtatgat	gaccacatag	aggcccaacc	tcttaaacac	atcaatacca	atgatcacat	300
ttcaatctag	acttctaagc	aacggctgaa	atctctccag	gccaaaggag	agtttgtatc	360
accttaccag	aagcttctcc	ggaacaattg	gccagaagcc	tagagtccag	aaaccagac	420
acatgcagta	agcaatttcc	agtttctcta	taatttagaa	gaggacacca	tgatatgtaa	480
tgcggggctct	ggaggttggg	atgcctccat	aaaacacctg	ccatattttt	tgggtccaagc	540
cttagtggtta	taaatcaaga	aggctgtaaa	taagacttca	gcyttttgtg	ctggtgaaat	600
ttgtttcctt	taacttatcc	tccaagagta	ccgaggcacc	gagatctacc	atttggccacc	660
tcacccattt	ctatggcaga	acaccgcctg	gggagaggaa	ttcgattccc	cgaatcagga	720
tgactgtgtg	gggcttctgc	aaagggttgca	tcacgagtc	tatttctgag	ctatctgaga	780
tccccattaa	gaatttaaaa	gcaataaaaat	aacggagatt	tttgactatc	aacatgaatg	840
ctgtgtgggc	ttttacagtt	aatgattgcc	cttgagtgtc	gaataatctg	tggcctgaaa	900
aaagaaatgt	tcttatcttc	taaatttggg	aatcaagaac	aagatagagt	aatgaatgta	960
aaggaaact	gttgcaagtt	gagtgtttcc	aaaaaaaaaa	aaaaaaaaaag	ggcgccgct	1020
ctagtaggat	accaagtctt	tacgt				1045

<210> 64

<211> 1051

<212> DNA

<213> Homo sapiens

<400> 64

ggtttttccc	gggatacatc	tgtgttgagt	cacttttgc	tcaacagtgc	ctcgccacca	60
aaatcataca	taagaggaaa	actaggactg	gaagaatatg	ctgtctttta	cccaccaa	120
ggtgttatcc	cttttcatgg	attttcaatg	tatgttgac	cactttgttt	tctataccat	180
gaaccttcca	aattgtatca	gatattccgt	gagatgtatg	tgcgtttttt	cttcagactc	240
cattccatct	cttctcatcc	ttctgggtatt	gtgtcactct	gtctgctgtt	tgaactctt	300
cttcaaactt	atcttcccca	actcttttat	catctacgag	aaattggggc	tcaaccactt	360
cgcataatcat	ttaagtggat	ggttcgagct	ttctctggat	acttagctac	agatcagctc	420
ttgctttttat	gggatagaat	cctaggatac	aactctctgg	aaattcttgc	tgtgctggca	480
gctgccgtgt	ttgctttccg	agcagtgaac	ctgatggagg	tgacatcact	ggctgcagct	540
gaaaatctag	ctgcccacag	tgaacagttc	tgcactgctc	ctctattccc	tgagctttac	600
agagtccaga	tccctgtact	gctgaactca	ggcagaaaga	agagtgcagt	ttattggact	660
ccaatctcat	tcaacagaac	aaagaagttg	aggttgcaag	gaagaaccta	taatgatggg	720

3>

tcattggaata	taacctagaa	aagaagagaa	ataaaagaga	ctgtgtttca	ccatgttgcc	780
caggctggtc	tcgaacttct	gagctcaagc	aatccaccct	cctcagcctc	cagaagtgtc	840
gggattacag	gcatgagaca	ccaagtcag	ccataagggt	cttattctat	atatacatga	900
aatgatataca	cttgaaggta	gactgtgata	agttaaatac	gtatatTTTT	taaatcttca	960
aacaaccact	aaaataaaaag	aacaaagagt	tacaactaaa	aaaaaaaaaa	aaaaaactac	1020
gtagggggggg	acggcggtacc	caattacgcc	c			1051

<210> 65

<211> 1182

<212> DNA

<213> Homo sapiens

<400> 65

ggcagcagcc	cccagcacat	ggaagccctg	ttacagtccc	tcgtgatagt	cttgcttggg	60
ttcaaactct	tcttaagtga	agagctgggc	tctgaggttt	tgaacctact	gacaaataaa	120
cagtatgagt	tgctttcaaa	gaaccttcgc	aagaccagag	agttgtttgt	tcattggctta	180
cctggatcag	ggaagactat	cttggctctt	aggatcatgg	agaagatcag	gaatgtgttt	240
cactgtgaac	cggctaacat	tctctacatc	tgtgaaaacc	agccccctga	gaagttgggtg	300
agtttcagca	agaaaaacat	ctgccagcca	gtgaccggga	aaaccttcat	gaaaaacaac	360
tttgaacaca	tccagcacat	tatcattgat	gacgctcaga	atttccgtac	tgaagatggg	420
gactgggtatg	ggaaagcaaa	gttcatactc	cagacagcaa	gggatggccc	aggagtcttc	480
tggatctttc	tggactactt	tcagacctat	cacttgagtt	gcagtgcctc	ccccctccct	540
cagaccagta	tccaagagaa	gagatcaaca	gagtgggtccg	caatgcaggt	ccaatagcta	600
attacctcaa	caagtaatgc	agaagcccg	caaaatcctc	cacctaacct	ccccctggg	660
tccctgggtga	tgctctatga	acctaaatgg	gctcaagggtg	tcccaggcaa	cttagagatt	720
attgaagact	tgaacttgga	ggagatactg	atctatgtag	cgaataaatg	ccgttttctc	780
ttgcggaatg	gttattctcc	gaaggatatt	gctgtgcttt	tcaccaaagc	aagtgaagtg	840
gaaaaatata	aagacaggct	tctaacagca	atgaggaaga	gaaaactgtc	tcagctccat	900
gaggagtctg	atctgttact	acagatcggt	gatgcgtcgg	atgttctaac	cgatcacatt	960
gtgttgga	gtgtctgtcg	atcttcaggc	ctggaaagaa	atatacggtt	tggaaatcaat	1020
ccaggagtag	ccccaccggc	tggggcctac	aatcttctgc	tctgtttggc	ttctagggca	1080
aaaagacatc	tgtatattct	gaaggcttct	gtgtgacagg	aaacccaagc	ctaagaaaca	1140
attaagtggg	tctcatctct	aaaaaaaaaa	aaaaaaaaaa	aa		1182

<210> 66

<211> 675

<212> DNA

<213> Homo sapiens

<400> 66

ggcagcagct	gctcttcttc	ttcaaatg	tcttctgggt	gatttccatg	gtgatgggtg	60
ctgtgggtgt	ctacgctcgg	ctaatgaagc	atgcagttct	ccctctgcct	caccgctgtg	120
ttcctgctgc	agctggccgc	tgggacccg	ggcttcgtct	tctcagacaa	ggctcgaggg	180
aaagtgagt	agatcatcaa	caatgccatt	gtgcactacc	gagatgactt	ggatctgcag	240
aacctcattg	attttggcca	gaaaaaggta	tgggtcagcc	agtggctctg	gggactgtgg	300
gtaaaagtga	atgtcatccc	aagagatgcc	tcacctctta	tgctgtggg	gctcttcatt	360
acctgccagg	taatggcttc	tgggaagggg	tttggcaaaa	aaagcacacg	tagcagagt	420
ctttaaatgt	acttttaaa	acacagaaca	gtatatatag	taatctactg	tggtataaat	480
ggttacttac	aggggtgag	gaactgggca	gattcttgaa	tattacctct	tcaaaagtga	540
catttttaggc	tgggtccaaag	ggagtgaagt	atctcatttg	attgttcaca	gtcagctaca	600
gatccaactc	cttgttctac	tctttccccc	cttctcagtg	ctgcacttga	ctagactaaa	660
aaaaaaaaaa	aaaaa					675

<210> 67

<211> 1105

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (797)

<223> n equals a,t,g, or c

<400> 67

gggggaaaaa	aggacacggt	gaattctgtt	gctttaaatg	tatatTTTTT	tattgtgcta	60
aaatgcacag	aacataaaat	ttgccattag	taacactgag	tacattcaca	gtgtcgtgca	120
accatcagca	ctgtctagcg	ccagaacttt	ttcatcacc	caaagggaaa	ccccgtatcc	180
atgaaggact	cactccccat	tcgccccttc	cagcccttgg	cagccaccag	aatgctttct	240
gtctccataa	attcattttt	aataagtgc	attctgtgtg	actttaaaat	aaataaacat	300
gagcacgatg	agttgcttat	tggaaggata	tccatgcggg	gaggccggcg	tgtggagtgc	360
gtargcctcc	ggacgggcag	gagttgaagg	ggcgtggatg	tgccgccctc	tcctccccct	420
gtcttttctt	tgggggtcact	gcctgagtat	ccctctttgc	aaatggcccc	aaataatgtc	480
tcagccccc	cgtctgcata	gcctcctagc	ttcaggaccc	tccaccaaaa	aacattccaa	540
gcttcagact	cactcctggg	aaaattccaa	tgccctcact	ctcccttttg	agccagccag	600
atcccatggc	ctgtggcggg	ctgcctttga	gtcctgagca	cctgtgagyt	aggggaagcag	660
gacaggcaca	cccagggaag	gggaagagtc	gtcgtcagtc	acagtaattg	atatcttttg	720
aatcgtctaa	gagatactta	gcgtgtgcct	aaaacattca	tttctttttt	tgtttgtttt	780
ttgwgacgaa	gtctcgntct	gtcgcccagg	ctggagtgc	gtggcgtgat	ctcagctcac	840
cgcaacctcc	tcctcccggg	ttcaagcgat	tctcctgcct	cagcctcctg	agtggctggg	900
actgcaggca	cacactacca	cgccctggta	gttttttgta	twtttagkgg	agacgggggt	960
tcactatgtt	ggccaggctg	gtctcaaact	cctgacctcg	tgatctgcct	acctcggcct	1020
cccaaagtgg	tgggattaca	ggcatgagcc	actgcacca	gccaaactag	cttaaaaaaa	1080
aaaaaaaaaa	aaaaagggcg	gccgc				1105

<210> 68

<211> 1279

<212> DNA

<213> Homo sapiens

<400> 68

ggcacagggt	aaacgttgta	taaaatgttt	cagtggggct	ggagacagtg	gcttcttggc	60
atcagctgca	cagggttgg	ccacagagca	ggccccgcga	agtgttatcc	cgagcagccc	120
cgccctcagct	cgagcctgg	actccaagt	ttactgtgtc	catgctgagt	ataattgatt	180
tgtttttcct	actttctcca	actttcgggt	taatcacaga	attgcttttc	agtcagaaag	240
ttcccaaagc	tctttcctgc	cctctgaagg	ctctgggagg	tggttctcat	tcacatgagc	300
ccctgggrat	gtttgtccca	gttcctccag	gctgtgaatc	aagtactcca	ttccccaagg	360
gcctgggggc	cagtaagatc	ctcaccctgg	gggctcaggc	tgaattcagg	aggaggagtc	420
actgaactgc	attcaggaaa	ataaacacat	cttgctcagc	tgagtgagcc	tttgaatttt	480
tctccctggg	aagtcacatg	tggggtggcc	araggctacc	gtgtggtcag	ccagggatct	540
tggctgggca	acgatgcctc	ccacagaccc	ttatcaagac	cacggcagtc	ttcacttttg	600
ccatctcctc	tgggtgtgtg	gaagaagcca	ttctctgggt	tgaactgagg	ggcttcagat	660
gaacaattct	gttccttcac	ccttggtctaa	tggcccaactg	gcagtcagac	ctgctagtga	720
attcaggagt	ttcccagctc	taagggtgagt	cctggcagct	ctactgggccc	tggccaactc	780
catctccagg	tcagggtttc	cgtcacaagg	caactactag	ggagaccccc	gamccccgtc	840
cagctcccac	caggctgaat	taragtcctc	stkgtgccc	ctcaccctgt	cccccatatt	900
caccattcca	gttacttttg	ctgtgaagta	ggccacccaa	actagggaga	ataaaacaac	960
aaccatttta	tgcagctcat	ggctacagt	tgggaagaat	ttaaacaggg	catggaggag	1020
atgtctcctg	tctgtctcaa	aatgtctgga	gtctcagcta	ggaagactcm	aagtctggga	1080
gtartttggc	atctgggggt	tgggatcatc	tggagactcc	tttactcctg	ggcttttctg	1140
gtccacaggc	tgggaaggck	tcagaattag	gaccgccagc	cacagtcctg	agcttggcgt	1200
ctccatgtgg	cttggcatcc	tgcagcacac	acagcagcct	ctgggctcga	ggggggggcc	1260
cggtacccaa	tcgcctgag					1279

34

<210> 69

<211> 1638

<212> DNA

<213> Homo sapiens

<400> 69

ggcagcagag	aaggtgctca	tgggtggagct	gttcatgcgg	gaggagcaag	acaagcagca	60
gctgctggaa	acctggatgg	agaaggagcg	gcagaaggac	gagccgccgt	gcaaccacca	120
caacacccaaa	gcctgcccag	acagcctcgg	cagcccagcc	cccagccatg	cctaccacgg	180
gggcgtcctg	tagctatgcc	agcggggctg	ggcaggccag	cggggcatcc	tgaccgatgg	240
gcaccctctc	ccagggcagg	cggtctcccg	ctcccaccag	ggcccgggtg	gtcctgggtt	300
ttctgcaaac	atggaggacc	actttctgat	aggacatttt	cctttcttct	ttctgttttc	360
tttcccttgt	ttttgcacaa	agccattatg	cagggaatat	tttttaattct	gtagtattca	420
agatgaatca	aatgatggc	tggtaatacg	gcaataagg	agcaaaggca	ggtgctttgc	480
agaaagaatg	cttggaact	tgagtctccc	tagaagtga	aagtgagcag	agggccctag	540
aaaccctcct	ctgaatcctc	ctaattcctt	aaaatagatg	caaaatggta	agccgaggca	600
tcgcgcaaaa	gctgggtgca	tgcttcaggg	aaaatggaaa	acccacgcaa	gaataatgat	660
tgattccggt	tccaaaagg	gtcacctacc	tgtttcagaa	aagttagact	ttccatcgcc	720
ttttccttcc	atcagttgag	tggctgagag	agaagtgcct	catccctgag	ccacacaggg	780
ggcgtgggag	catcccagtt	atccctggaa	agctagaagg	ggacagaggt	gtccctgatt	840
aagcaggaaa	cagcaccctt	ggcgtcccca	gcaggctccc	cactgtcagc	cacacacctg	900
cccccatcac	accaagccga	cctcagagtt	gttcatcttc	cttatgggac	aaaaccggtt	960
gaccagaaaa	tgggcagaga	gagatgacct	cggaagcatt	tccacagatg	gtgtcagggg	1020
tttcaagaag	tcttagggct	tccaggggtc	ccctggaagc	tttagaatat	ttatgggttt	1080
ttttttcaaa	tatcaattat	atggtagatt	gaggattttt	tttctgtagc	tcaaagggtg	1140
agggagttta	ttagttaacc	aaatatcgtt	gagaggaatt	taaaatactg	ttactaccaa	1200
agatttttat	taataaaggc	ttatattttg	gtaacacttc	tctatatattt	tactcacagg	1260
aatgtcactg	ttggacaatt	attttaaaag	tgtataaaac	caagtctcat	aatgatatg	1320
agtgatctaa	atttgcagca	atgatactaa	acaactctct	gaaatttctc	aagcaccaag	1380
agaaacatca	ttttagcaaa	ggccaggagg	aaaaatagaa	ataaatttgt	cttgaagatc	1440
tcattgatgt	gatgttacat	tccctttaat	ctgccaaactg	tgggtcaaagt	tcataggtgt	1500
cgtacatttc	cattatttgc	taaaatcatg	caatctgatg	cttctctttt	ctcttgtaca	1560
gtaagtagtt	tgaagtgggt	tttgtatata	aatactgtat	taaaaattag	gcaattacca	1620
aaaaaaaaaa	aaaaaaaaaa					1638

<210> 70

<211> 887

<212> DNA

<213> Homo sapiens

<400> 70

tttttttttt	ttttttttat	tagaataaat	gttgttgcca	aatgaaaaca	tgacactgta	60
tataactgct	gaggcctatt	ctcatgttta	accttcctaa	gccagttttc	ttaagttggg	120
ggagatggaa	gactatagta	attttcctag	catgtctggc	atctgctgct	ataaaagaga	180
cagcagtaag	catgaagact	gtgtttccca	tatttgtcca	aatcactttg	attttgcttt	240
tagagtcaag	agtcttaaaa	attgggtgatt	tttccaaatt	tttctgctaa	tagtatattt	300
aaaaatttagc	atatgcttta	ggtacatgaa	acttttaaaa	agtaattata	atgtacagtt	360
aaaaatttat	agtaagtggg	tctcacaatt	cattttctaa	ttagtaattg	atagttttac	420
ccattaaaaa	gacaatgaaa	atatttttac	tatgggtttt	cctcccattt	gtttctaatc	480
atacctttga	taatatttta	taaaggctta	taatcâtagc	agggaaattaa	tttactactt	540
aaattttatg	tatatgtaca	gtacatttaa	aaattaagta	gtcagggcat	ggtggctcac	600
gcttgtaatc	tcagcacttt	gagaggccaa	ggtgggtgga	tcatgaggtc	aggagttcaa	660
gatcagccta	gtcaacatgg	tgaaccctg	tcactactaa	aaatacaaaa	attaccggg	720
cgtgggtggg	catggctgta	atcccagctc	ctcaggaggc	tgaggcagga	gaatcacttg	780
aaccaggag	gcggagggtg	cagtgtgctg	agatcacacc	attgcactcc	agcctgggtg	840

acatagtgtgag actctgtccc cctctcaaaa aaaaaaaaaa aaaaaaa

887

<210> 71

<211> 864

<212> DNA

<213> Homo sapiens

<400> 71

ggcagcagcg	gaccggggccc	gcgggggctgc	tgcgggggcga	tcggggccggg	ccgctgccgc	60
gccatggact	cccgtgtcca	gcctgagttc	cagcctcact	gagtggccac	cccccagtg	120
ctgccagccg	aggaagcccc	cagcactgac	catgtctatt	atggaccaca	gccccaccac	180
gggcgtggtc	acagtcacg	tcatectcat	tgccatcgcg	gccctggggg	cctttgatcc	240
tgggctgctg	gtgctacctg	cggtgcagc	gcacagcca	gtcagaggac	gaggagagca	300
tcgtggggga	tggggagacc	aaggaaccct	tcctgctggt	gcagtattcg	gcccaarggac	360
cgtgcgtgga	gagaaaggcc	aagctgatka	mtcccaaacg	gscgggaart	ycacggstga	420
vccaggatgc	aaaggccyc	tggtccctgt	ttgcaagccg	gccaagargg	ggctgggagg	480
ggcaaaamcc	atacggatgc	gctgctgtct	gagaggaagg	gctgacactt	gctggcatgg	540
cctctgcggg	tttcgtccat	cgcatgcact	gatgcccggg	gacttggtcg	tcctgggctt	600
cccctcgccc	tccaggtgag	gctgcccatt	gcaggcactg	ggtaggcctg	accttgctgg	660
ggctcatggc	cctgtagcgc	ttttgttact	tgaatgtcta	gctgagcctg	tttttgatgg	720
agctactact	gtaatgcgtg	aactaacaaa	cctgtgaact	gtaaataggc	ccctggaagc	780
acgtgcttaa	gcccttttgc	tgattttttaa	aaatatcatc	tagcgcaaaa	aaaaaaaaaa	840
aaaaaaaaaa	aaaaaaaaaa	aaaa				864

<210> 72

<211> 1217

<212> DNA

<213> Homo sapiens

<400> 72

tctggaacct	ctctcttaat	tcataatttc	cgtaagtctg	tccatctgtt	gtgtggaatc	60
tcagtttgtg	atattggata	gatgcaaata	agcaaagctg	ttacttttca	tagttttcaa	120
tgaaaaactc	aacatcacta	ctgtataaat	tattttctag	tctatctgtg	tttattttta	180
aattcctttt	actattctat	acattgcaca	ttgctctggg	ggtaaaaatc	cartataaac	240
cattagctca	ttttattgac	cattcttgta	ttcagcaagt	atccaagta	cagtgggtcca	300
taccttgaat	tttttttcac	tttttaagt	agatataatt	tacataccat	aacaacttag	360
tgggtttcag	ttattttcaa	tacaagggtg	twcatatatc	atcactgtct	aattccagaa	420
catttttatt	ttattttttt	tcagcagtg	ggctctgcca	tgttgcccag	gctgggtctg	480
aactcctggg	ctcaagtgat	cctcctgcct	cagtctccca	aagtcctggg	attacagggtg	540
tgcgccacca	cacccaccct	caaaacattt	ttatttccta	aaaaagaaac	cccacatcca	600
taggcagttc	cacattccgt	tcttcctatg	atccagctct	tggcagctac	tatagttkgt	660
tttctgtttc	tgtggatttg	tctattctgg	acatagcatg	taattggagt	catacaatat	720
atggcttttt	gtgcctggct	tctttcactt	agcataatgt	ttttaagatt	cattcatggt	780
gtagcattat	cagcaacttg	tttcttttat	ggctaaataa	cactgcattg	tgtgsacata	840
ccacattttg	tttatccgtt	aatcagttga	tggatatttg	ggttggttcc	acttcggggc	900
tattatggat	gatgcttctc	tgaatatttg	tgtacaagtt	tttgtgtgga	catttggttt	960
tagttcactt	gagtatgtac	ctaggatgga	attactgggt	catgtggtaa	ctgttttaat	1020
ttcgtaggaa	ctgccaaatt	gtttcctaaa	gtggctacgg	tattttacat	tcccatcagt	1080
actgtatgac	agttccgttt	atccacatcc	actccaacac	ttgtgtttat	ctgttttgat	1140
tatataatag	ctatttttagt	gggtatgaag	tcgtatttca	aaaaaaaaag	gaattcgata	1200
tcaagcttat	cgatacc					1217

<210> 73

<211> 1717

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (712)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (721)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (903)

<223> n equals a,t,g, or c

<400> 73

gctcctaggg	gcaggacttg	gcagttgctc	aggagatggt	tgagaggtgg	gcctttccgg	60
agaagtggcc	ctttaaccct	ggactgtggg	ccatttagga	gcatcacacc	agtttttagaa	120
tgctagtagg	gacactgcga	cacagacagt	gacctgggag	caggatgcag	gagccacatg	180
gcaagtttct	gtcctggggc	aggtggctct	ggtggtgggc	tcttgccgca	ccagctctgg	240
ttcaggctgt	taacatgcct	cccgcataca	tccagataga	gaactggtac	atgatgctcc	300
tgatgggctg	ggagacaaaa	tgttgccatg	tcaggagtct	gtgggtggga	acataatgaa	360
agggaccccc	acgtggggac	agtgtggcca	ggcagcatgg	agaagtgggt	gccactggct	420
cagcctgggc	ctgagctcac	tggggaccgg	agctactttg	catgttgctg	agggatgggg	480
agggcagggc	tcttccttca	cagaatgctc	tttctcttgc	ttggtaggat	gttctttggc	540
aaaaacaagg	tgatgatggg	ggccttgggt	cggagcccat	ctgatgaata	caaagacaac	600
ctgcaccagg	taagtctctg	gccctcacgg	ggtggagcta	aggcaaagcc	gccctctctc	660
ccactactcc	catgtgacag	cctccccac	cattggcagg	ccagaagctc	anggararca	720
nggcctggct	taaactctgg	gttytccytt	macytccgra	cccaaggaat	arcasttcc	780
gcytcccayt	cyttcttgct	atgacaacca	aaaastcttt	aatgttgcc	aatgtaccc	840
ggtgagcaaa	aacgtgctta	gtagagaacc	aytgttctaa	tgtgaccaag	ctgtcctcac	900
tcntgatttg	taggtcagca	aaaggttgag	gggtgaggtg	gtctcctggt	caccaaccgc	960
acaaaggagg	aggtgaatga	gtaagtactg	ctgaggaacg	gagggaaaga	ggcgggtgag	1020
aggggatttc	agggagggaa	tgatgcagga	acttcgttcc	atatgtggca	tcaagttgga	1080
actgctcctg	gaagcctgtc	actgagctag	tggcctggga	ccacagccct	ctctgtcctc	1140
tccagcatgg	tgctctgggc	tggctgctgg	ccagcagcca	aggtctgtga	tgggggaagg	1200
cccagggcca	cgggacactt	gggggagcct	gactcatggg	ctggtggcgt	ggccaggagc	1260
atctcctcct	accaatttct	ttttccctta	ggtggttcac	gaaatacaca	gaaatggact	1320
acgcccagag	tggtataaaa	gcagctttca	ctgtgagcct	ggatccaggg	cccctggagc	1380
agttccccca	ctccatggag	ccacagctca	ggcagctggg	cctgcccacc	gccctcaaga	1440
gaggtatggg	cagccctgga	gccaaaaggt	cacagcctag	agtccaagag	cacgggcctg	1500
caagctcatc	cctttctagc	tgagcaagtt	atttctacaa	gcttccattt	tctcatgaga	1560
aaatggggct	aagagtgcac	gcctcacagt	gggggtgaaa	gcagacagta	acttacagcc	1620
aaaatgcagc	ccttgaggcc	catcaggggc	cttgcgcttg	tttaaaaaaa	aaaaaaggga	1680
atcgcatac	aagcttatcg	ataccgtcga	cctcgtga			1717

<210> 74

<211> 1276

<212> DNA

<213> Homo sapiens

<400> 74

ggcacagtga	aaacttggta	tttaaggaaa	atgttggggag	aagcccacag	ataacacact	60
tgctctgttt	aatgctaata	aattgcattt	tttctttatt	gttattattg	tcacatgcag	120
atgggatgca	tttatttata	agttctgggg	ataggatact	tttttgcctg	tactttttac	180

37

attccagagt	ttgtgcttga	cttcattcttg	actaagacag	atgagaatat	ttcctagaat	240
ttgttttttag	tcttattttgg	atcctgtcac	actgcagcat	tttattgtcc	tcagctgtct	300
gtgatcctgg	aaacatacgt	gtgactgaag	ctcccaaaca	cccaatctct	gaagaactgg	360
aaactcccat	aaaagacagc	cacctgatcc	ctacgcctca	agcccccagt	attgcctttc	420
cactcgccaa	ccccctgtg	gctccgcacc	ctagagaaaa	gattataacg	atagaggaga	480
ctcatgaaga	attaaaaaaa	cagtacatat	ttcagttatc	atctctgaat	cctcaagaac	540
gtattgacta	ttgtcatctg	attgagaaac	taggtaccag	tattttactt	aatccaaaa	600
tgtcccatat	aataaccata	tttgggaagtc	aaatgtagtc	aatattaata	tcatttgtaa	660
agtctagaag	tagaggaaaa	gaacatgggt	aacttgtcta	aggctataca	ttgagtcagc	720
attggaactt	tgattttaaa	aagctctaaa	gtcagtagat	catgggattt	aaaattcaaa	780
tggaaatctag	cagttgaggt	ttaagaatct	tctgtgctgt	ttgactattc	attcactaat	840
gaagctttaa	agacttacta	tataaataaa	accacatttt	aatgaacttg	aaagggttaat	900
aatatctaga	gaggagctcc	ctcttgTTTT	tttgtttgtt	tgtttccagt	tagaaaacagt	960
gtggaggcca	ggcacggtgg	ctcgtgcctg	taatctcagc	actttgggag	gctgaggtgg	1020
gcggatcacc	tggggtcagg	agtttaagac	taggctggcc	aacatggtga	agccctgtct	1080
ctactaaaaa	tacaaaaaaa	tttgcgagac	gtggaggcac	atgcctgtga	tcctagttac	1140
tcgggaggct	gaggycagga	gaatcgcttg	agcctgggag	gcggagggtg	caatgagcca	1200
agatagcgcc	actgtactcc	aggctgggtg	acaaagcaag	actctgtctc	aaaaaaaaaa	1260
aaaaaaaaaa	ctcgta					1276

<210> 75

<211> 1144

<212> DNA

<213> Homo sapiens

<400> 75

gcacacatac	gtatgcatat	aaggattatc	atatataaat	ttatataaca	atTTTTatgc	60
atgagtgtga	ataaatatat	gcatatatat	gtctgtatat	gtaaacataa	tgcataatag	120
aattttacata	tatctgtgtg	tatatatgtg	tgtggcacag	tcacacacac	acacacaaat	180
atgtatacag	atgcttctctg	gcttacaata	ggatttcatac	ctgataaatt	catcgtaa	240
caaaagtatt	gcaagttgaa	aatgcatttc	ataccccagt	aagttcatca	tttghtcaaa	300
agtattgtaa	gtcagaatac	atTTgacatc	tggataagtc	cattataaag	tcaaaacatt	360
ttaagtctaa	tcattgtaat	ttgggtaccg	tctatgtaga	tacgtaaatc	atacatTAag	420
ggtgactagg	tgccagggtg	aatgtttatga	aaatgaattt	caagtctcac	aggcacattc	480
acccattaca	aatatgtacc	acattcacct	attacaaata	tgtacacatg	tatgtgttca	540
tgttcatact	acaatggcag	agttgcataa	ttgtgacaga	aatcaaatgg	cttacaataa	600
ctaaggcatt	tctacatagc	cttttaaagt	aaaaagttta	ttcattgttg	gtctacataa	660
cgtggaggaa	tttgtagcgg	acaggctatt	acagtcagtg	aattgaaagg	aagggagaag	720
ttgggggaga	ctagtagctt	tttgaaggta	ttatttttaga	gattttatgaa	kttttggaga	780
acaagggatg	aggaaaaagt	attgaagaat	ttgggagagc	aggatatcaa	ttagtttctg	840
actttatttg	gaatgcagat	cagagaaagg	ctgggtaga	aaactgaaat	aataattata	900
gccttcggtg	aatatcagca	ggactgatgg	gactataggg	agggtagact	agggtataga	960
gccattgtg	gcagtttctg	taggacatca	ttggtgtata	cgtatatgtt	atTTgtgatt	1020
ttgtttatct	ttttttaata	agcaaaagga	aaagtgtcct	gatatgtttt	ggctttgtga	1080
ccccatccga	atctcacctt	gaattgtaac	aaagttttac	catgttaa	aggctagtct	1140
cgta						1144

<210> 76

<211> 918

<212> DNA

<213> Homo sapiens

<400> 76

ctgcaggaat	cggcagrgca	gttttcatca	tctatgatct	taagtttctt	catgctcttt	60
ctcattgtta	agacaatacc	acttattttta	gcttattgtt	acaatagtat	ttcatttttt	120
tctaacaact	tgggtattagt	caagatggga	tataataaca	aatgactctt	gaaatcttca	180

39

tgggtgccat	gggaagtttc	aagaacacaa	cttgaaagtg	aattgcatga	ctttgttctg	240
tgtttcattg	accactactc	attcagtgag	cctgaaagta	actgtgtata	tcactgttag	300
tatactatgc	atgccagata	ccaagactc	aaacttttcc	tttccttttag	ataccaccta	360
cttagtcatc	aattttgggt	caacctactc	cactaaatag	ctttgacttc	cattcactta	420
ccactttaga	ttagtgctat	agactcctat	tttacctcct	tcatatcaat	ccccttaaaa	480
ctcccaatag	cttccattgt	ttcaaccaa	agctcaaatt	cctttaatat	aaagtgttat	540
atgaactggc	accctatata	ctctatatcc	taatctctca	tctttcattt	atttctttaa	600
ctcctgactc	atgtaacaaa	aatgtattta	tctcagcacg	tatgtactcc	cttggtatgc	660
tctcccaa	gtccctgtat	ttttcattga	atagcaattg	ccacatttta	ttttatatgc	720
ttgtgttacc	atttatatat	atatttacta	cagactctgc	ttttagaagg	catacactgg	780
ccaggcgag	tggctcacac	ctgtaatccc	agcactttgg	gaggctgagg	ctggcgagtc	840
atgaggtcag	gagatcgaga	ccatcctggc	taacacgggt	taactccatc	tctactaaaa	900
ataaaaaaaaa	aactcgag					918

<210> 77

<211> 1065

<212> DNA

<213> Homo sapiens

<400> 77

ggcacgagag	agaggtgggg	ttaatgtgat	ggaaagttgg	cattgcttag	tttctggggc	60
tgctttctag	atagcttcta	aaccccaaag	cttctctcca	cagtgccatg	cgtctactga	120
gtactctgct	tagcttttac	cccttcagca	actgcttttt	gctcagtttc	tgtgattctc	180
accacccgt	gtggcttagg	aattcacaa	tgtttccaga	ggaagttgtg	tgaagggtgg	240
ctagaggacc	agaatctttt	tccctcgtgt	cctggctgtc	ctgtggccct	ggactccaca	300
tttttatcac	tagccactg	ggccccacc	ctgttgggtg	tcagtgtctc	ccaaactgac	360
aagtgcctg	aggaggaaac	gctcggatgc	aggcgagct	cagttcattt	cctttctatg	420
tgggacctc	accttctatg	tctttcttgg	tctccaaagc	ctcacacagc	tgttccttgc	480
tggttggcatc	acttagtctc	gcagtccctg	gtgtttccat	caggaaagtc	aggcaggcgt	540
tagcttcaact	gccaggagg	aatctcctgt	ggcctccatg	gcgctgtgtt	ttcttccagt	600
cttttttcta	tgcacagaca	catcatcccc	tttgtttcct	tttgtgatgc	tgtttctaaa	660
aggatcttta	tttctagaag	aaacttttagg	aggcaaaaac	agcgagccc	ccttaacaga	720
gtggctctgg	tggcaacttt	cccttggtta	tttgtcctgt	agccctact	ttcccaagcg	780
cttgcctgtt	gtggggctac	aggacacagc	agctgaaagg	ggctgtgggc	atcgccagca	840
tgtaccctct	tatcccattt	gctgacaagg	atcctgaagg	cccaagcata	gaaagaagtt	900
gctatggctg	ccatgtgtca	gcagcatagc	catggccaac	tcagggccct	gactcctacc	960
tgacccctt	ctgaatgaca	ctcaaggtaa	gggtcccctt	cccactcaca	ggtgaggtga	1020
aacatttcac	cttgaaaagc	ctcttgcccc	cagcctcccc	tcgta		1065

<210> 78

<211> 1126

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (1124)

<223> n equals a,t,g, or c

<400> 78

ggcaggactg	catgactggc	aagctttggc	tggttctgcc	taggttgggc	catgctgcag	60
ccgccccac	cactgcactc	tctggatcag	aactggaagg	tacctcgata	tctttgctta	120
tagctattga	cagatgaagg	gctggttacc	tgtcagggaag	tgaagagcgt	ggttaagagt	180
cctctatgct	aggttgtcac	agtcaccagc	tactagactc	ttggctacaa	catttctcac	240
caagagcagt	gtccttgggg	aaaaactaaa	cagggatgag	aaaggggtta	ggaataaaac	300
tytctcctag	agaccaggtc	agaatacata	atgggtttta	cttcgcaata	aagtgacaag	360

39

gtgcacttga	ataagccacc	ctgatacacg	gaaagcactg	ggcacagaag	taactttccc	420
attgaatcag	gagttgatcc	cataaacctt	actattagcc	aagtttacat	ttatgaacat	480
tttacacaca	ctactcagtt	atatattaaa	gacaaaaatt	gataaaatac	ttatactttg	540
gtaagccata	gagccaattc	tcttttcaac	ctagttgttc	atttcaccag	tgggcaaaaa	600
tcattatttt	taaagggtttc	caatttaaga	gcacagacca	cccagctatt	atagagctct	660
atagttttagc	cctcgaaggt	gagtcccaga	tgcagttcag	ggatgggtctg	aacctttgaa	720
cagggcaaat	ccaagcactc	taactcctgg	ttccctgctc	tatccctcat	ccatgccacc	780
ttttttaagc	atccatatgg	attagtgaat	ctcatttcca	aatctatgct	tggttagcat	840
aatctctcat	ccagaactcc	cctttgaatt	tgagatccta	tatctaaggg	cccacctcca	900
ttcctccctt	caggtctcac	agacacctta	aagtcaatgt	cttatctgtt	tccctttcac	960
gctcccaaac	gagtcattgt	tcattctcca	ctccttattt	cagtaactgg	aactccatcc	1020
ctccagaagc	acaaaacaga	acctgggagt	catccttgat	tcttgctatt	tcctcacctc	1080
ccatatccaa	cctatcccca	agtcctgatg	actttacctg	ctgnaa		1126

<210> 79

<211> 984

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (232)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (332)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (333)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (929)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (943)

<223> n equals a,t,g, or c

<400> 79

tcgacccacg	cgtcgcgcg	tccgtcggtc	ggtgcgtccg	ggcgcgcggc	ttcgmcctcg	60
ccatggcacc	ctggctgccg	ctgctgtckc	tgctggggct	gctcctgggc	rcgcgtcccg	120
ccccgccccg	ccgagcagcc	gacgctcagg	cccgggaggc	ggcgtaccgc	gagctgctgg	180
ggccccgccc	cttcgcgctg	gagatgtaca	accgcggccg	ggcagccggg	angcgggcga	240
cgctgggggc	cgtgcgcggg	cgctccgcc	gggcgggcga	ggggtcgctg	tactccctga	300
gggcgaccct	ggaggagcct	ccctgcaacg	annccacggg	gtgccagctc	cctgtgtcya	360
agagaccytg	ctctgcagct	ttgaagtctt	ggacgagcta	ggaaaacaca	tgctactgag	420
gcgggactgc	ggcccagtg	ataccaaggt	cacagatgac	aaaaacgaga	cattaggttc	480
agtccttcca	ctgttgaaca	aggaaccctt	gcccaggac	ttttctgtga	aaatggcctc	540
aatcttcaag	gagttcggtta	ccacctataa	tcggacgtat	gagtcgaagg	aggaaaccca	600
gtgggcgatg	tctgtctttt	ccaacaacat	gatgcgagca	cagaagatcc	aggcactgga	660
ccgtggcaca	gctcagtatg	gggtcaccaa	gttcagtgac	cttacagagg	aggarttcca	720

40

taccatctac	ctgaatcccc	tcctaagaga	gtaccatggc	aagaacatgc	gcctagacaa	780
gtctgtctggc	gactctgccc	catccgartg	ggactggarg	araaaggggg	scgtcaccaa	840
agtcaagaac	caggcatgtk	tggctcctgc	tgggctttct	cartcactgg	taacgtggag	900
ggccagtgg	tcctgaaaca	ggggcctgnt	ctscctctcc	gancargarc	tcttgactg	960
tgacaaggtg	gacaaggctg	cctg				984

<210> 80

<211> 1247

<212> DNA

<213> Homo sapiens

<400> 80

gcgagttcat	ctaactatgt	gtccacag	ggatctctcc	ttgagacttg	acagttgatt	60
tgcagacaga	agtatgccct	catggaactc	tcacctccag	gggttcccc	tagacctcac	120
atccattagg	agggggtgtg	gaaaggatgc	ccacgtggcc	acttttacia	ctgctgtcct	180
gtcatttcc	ttccctactt	tgtgaaacgt	tcactttctg	ctccaaagat	gaagtgtcac	240
gttggaaggc	gggatgcttt	gtgccccttc	cagcaagcta	acttccaaat	aaattctcta	300
wttttatata	agaccttggt	cttggttaatt	agactttaca	tgaagtgage	aactaagctt	360
ttctgttaca	agacttcatg	cccacagata	cattcaagtc	ccaggtggaa	ggatgatctg	420
gattaggcaa	ggtgctgacc	atgggaacag	gagacagtca	agggaaatctg	aggaagcaca	480
tgtttgtgtc	caatatagtc	ccctccttgt	gccactcaga	tgtgctcatg	ccctccaact	540
gtggctgggt	ttataagcag	atgccttgtg	tagttgcacg	gggtcttgtg	cttgaggggc	600
tttttgcttg	gattaatggt	ctgcacttgc	atttttatct	ttcagacagc	ggatactcct	660
ccactgaaga	gggaatagtt	ctgcagtaat	tccttagggg	tttgctttct	cctctcttac	720
gggcttgatg	gagacaggca	cagagtccctg	tgatgccac	gtkgeatgcc	tctagcagct	780
ttgaattctg	ctggatcatc	tggcacaatg	ggccgagcag	gttggggccag	accctgtatc	840
tgctgtagag	atattatatg	ttctggattc	cacttgaaac	caggagtctt	tgcattcatt	900
ctgtaaccca	tctgggtttg	gcagaggcta	ggtagggttaa	ctgaacaggg	atttaaaaag	960
gactttatta	acacatcaga	gcaatatact	ctactgtggg	atttttggac	tccaaaatcc	1020
ctagaggctc	aataaacgcc	gtgcttctct	tttagtgata	ggaagtatat	aggaagccat	1080
ttaccttaaa	caggatgttt	ctgctcggca	tgggtggttca	cgctgtaat	accagcactt	1140
tgggagacca	acgcaggcag	atcactggag	gccaggagtt	tgagactagc	ctgagcaaca	1200
tggtgaaacc	ccatctctac	ttaaaaaaaa	aaaaaaaaaa	ctcgtag		1247

<210> 81

<211> 946

<212> DNA

<213> Homo sapiens

<400> 81

ggcacgagtg	agattgcac	cagacagagt	tttaaaagtt	tcccgggtga	gtttaatgta	60
cagttgaagt	tgagacatga	atctctgcat	gtaggggaaa	ttttgtgtct	ggtagtcaa	120
gaaactatgg	aaaccaattc	ttgatatttt	gaaccattca	cgaagatagt	ttgagtcag	180
agcatgctgt	tgtctagagt	gggcggggat	gactcattgg	agtggatgcy	ctgctctgta	240
cttgattttt	ttgagtctga	aattagcttt	ccaggctggg	gcaggaggag	gagcacagg	300
gggatcagta	ctgcccccaa	gcggtggagc	tgtgggtggg	gatcaatact	gctgccgcct	360
gtctgcacaa	acataattct	ctcttccagc	ccttcagaag	tgtattggaa	tatgtcgata	420
acaataatga	tggtagtgaa	gatgatgatg	atgtgggtaa	ttctggctac	cttattgggt	480
ccaagctccc	cacaattcgt	tgcacaaagc	actctacata	cattctcttt	agtcctgatc	540
aaaccacctt	tcagagtagg	atcttagtgtc	ctattttaaa	gatgaaggag	ctcgggctca	600
gagagagatc	gttttagacac	acacacaact	ttggaatgaa	acatttacag	ccgggcgcgg	660
tggcgcgtgc	ctgtagtccc	agctacttgg	gaggctgagg	ctggaggatc	gcttgagtcc	720
aggagtctctg	ggctgtagt	cgctatgccg	atcgggtgtc	cgactaagt	ttggcatcaa	780
tatggtgacc	tcccgggagt	ggaggaccac	cagggtgcct	aaggaggggt	gaaccggtcc	840
aggctcgaat	gaaacattta	caaaaattga	catttcctta	tgcatagata	tttactagg	900
tccttaaaac	ccacgtgaat	ctgtgattaa	aaaaaaaaaa	aaaaaa		946

41

<210> 82
 <211> 1392
 <212> DNA
 <213> Homo sapiens

<400> 82
 attcggcaga gcagaaaacc agactgcact tgctttataa aacagagcct tatttttcct 60
 tcataataag cagagttgca gtgttgctgg tattgattca ctggcgtggg ggtatcagga 120
 cagatgtctc tatgattaat ttttggcctg tcaactcatgt ttgcataatg ctgttggtggc 180
 tccaagcatt ggaagcaaga ggacagggaa gcaacattga ctgtaccagg aactccaaaa 240
 cagtcttcac atcttaattg ttggacaatg ccaaattggc actcttttct ggaagttgac 300
 tggggacaag atagtggtaa ggattagatt tggccagaaa gtttctgcca cagtgaagctt 360
 tcctgtctaa atccttattt taactgttgt cacttaatat tcacactttg gaaggacatc 420
 tactgttggg tacaattatg aaaccaactt gaatactttt tagttgaaca tttcagtagt 480
 cttaattatg tttaaatagg tttcacaatt tactgttttt agtttagttt cgggtcccc 540
 ccaaccccc aacttttgyta gagagttact ctcttaactt ttgctagaaa gtagcaaaagt 600
 tctctactct acatgttcag ggctggctgt agaatttcgt tttttaagga aacaggaaga 660
 cagaactaat tatgcaagtc ttcatttagc tttttaaaaa aacagcttta ttgagttaga 720
 attgacatgc agtaaattgg acatatttaa agcgtacaat ttgttaagtt ttgacataag 780
 tatacatgtg gaaaacatca gtcaccacaa tcaggatact tattttaaaa aacaacttta 840
 tttaggatta gtatactgat aatgtgtcca ttgtaagtgt acattttcag ttttgacaaa 900
 tgtatagatt tttgttaacta ccaccaccag tcaagatgaa aacgtttcta gcactccaga 960
 aagttccctt gtgtcccttc ttggtcagtt attcccacca tgctctcagg caaccacagt 1020
 tctgcttcta tcactatata agtgacagaa tttttctaca gaatttcaca tagatggaat 1080
 catacaatat gtactgttct gtctggcttc ttgaggtaaag ccaaattgtc ttttaagagtc 1140
 atgcatgttt ttgcatttat tagtagttta ttcttttttt gttggtgagt agcattcatt 1200
 gtatggatat attccagtcg gttttattca ttcacttttt ggacatttgg gttgttatca 1260
 attttgggct cttttgaatt aatccctccc tccttccctc cttcccyccc tccctccttc 1320
 cctccctccc tccctctctc cctccctcct tccttccctc cctccctccc tccctttttt 1380
 ttttcggcac ga 1392

<210> 83
 <211> 1155
 <212> DNA
 <213> Homo sapiens

<400> 83
 ccgggtcgac ccacgcgtcc ggtgagtgca ctctaggatg ttcacatgat gacaaaatca 60
 cctaacaatg tagacgttc agaacatata ccctttgtta atcgatgat cactgtatat 120
 atgtgtgtat atacacacat atatgtacat atatttaata catttgtgta tgtgtgtgta 180
 tatatatata tatatacttc tcattattta tactctagac ccagagcctc ctagtgggc 240
 tccaaaattg gactctcatc tctctttgag acagccttca aatgatcgtt tttaaagtgc 300
 taattaactc ctcttctcaa aatgcttcaa tggccacta atctctaccg aatcaaggaa 360
 tcagccata ctgtcccaag atatctttcc ttggccagtt ggagcctcat ttcagctgct 420
 ctgggtttat cccctgtctc ttctttccca cttccaagcc tgtgctcagc ccacctctc 480
 ttctggggat gccccacacc ccaactctgc atatctgcca aacctttcat ctccccgtga 540
 agctcttgac accaaataca gtttacttta gaaaatgtat tttttccact ttctcaacta 600
 aacttttcc tgtgtgatct gcttttccgc tgccaaggca catcggtttt aattctctac 660
 agcactgctc atatcttgcc cagtattata gcttctacat attggtcttg cttcttattt 720
 ttgagcacia aaactaagcc actccacttt ctcttaccag tgaatccagc ttaaaaaaac 780
 tgtgagcaac ctatcagtat tttgttgaca tgaactctat agaaacctta gtccctggat 840
 cttcgactct gcctccctg acatttatct gctccacaa agcacgcagg tgtgggaaga 900
 gaagtggctg ttttttgagg tcacatttca gccctgatc atcctaagt cttcaccctt 960
 tttatccttt ggscactgtg tycctagaga tgtgaattca attccgcacc attctctcct 1020
 ttacaatgat gccaatattc tcaggctttt aagactaaat ttttaattac gagaaaattt 1080

12

gatcttcaaa cttaagttgg acctagaaag aacaatctca tgaactcaaa aaaaaaaaaa 1140
 aaaaaaaaaa aaaaa 1155

<210> 84
 <211> 1373
 <212> DNA
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (877)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (897)
 <223> n equals a,t,g, or c

<400> 84
 gatgtgctgc tgcctctgct gtacctcctg gtccggaagc acatcaacag agcgggtatc 60
 gggaacacgt ttcagggagg tgccaactgc atcatgttcg tcctctgcac ccgcgccgtc 120
 cgaactcggc tcttctctct ctgttgctgc tgctgctctt ctcagcctcc caccaagagc 180
 ccggttgga cteccaaggc tcccgcgcct tccaagccag gagaatctca ggaatcccaa 240
 gggacccag gggaacttcc aagcacctgg agcttttgct cttctagtt ctgtggcata 300
 ggtgctgct tcctgggggt aggcamtct gtgagtcct gstgcagggc agaagtgcatt 360
 gtctgctgca ccagaagccc ccgctattga tgctctcggc tccattggga gagcagctgc 420
 caactcagct tcttctacct cctagatcct caggcagcaa gttccamcgg taccagcgtc 480
 ctggccccag ggtgggcgtt catctgcata agggtagcag tgagattcgg gaagccgggtg 540
 gccacacag gtggccacag tgccccacc cagtggacct tgatgtcctg aggaccacac 600
 agcactgttt acaatctgag ggaccacat ctgtgcacct atcctctgtg tgattaaata 660
 agctgtcagg ggcctcaga gatgtgtccc catccacaca tcatggagca gattcccggc 720
 tggtcagggt gttaagattc tctggagaac ccggtttcc ttggagactt tgctcaacag 780
 aggaaccakt tgccattcac caggttgaag gtyccctgar gttgcatggg gaccaccagg 840
 ggcaagcaca gcccggcgca ctgctcaggg acagcantct ccccggsct cytggcntct 900
 ccytccatgg gctgctaacc cttgccgtag cttgggtttg agtgtcagca caggccagaa 960
 tccacactcc tgtcccaact gtcacactt cagacacagg tccctcctt ccaattccaa 1020
 agctggaaca agggcaggaa gaggcgtctc agggaaacccc cgaagacccc actgtctctc 1080
 cagctccttt cttagaatcc tcctcttctt ggggagacca aaggggatga gtagggaggg 1140
 gttctggctt atttacttct tagtttagtt ttttgtaagt aaagtggccg gagagatgag 1200
 taatccaaat acgattcctc aggcggggca cagtggctca cacctgtaac cctagcactt 1260
 tgggaggcca aggcaggcag atcacctgag gtcaggagtt tgagaccagc ctggccaaca 1320
 tggtgaaacc ctgtttctac taaatataca aaaaaaaaaa aaaaaaactc gta 1373

<210> 85
 <211> 1258
 <212> DNA
 <213> Homo sapiens

<400> 85
 tgcactgtgg gtgtccaccc acaccttgtg ttcctcatgg cctaccccag cctttcttct 60
 ccactgggtc ccactgttcc ctggagacag agggctagca tgctgtcatt tatctgaagg 120
 ttgtggctga cccattctcc tgggatttcc caggccacct ctcctttccc tttccctcac 180
 ttaaccaga cttgctcagc tgaggctatt gtccctgatg ttggctttac ttgtaggagg 240
 tttagtggct gctctggcct gccatggaat tttggctgca attttggcag tgtgtggaga 300
 actggtatca ggaaagggaa ccaggagtag tgatgaagat gatggtgggg atggggacag 360
 aggacatagg ggactgtccc tcttgaactc tgctttggg cacatgggag atggggacag 420

43

gaaagatgat	aacagtggaa	ccctgtgaag	aataggagaa	tgggaattct	aaaataccag	480
ttcccaaaca	aaccatctct	ccattggaaa	aggagctggt	gcaaaccgtg	ttcattttaag	540
cttctgatga	aaagcaatgt	atgtttataa	gcttgcaagt	aacagacaag	atgttccaag	600
gccgatgagg	aaaatatcgc	ttccctgtct	ctgcctctac	cttgctgtgc	ctctctggct	660
ttagcagcat	tggatgttga	attggtttgg	ttcattttctc	tgaagtcgga	ctgtcaggag	720
gagccagagg	ttggctttct	gggatgtgaa	ggggacagtg	tcagcctaag	cctctggaga	780
tgcttaggat	gctgaatttt	aaccctttct	ttcactgcat	ggcccttcaa	acagggattg	840
gcaggctccg	cgagttacag	agtgggtgag	tcttgaggag	aaccaacgac	taaggtagct	900
tagctctctg	ctttgttctt	gggagccaag	agaatgcaag	atcctggtgg	gtgagcagtg	960
gtatgaagag	gggacactgg	tgggtgaaca	gtggcatgga	agtcactgat	gcagcctctg	1020
gccctcagtc	tccctatcgg	caaagtgggg	ggctcttctc	atctctaaag	agtttcatcc	1080
ctgacaggtg	gatgggtgaa	agatccagag	agtgggggtg	aggggtgggc	cctgaagact	1140
ttttgtctgt	agctcttgca	tattgagtgt	ataaacccgg	cttctggacc	aaccaagat	1200
gaataaactg	gggcagaaaa	ttaaaaaaaaa	aaaaaaaaaaa	aaaaaaaaaaa	ctcgtagg	1258

<210> 86

<211> 1318

<212> DNA

<213> Homo sapiens

<400> 86

cccacgcgtc	cgaacagctc	ttaattgggt	ctttcagttt	ttaatggact	tcctgatgtt	60
aagagagtga	gagagcagtg	gtttacttaa	tcaccggggc	ccttttcata	gtttcgtctt	120
gtgtgctgtg	ctttttaccc	agctctagga	gggagatgtt	tgtgggtacc	aggggtttgt	180
tagtccctct	cccccttttt	agtataagtg	gcatgctggc	aatagacaaa	taccttcata	240
aaaagctgct	tctaaatgaa	ataattacta	cctccacttg	ggcattatga	taattatcaa	300
tagaatattg	aatcagcttg	gtttttcagg	aaaaaaaaat	ccaaacaaac	cttccttaaa	360
caaggaaggt	tttatttcag	gtctccaatt	agagtaaaca	agcatttggg	gcttttcctc	420
ttgttttctg	tgttaggatt	gtgaatacta	gtctgctata	gaaggttgct	agttactttc	480
tgtaatgtag	gcagtttctg	gggattggag	acacagatgg	gagtaaaggg	tctggattct	540
tctctttccc	acctgcctgt	ttggctttcc	ctcagcctcc	cacctgtgc	ccatgtgccc	600
cttctctgat	ctctgtgctc	agatggcagg	gaccatagga	gttcatgtgt	ccagaggtaa	660
cactacccga	gtctgtgata	ctcagaagct	tccagttcct	tattaaaacg	ggaaattgaa	720
cagtaaattg	ttactgccta	ccagatttca	cagcaggaga	ctgtcattac	acctttcatt	780
tgactcagag	agagtgtagc	catcagatgc	ccagattttg	ccattcttta	gtcacttgga	840
aaatgcaccc	attgtcagca	ttttgagtta	ttaggattcc	tcaaatgaaa	aaaaggcaaa	900
caatatattt	tgatggattg	aaatccacac	tgttgtgctg	aaaatagggg	aagggaaaaa	960
ggaagccaag	attttggcag	aaaactgaac	actcttaaga	ataactcctc	agccggggcg	1020
ggtggctcgc	gcctgtaatc	ccagcacttt	gggaggccga	agtgggcgga	tcacgaggtc	1080
aggagatcga	gaccatcctg	gctaattgtg	tgaacccccg	tctctactaa	aaatacaaaa	1140
aattagccgg	gcgtgggtgg	gggcgcctgt	agtcccagct	actcgggagg	ctgaggcagg	1200
agaatggcgt	gaacccggga	ggcggagctt	gcagtgagct	gagatagcgc	cactgcactc	1260
cagcctgggt	gacagagcga	gactccgtct	caaaaaaaaa	aaaaaaaaaaa	aaaaaaaaa	1318

<210> 87

<211> 978

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (977)

<223> n equals a,t,g, or c

<400> 87

ggcacgaggc	gcgccaaggc	gtcagtcgag	gagtcaaggc	agcaatgaat	cgtgtcttgt	60
------------	------------	------------	------------	------------	------------	----

44

gtgccccggc	ggccggggcc	gtccggggcg	tgaggctcat	aggctgggct	tcccgaagcc	120
ttcatccgtt	gcccgggttc	cgggatcggg	cccaccctgc	cgccgaggaa	gaggacgacc	180
ctgaccgccc	cattgagttt	tccctccagca	aagccaaccc	tcaccgctgg	tcggtggggc	240
ataccatggg	aaagggacat	cagcgggcct	ggtggaaggt	gctgcccctc	agctgcttcc	300
tcgtggcgct	gatcatctgg	tgctamctga	gggaggagag	cgaggcggac	cagtgggtga	360
gacaggtgtg	gggagagggt	ccagagccca	gtgatcgttc	tgaggagcct	gagactccag	420
ctgcctacag	agcgagaact	tgacgggggt	cccgcctggg	ctggcaggaa	gggagccgac	480
agccgcccct	cggatttgat	gtcacgtttg	cccgtractg	tcctggctat	gcgtgcgtcc	540
tcagcactga	aggacttggt	tggtggatgg	ggcacttggt	tatgctgatt	cgcgtgaagg	600
cggagcagaa	tctcagcaga	tcggaaactg	ctcctgcctt	ggctcttgat	gtccaaggat	660
tccactggca	agacttctca	gatccttggg	gaagggttca	gttgcactgt	atgctgttgg	720
atttgccaag	tctttgtata	acataatcat	gtttccaaag	cacttctggt	gacacttgtc	780
atccagtgtt	agtttgagg	taatttgctt	tctgagatag	aatatctggc	agaagtgtga	840
aactgtattg	catgctgcgg	cctgtgcaag	gaacacttcc	acatgtgagt	tttacacaac	900
aacaaatgaa	aataaatttt	aattttataa	tatgggaaaa	aaaaaaaaaa	agggcggccg	960
gtaaccatt	gcgccna					978

<210> 88

<211> 1863

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (82)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (112)

<223> n equals a,t,g, or c

<400> 88

tgggggttgg	gatgcagagg	catgaccccc	tttcttcccc	tgcccccttc	tgacatgaga	60
tgctctatga	tagaggtagc	tnngggcggtg	agtgaacacag	aatgtgggtt	tncggggagg	120
aagtgttggg	gtcacatgct	gcttctcctg	ctttcttaca	taggtgcttc	agcgaagagt	180
cctgcgtctc	catccctgaa	gtggagggtc	acgtggctgt	ccttcagcct	gacgcccccc	240
agatcctgct	gagtggaact	gctcattttg	cccggccagc	tgtggacttt	gagggaaacca	300
acggcgctcc	tttggtccct	gatcttcaaa	tcacctgctc	catttctcac	caggtggagg	360
ccaaaaagga	tgagagttgg	cagggcacag	tgacagacac	acgcatgtcg	gatgagattg	420
tgacacaacct	ggatggctgt	gaaattttctc	tggtggggga	tgacctggat	cccgagcggg	480
aaagcctgct	cctggacaca	acctctctgc	agcagcgggg	gctggagctc	accaacacat	540
ctgcctacct	cactattgct	gggggtggaga	gcatcactgt	gtatgaagag	atcctgaggg	600
aggctcggtt	tcggtgcga	cacggagctg	ccctctacac	caggaagttc	cggctttcct	660
gctcggaaat	gaatggccgt	tactccagca	atgaattcat	cgtggagggtc	aatgtcctgc	720
acagcatgaa	cgggttgcc	cacccagcc	acgtgctcag	ctyccagcag	ttcctgcacc	780
gtggtcacca	gccccgcct	gagatggctg	gacacagcct	agccagctcc	cacagaaact	840
ccagtacgta	agcctgggtg	ggctgggcag	ggaggggcag	gtggcagggtg	agtgtgttgg	900
gacaggtatc	ctccccctcc	acctctggga	gaggacaagg	taggtggagc	aatgggttct	960
gttcctgtg	gtacctgct	tagttgacag	ctatggacca	atccctctct	ccctttggat	1020
cattcacttt	cccttgtgaa	gaacatgagc	cttgtgtcat	tgtagggatg	cttatgcctg	1080
cartcataaa	gtcagtgtgg	gtttaggcgt	gtttgttata	cctgatgatg	cctccttccc	1140
cagccccaag	caatatcaaa	ttagccctgg	acctaggatc	ytctgtgggtg	aaactatcaa	1200
gtatttacta	agcacytact	atgtactagc	actgtcctaa	gatgatctaa	atgatttata	1260
ggatagatca	gtccctatcc	ccaaggagag	aattgggtact	attacatgta	aaattatgtg	1320
ctacgtatgc	agtcacaaga	atattttctaa	aaaatatatg	caaatatgca	aagcgttaca	1380
ttcagatgga	gtgctgccct	tcaaagctgc	ccttcaaagc	aacctggccg	tggtcccaaa	1440

45

ggttccccggt	catgggcctt	tacgaatggg	gcaattttat	tttcaacttt	tgagttgcgt	1500
aattgtagaa	aaaccagaaa	atatcaaata	tttcttttaa	agaagaaaaa	tcaggccagg	1560
cacagtggct	catgcctgta	atcccaacac	actaggctga	gtcaggaggc	tcacttgagt	1620
ccaggagttt	gagaccagcc	tgggcaacac	agtgaagactc	tgtctttaca	aaaaataaaa	1680
aaaattagct	gggcatgggtg	gcagggtgct	gtagtcccag	ctacttggga	ggctgaggta	1740
ggaggatctc	ttgggcctgt	gaggccaagg	ctgcagttag	ccatgatcac	accactgcac	1800
tccagtctgg	gagacagagc	aaggctctgt	ctcaaaggaa	aaaaaaaaaa	aaaaaaactc	1860
gag						1863

<210> 89

<211> 2086

<212> DNA

<213> Homo sapiens

<400> 89

cgaggaatgg	agccggtagc	tgctgcggcg	agtsccgcgg	ctcctccgta	gacccgcgga	60
gcaccttcgt	gttgagtaac	ctggcggagg	tggtggagcg	tgtgctcacc	ttcctgccc	120
ccaaggcggt	gctgcgggtg	gcctgaatgt	tcgcacttta	ccacatacag	ttctttacat	180
ggctgattca	gaaactttca	ttagtctgga	agagtgtcgt	ggccataaga	gagcaaggaa	240
aagaactagt	atggaaacag	cacttgccct	tgagaagcta	ttccccaac	aatgccaagt	300
ccttgggatt	gtgaccccag	gaattgtagt	gaytccaatg	ggatcaggta	gcaatcgacc	360
tcaggaaata	gaaattggag	aatctggttt	tgctttatta	ttccctcaa	ttgaagggaat	420
aaaaatacaa	ccctttcatt	ttattaagga	tccaaagaat	ttaacattag	aaagacatca	480
actcactgaa	gtaggtcttt	tagataaccc	tgaacttcgt	gtggtccttg	tctttggtta	540
taattgctgt	aagggtgggag	ccagtaatta	tctgcagcaa	gtagtcagca	ctttcagtga	600
tatgaatatc	atcttggctg	gaggccagggt	ggacaacctg	tcactcactga	cttctgaaaa	660
gaaccctctg	gatattgatg	cctcgggtgt	ggttggactg	tcatttagtg	gacaccgaat	720
ccagagtgcc	actgtgctcc	tcaacgagga	cgtcagtgat	gagaagactg	ctgaggctgc	780
gatgcagcgc	ctcaaagcgg	ccaacattcc	agagcataac	accattgggt	tcattgtttgc	840
atgcgttggc	aggggctttc	agtattacag	agccaagggg	aatgttgagg	ctgatgcatt	900
tagaaaagttt	tttcttagtg	ttcccttatt	cggcttcttt	ggaaatggag	aaattggatg	960
tgatcggata	gtcactggga	actttatatt	gaggaaatgt	aatgaggtaa	aagatgatga	1020
tctgtttcat	agctatacaa	caataatggc	actcatacat	ctgggggtcat	ctaaataata	1080
attaaagtgg	ctttcataat	atgtaacttt	tgggttctgc	ctttttcaga	aaatggaaac	1140
ttggggccatg	tgtattttcaa	acaaaaataa	ctttagatat	atcttttttg	tagcyttgat	1200
tgatgctcta	agatcacatg	agggtagtat	ttaatatatt	agatgaagga	caactttgga	1260
cataacactg	actaggagtt	gagagctttt	gcactcaggca	gaagcaaact	gattatagtt	1320
gtgttgccacc	agatcatgta	gctgctgtgt	aacatgacct	taaatagtct	tctgcatag	1380
gaagagcaaaa	aggggtattca	tcaataggat	atagatttaa	gacattccct	gactaccctt	1440
tgcgttggtta	ggatgatgtct	tttagcagaa	tcatagaagac	cttttttctc	ccttaataaa	1500
ggagaaaaaat	atactgatgg	ctggagaaat	ttttctctgc	ctttcagttt	tatgaatttt	1560
ttcagaagta	acaatattat	tattgacttt	ttacttattt	gataaaaatt	aaagaactat	1620
ttttgttttg	gtcagataaa	ttgacaagac	taatcagtat	tttattataa	gtaaaagatt	1680
tttcttcttt	ccttaaaaaat	attttttttt	cacctaggtc	taaatagcta	actaactggt	1740
agaccagagt	attacatcat	cttatttttg	ttttatacca	ataaaacata	gcgtgggaact	1800
cattcaggta	atgtttttgca	tttcattgct	tttggatgaa	caaaggaagt	aaactaatcc	1860
tttataaatg	aaaaccagaa	atagttggta	tgtcagctag	tcattcctgt	catattccca	1920
gtagaatgat	tttcaagttt	gaatttctgt	wcaaatatct	aaataagaga	tgtgcagaga	1980
gcaccaattt	tccttcaata	tccattcttt	acttttcaca	taatgataga	acctttgatt	2040
tttcaagtgg	gtatgcctcc	tagaataaag	actacatttc	ccgagt		2086

<210> 90

<211> 891

<212> DNA

<213> Homo sapiens

<400> 90

gaaattatgc atgagctgtg attgcaagta atttttaaac gtctgcatgt ggacaaaagta	60
aatacataaa aatgaaaatg gctgtgtaag agagtaggat gaacaataat aattgttttc	120
cccactgttg tgaattatga tgttatcttt taacttcaca tggatggttt gggtttctct	180
agttcttaaa agtcaaagag ccaagttagc cctccattcc ctccatcttc atcaagaggt	240
caggctcaga atgtcacgga gagagtcgcc tggcaggcct cttcgttggt gggtgagagg	300
gaacatgggt gccagaacct cagtgccaac cgctgactat ccttctccct acaggacatt	360
gccgagaatg gctgcgcccc caccacagaa gagcagctgc camagactgc accgtcccca	420
ctgggtggagg ccaaggacct caagctccga gaagaccggc ggccaatcac agtccacttt	480
ggacaggtgc gccacactcg tccacatgtt gttaagagac caaagagcaa catcgcagt	540
gaaggccgga ggacgtctgt gccgagccct gagcaaaaca ccattgcaac accagctaca	600
ctccacatcc tacagaaaag cattacccat tttgcggcca agttcccgac gagaggctgg	660
acctcttcac cacattgact tacgccgttg cttttccaga ctgggcagag gggctgactt	720
cgcagtgtgt gccaaagarc cgggtgtctga taatcccatt ttctgtctta tcacctgaac	780
tgtgtcagta tcacttttag tttgttgggt tggttggttt gttgtttgtt taatatgccc	840
tgttttctac ttctgttggg aaatatttgg ggttgaaata aaccagtggg a	891

<210> 91

<211> 1974

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (654)

<223> n equals a,t,g, or c

<400> 91

aaaattgcta attaatatta gtatttgggt gctttatctt aaaacaatac ctacctctgt	60
taataggaaa attagaaatt atatctattg ttcttaaaga tacaatttgt ttctctaact	120
ccaagggaag tgaatgaata tgtttcagac tatacttgct tgtgtattat ttgtttttgt	180
aagggtggtt ttctttttgc ttcaaattga gagcatacaa acaaaattcc actgtataag	240
cagtcaattt tggtataaat aacacatgcc tagagcactc agattaaatt attacacata	300
ctaataaaat tattcacagt agtactcctt ggtaattgtg gtattaggga acagggatag	360
aacctatctt caaagtcaac cctattactt caagcctttg caggtcctgc tgtttccaag	420
gatacaatcc aatgcttaat ttacaggaat ccgaaatcat ttgctgtgtt aaaaacttga	480
gattagccct tgctgcccc ttccagggtg gcctttggaa gcgaaaggtc tttaagctta	540
agtaaaactg agaataattc tgagtaagca ggctccagggt ggagggtaca ctccaagata	600
ccttcacttg ccaaaaacaac actccctgca aggttatcag cgaaaaggca gtanaaagag	660
ccctcctgct gccagaagggt cttgttttgc aaggcttctc agcctgcctg cagccattgg	720
gtgaggagag gaaaggacac ctcacctggc agagcagcac aaaggtcagg catgtttccc	780
tcctgacctg gaagctgggt cctgctcctt cacatgaact gtccagggtt tggatcata	840
cccagggtccc ctgactcctg ccagartga ctccactgggt ctccccagaa aacaccaaca	900
gcagararaa atgaartgtg tctgttttct ggtttcataat tgcactcctt cttgtctctc	960
ytaggacaga aataacatct gcmatgcar aramctgtga agtaataaac aacctttccc	1020
ccagctcttg acaagcatta ccttgcttkg aggtgcaatc agttypgaca agtttatggy	1080
tttgtgtcyt tcaacccaat ggacattacc ytcctggggga gtttaatgac ccytytcagc	1140
ttcacactca cttgaaggaa agagctagaa gtcagtaatg atgtgttatg acttgcatgg	1200
cttcaggtag ttcagaacct gcaacctgta gttcagatta caggagaaga actcagacga	1260
taaattggct tccaaagaag gccaaataga aaaaaaaaag gttcttcaga ccaatagctg	1320
agtgacctga aatttaaagt caaaaaatgc attaaaaccc acaaaaccac ttcctttaaa	1380
attcttgata ccaaacatgt tacagcatta agtgaacctt aatctaagca aaattcagag	1440
aagagcatte acacctttga tagaagagcc agccagccag ggtgctgggg ctcagctctt	1500
gagaaaaatg ttgatgtaaa agccttgggtg gcctgggttca gcccctgagc ctccctggta	1560
agcacatggc aggtcagaa catccataac tgaatgctct gagccctgta aaaatgcagt	1620
tacaatggga aatcaactct taagtatgct atgggatgct gtgataaaat ttgagtttct	1680
ctaccttctg tttgcacttt ggttcattta gaaaagagtc cataattgtg tgcttcaaga	1740

47

gtgcaattcc	atggctgtaa	cagaatagag	tctcttttct	tctggccttt	gcctcttttg	1800
ggatattcca	actctgtgga	aggtgatcat	ttgcgattat	gatcagttat	ttatatattga	1860
ctgtaaata	aaacttcaga	gtcagtttca	aaaaacaaga	gatggacata	aggacatgtg	1920
cttatgagta	ggggacaaat	aactagagac	aaaaaaaaaa	aaaaaaaaact	cgta	1974

<210> 92

<211> 1423

<212> DNA

<213> Homo sapiens

<400> 92

ggcagcagat	actttcttat	ggaacttgta	tggtttcggt	ttttacattt	aaaccttctt	60
ccccgtggtg	tgtgttggtg	aatctgtgtt	tgtgtgagga	ggggcatggt	gctctcagaa	120
cccacctcct	gtggccagag	agccctgtcc	tgtgaggggtg	gttgtcacag	tggcaggggt	180
caattcagaa	gaccttgagg	gcaggctgat	gtttcctgaa	tgggcccctg	gttggttgctt	240
gtccctgact	ctccatttcc	ccatctgagt	ggatttggac	ctaatagggc	actggagctg	300
gttcgaatcc	tgactggact	acttggcaac	tttatgtctg	ggagcaagtt	acttaacctc	360
cccaagcctg	tgtctgtgaa	atgcgggtaa	atgaatgtag	atgtttggca	gcagctactc	420
cttggtgagc	tctcacagt	aactctcctg	cctctgccct	ccttccccgc	ctccccgtgt	480
gcctagcgtc	aggtctagcc	acttctcctc	gggcccctct	cccttttctg	tggctggctg	540
cctgcccgcc	tggcgtgga	cctttcatgt	aacgggaatc	agcatgtata	ttctggctctg	600
gtctgtttct	acacttaatt	ttgtttccag	tagtatttcc	ctgtaccggc	agagttcaca	660
aacacatttg	aagaggcttt	ttctcaggat	tcttaacctt	cccaaaggaa	gtcccatgga	720
tgggtttcta	gaagtctata	aaatgctctg	aaattgtatt	tttctgtgtg	aaagcataac	780
tttcatctgc	ttgttcgtgc	tcaaaaaaga	tcatgaatga	atgattgcat	gattttatgc	840
cattgtgctt	atactaaagg	atatgtagcc	catctcttga	gctgktaaac	tgttttgact	900
actttaaatc	gtgcagctgt	gagcatctct	gtaaatttag	tgtacacatg	tatcccctgg	960
agtggcattg	cctcggcagt	gagcacttat	ggttttataa	ctctcttcac	agactcaaat	1020
gactccagaa	agctacactt	cctgttggtga	gtatatgata	tccatttccc	tacatagcca	1080
ctaacatcag	gtttttacaa	ttttatttat	ttcttgctac	tttaagaaat	ttttgtgggtg	1140
aaatacatat	aatagaagtt	gactatctga	atcattttta	agtatacatt	cagtagtggt	1200
aagtatgtcg	ccattgttgt	acaaccaatc	tccagaactt	tttcatcttg	caaaacaaac	1260
tctgtaccca	ttaaataaca	ttaaacattc	cattccctcc	agcctcagca	acccctattct	1320
actttctgtt	tctgtgagtt	tgactatttc	aagcacttca	tatcagttaa	atcatgaagt	1380
atttgtctgt	ctgtgactgg	cttattttct	tgagcacagt	gtc		1423

<210> 93

<211> 1365

<212> DNA

<213> Homo sapiens

<400> 93

ggcagagcta	acccgagtga	agccacttcc	gggcttcccc	ggcgccttcc	gcagtcctct	60
tccgggtgat	ggcgccggg	tgccccgat	gtagccctgg	cgcaagatct	cttctttttt	120
ccacctcgcc	ttccgcgat	tcccagcttg	agaaacacct	ctttgccccg	tcatgccaaa	180
gaggaaagt	accttccaag	gcgtgggaga	tgaggaggat	gaggatgaaa	tcatgttccc	240
caagaagaag	ctggtggacc	ctgtggctgg	gtcaggggggt	cctgggagcc	gctttaaagg	300
caaacactct	ttggatagcg	atgaggaggga	ggatgatgat	gatgggggggt	ccagcaaata	360
tgacatcttg	gcctcagagg	atgtagaagg	tcaggaggca	gccacactcc	ccagcgaggg	420
gggtgttcgg	atcacaccct	ttaacctgca	ggaggagatg	gaggaaggcc	actttgatgc	480
cgatggcaac	tacttcctga	accgggatgc	tcagatccga	gacagctggc	tggacaacat	540
tgactgggtg	aagatccggg	agcggccacc	tggccagcgc	caggcctcag	actcggaggga	600
ggaggacagc	ttgggcccaga	cctcaatgag	tgcccaagcc	ctcttgaggg	gacttttgga	660
gtcctatttg	cctagagaga	cagtggctgg	ggcactgagg	cgtctggggg	cccaggaggg	720
aggcaaagg	agaaaggggc	ctgggcaacc	cagttcccct	cagcgcctgg	accggctctc	780
cgggttggtc	gaccagatgg	tggccccggg	caaccttggt	gtgtaccagg	aaacaaggga	840

48

acgggttgct	atgcgtctga	agggtttggg	gtgtcagacc	ctaggacccc	acaatccccc	900
acccccaccc	tccctggaca	tggtcgtga	ggagttggcg	gaggaggaac	tgagagcccc	960
aacccttacc	cagagaggag	aagcagagtc	gcggggagat	ggtctggtgg	atgtgatgtg	1020
ggaatataag	tgggagaaca	cgggggatgc	cgagctgtat	gggcccttca	ccagcgccca	1080
gatgcagacc	tgggtgagtg	aaggctactt	ccgggacggt	gtttattgcc	ggaagctgga	1140
ccccctgggt	ggtcagttct	acaactccaa	acgcattgac	tttgacctct	acacctgagc	1200
ctgctggggg	cccagtttgg	tggggccctt	tttccctggac	tttgtggagg	aggcaccaag	1260
tgtctcaggc	agcagggaaa	ttggaggcca	tttttcagtc	aatttccctt	tcccaataaa	1320
agcctttagt	tgtgtaaaaa	aaaaaaaaaa	aaaaagggcg	gccgc		1365

<210> 94

<211> 756

<212> DNA

<213> Homo sapiens

<400> 94

agcagcaggg	tgggaatgtg	aagagggcag	cccaggccct	gtgtttcggg	ggtgtgcctc	60
tccccgcact	cccgtttctg	ggaatgctgt	tccttctacc	ttcagggcct	gcccgccttg	120
cgggtgtagc	gcactcctcc	cgggtgtcat	ttcttcagag	tctttccttc	atgccctttt	180
tcctcctcac	gttccccctc	gtgctttacc	cacatctgtc	ccgtgggttc	gatccagttc	240
tcccgctcgt	tatgggtatc	cacgtgtttg	gtcttagcca	tcactccagg	aaagtggctc	300
ctccctgagg	acagggctct	gtctctgatg	ctgctggcct	ccctccagtg	ctgccctcct	360
ccttttgggg	cttgggtgat	gcaagtgctg	acacacaaag	gcaggcaggc	aggcctcggg	420
ccaggggtgt	cctccaggcc	cctctgatgg	cagggaccag	cgagacctgg	ggaaactgca	480
cagtctgtgc	ctgtctgtgc	tctcaggaga	ctgagggaat	tagtgggtgg	gagagatttt	540
atgttggatt	aaaaactgca	cctccaggcc	gggcatggtg	tctcatgcct	gtaatcccag	600
cactttggga	ggccaagggtg	ggaggatcac	ttgagccctg	gagtttgagg	cttccaggaa	660
gctatgatta	caccactgct	ctccagcctg	ggcaaacaga	gtggcaccct	gtctctaaaa	720
acaaacaaac	aaacaaacaa	aaaaaaaaaa	aaaaaa			756

<210> 95

<211> 938

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (479)

<223> n equals a,t,g, or c

<400> 95

ggcacgagtg	gaggttcagg	gtcagggtcc	aggggcgaag	gtggacgctg	gagaagggca	60
gatggacagg	gtcaggttca	gatcttggct	cttgtatcct	tgctgtgtgg	ctctggggcca	120
agaacttggc	ctctctgcgc	ctcagtggct	cattacagaa	aatgggatgc	cagcacttgc	180
cttagtgggt	tgttttgagc	caactgcagg	ctcaggggagt	agctggcatg	atgtgttccct	240
accctgaagg	gcagaaaaag	gggaaggagg	ccaccagatc	ccacagggtg	gtccccagggt	300
ccctccccgg	gatgggcagc	akcctggctg	ctccccacag	caacccttgg	ctggcccccat	360
tggcgtcgt	ggaaatcccc	camccagttc	tttgtgaatg	gaaaaggaaa	ctgattgcgc	420
tagaagaggt	ctccgaatgc	cggccggggg	tggggggcgg	ggggggsttc	ctctcccant	480
gcaggagggg	ccacctcagc	ttcctttctg	gggccccata	cccccttttt	cctatctccc	540
ctctgasttg	aggaggaggc	tgtggccccq	gcccactgtt	tggcctcttg	aaccactggg	600
tgacagtgtg	catcggggcg	attaagggcc	acctggcctc	tggacctccc	tcgctgtggc	660
tgctgcctgg	gacccccccac	cccatgcctg	actcgaacct	caacctcagc	tctacacaca	720
ggggcaccac	tgtaggagga	gagagaaaacc	tctctctagg	gaaccacctg	gaagggggcc	780
ctgccctgtc	ttcatctggg	taccccatgt	aatctaggaa	actgtcttta	acttgccgag	840
ggcctccatg	tctgagttcc	taatgttttt	tattttgctt	tttcaattaa	taaagctcat	900

ggcagaaact atttaaaaaa aaaaaaaaaa aactcgta

938

<210> 96

<211> 928

<212> DNA

<213> Homo sapiens

<400> 96

ctgcaggaat	tcggcacgag	gcagggtggg	tgcatacaagg	gggttggtg	gtggctgaca	60
agaggttgta	atgtgggaa	gggcaggagc	gggctggtct	ttaccactg	ccatccccgt	120
cctcttatta	aatagagatt	cctagtattg	ctgttataaa	aaatgtcctg	ctaaacatag	180
ttctctcaaa	aatatttttc	tgagattctc	tgacaattaa	atctggatcc	cagccccaat	240
attcacctgc	aattacatta	tggatgaaat	taaatgtgca	ctttctatgg	tgtacattta	300
tttttcaaac	ctcaggaagc	catattgagc	tcttaatctc	aggtaagtg	tcttcttata	360
tccctccct	tgatttttgt	actcataagg	ttgtatccag	agagaagttt	gaggaataat	420
tcattacctg	ggaattaaag	taaaccttga	gcttgggagt	cctaacatat	aaccatctct	480
aaaattctgc	aactgtagat	ttttaatcat	ctaattttag	ggacttcaaa	atatttttct	540
gactttacct	acattcgaat	taagttaaaa	tagcactgat	aatggatagt	aggatccaaa	600
cagaaacatt	ttaaatgaat	ctagttaagt	attgagccgg	gcacagtggc	tcacacctgt	660
aatcccagca	ctttgggagg	ccgaggcggg	cagatcacct	gaggtcggga	gttcaagact	720
agcctgaaga	aaccccatct	ctactaaaaa	taaaaaatta	gctgggcgtg	gtggcacatg	780
cctgtaatca	cagctactcg	ggaggctgag	gcaggagaat	cacttgaacc	cgggaggcag	840
aggtttcagt	gagccgagat	cacaccattg	cagtcacccc	tgggtaacaa	gagtgaatc	900
cgtctcaaaa	aaaaaaaaaa	aactcgta				928

<210> 97

<211> 1715

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (17)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (34)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (40)

<223> n equals a,t,g, or c

<400> 97

cacggcctat	ggtttangtt	aaaaccctta	attntttttt	cggtaatcac	ttttaacact	60
gctaattgaa	attatgtttt	catgtgtttc	acatgaaaaa	gtatatatat	acaagggtta	120
tatagtgggt	aaaatacttt	acatagtcac	acattttaca	atttttcaag	aggtttagcca	180
ctaagacttt	aataatttta	caagggaaaa	agcctttttt	tttytttgat	atacagtttt	240
ttcttcttag	ttctgcatta	gaaatggcat	ctgttttagg	tctcaaaata	taactcggct	300
gtttcactgt	atatgtacat	tgttttctgt	aggaataggr	taatgatata	taggatcatg	360
atattccttc	tatccatgtg	ccaaattggg	gtaatgttta	tttactgatg	ctttatgtta	420
ccaaaacata	cagtaaaaaa	gtagaaattt	atgaaatacy	tttgataaaa	agttttattt	480
gtgcttacca	aaaggaatgc	tttcacaata	gtgtatcagt	tcttttggtt	tgttaaagtt	540
ggaattttatt	ctgttgccag	catttaagta	gtcatggcaa	gtcctgtttt	taagaccttt	600

50

tgagagactgg	agcttttctgt	tccattaagt	cttttgttta	tactacaaat	tgacacctca	660
cttagttcag	atgaaatctg	ttactctaca	aggaagggtg	tcatcattag	gaggcagctt	720
tactaagctg	gtgctttgca	tggtagcaag	tgctgccctt	tatcagcacc	ctgggtcata	780
gtgtaggcta	grgttaaggc	actggcagac	ttagggatgc	tgacacagacc	tgtagttcgt	840
tttaagtcac	gttcacagga	atttctacaa	taataaaccc	atcatctcca	taggtcagat	900
cgaagtgcac	tccaatgcta	aatagaagtt	atgagtgggt	ttaacaattt	tagatgattc	960
agcttttgtt	ccattactgt	tgaactatat	gaactattcc	attactgcag	agattttaagt	1020
atctgtttta	ataagctctt	tttggtattt	aaaggctgcc	catgggtttc	tgcctagtgg	1080
taaagctgat	tgttaccctc	ctttgaaatc	ccttctagtt	ctgagatgct	ttgagggtaa	1140
ctggattcga	ttttgggata	tcttttctca	cattcagact	tacacttaat	ggtgttagaa	1200
atcaacaaaa	ctccttttta	aaaagaaaag	atattaagcc	tgccacttcc	tacaatgcac	1260
tctgtttacc	atttgaacag	tatgtttgta	actatggcaa	tgaagtcagt	agataggaaa	1320
ccagttattc	cttctacctt	taaaaatttt	gagaacttgc	caaccaggga	ttaaagctat	1380
tatcttgaac	agagtcccta	aagctagtct	agtttttgcc	acatctgcaa	tgattattgt	1440
ttaatttcaa	aagaatcttc	aggctctaca	atctaggggt	ggtaaatgtg	tttccactat	1500
acttgggaaa	aggctcagtag	gatgtgcac	ctagggaaga	taaaatcgta	tatggtaaag	1560
gcatttgagt	taattttgca	ttatatctag	gaaccatatt	attttaaatt	tgaatcctat	1620
taatgctgag	agatcctaag	agctagtatg	ttgtaaaacc	tgccacctga	ataaaatgaa	1680
aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaac	tcgta			1715

<210> 98

<211> 678

<212> DNA

<213> Homo sapiens

<400> 98

acattttcta	tgtagtaaac	atgtatgtgt	atgtgtgtgg	gtgtgtgtct	aattactttg	60
ttggacagat	tcctgtgggt	tggaactgct	ggggccaagt	ttatacaaaa	atctacattt	120
ttgagtaaac	tccccatgac	tctagtaagt	ttccacagta	tttgagtatt	ctttatttcc	180
taaagctgta	caaacatcag	ataatcaata	tttgtaagga	tcgataatat	aattttataat	240
gaaattgctt	tatcagctat	taatgttaac	tacatcatct	tcataatagcc	ttataactca	300
tttgtgctat	tccattttcc	tctgttcctt	ttattttcac	ttcccttgta	atgttagtct	360
ctttgtactg	attttctgaag	agttcattta	tgattaaaca	tgtaaacatt	ttgtctagaa	420
ttgcaaatat	gtttttctc	attcatttta	ctatgggtgt	ttattttttg	gttatacaga	480
agttgtatat	tgaaatataa	tcttgttctg	ttttatgact	ttggagtttt	gtgggttttt	540
aaaacatggt	tacattggta	tcattttatt	gtattgcctt	ctatttcagc	agttagtttg	600
tcattttctt	ttccattaat	catcattctg	gtgttaatga	atgcattaaa	tattttaaagt	660
aaaaaaaaaa	aaaaaaaaaa					678

<210> 99

<211> 1541

<212> DNA

<213> Homo sapiens

<400> 99

ggcacgagtc	gcatcaaagt	cagaaccagc	gccctccgtc	cttgctctca	gcccagtgcc	60
aggcttcctt	gttaggagta	ttcttctcat	gaaattctac	tgattttctta	cattgtagcc	120
cagatatgat	ttgctgaggg	atatctgaga	gaaagcctat	gtgtcctttc	cataaagcgt	180
accttgactg	cttttttcaa	atctctctct	tgctactgat	ctttctaact	taccttgata	240
ttgggaagtg	tggtcttttg	agccatgaat	ggagaattag	ggaattaggg	aaacatgaga	300
ggtggtggaa	ttaattcatt	gtatcttgat	tatggtggcg	actatgcagg	aatatgtgat	360
gtcaaaatca	ttgacttggt	cacctaatgt	gtgtccatct	taatgcttgt	aaattatacc	420
ttaacaacgt	taactaaaga	agtaattaaa	ctccacatt	ggacaaggag	ggatctagag	480
cactgaatct	gcacagattc	tctgagctca	gtctaggaag	agaagtgggg	ctgaactata	540
cagaagatgt	tgaatttaga	aacgtacttt	ccaggaataa	attaaaaatt	gagtttttgt	600
ccacctacaa	taatcaggca	ctgtgccagk	tcctaggact	gtaaagaaaa	ctatctctga	660

51

tataagctta	taatctaaca	gagaagacaa	aaagtagaaa	gcttaaattc	agtgagtaag	720
tgctatgata	gaggcaggag	tgggttgcaa	aggagaagca	tcaaatacag	atgaatggca	780
caaaaggctt	tctgtaggac	attcatatag	agcctgagaa	atcaggagaa	ctcctccagg	840
atcactagct	gtattagttc	attctcatgc	tgctaataaa	gacaaacca	agactgggggt	900
wacttataaa	ggaaagargt	ttagttgact	cacagttcar	caggggtggg	aggcctcagg	960
cagcttacia	tcattggagga	aggggaagca	aacgcgtcct	ccttcacatg	aaggcaggaa	1020
agagaagtgc	ccaagcaaaa	ggggggaaaa	gtcccttata	aaaccatcag	atctcatgag	1080
aacagcatga	aggtaaccac	ctccatgatt	aaattacctc	ccaccagggtc	cctccacta	1140
cacctggaga	ttataggaac	tacaattcaa	gatagatttg	gggtggggaca	cagtcaaacc	1200
atatcactag	ccaaagcagc	aggtgtgtgt	gtgtgtgtgt	gtgtgtgtgt	gtgtgtgtgt	1260
gtgtgtgtgt	gtgtattcag	tatcttgaac	tagaccactg	ctcttgaaga	tctcttgacc	1320
atgtagaagc	caggataaca	catttagctg	gaagctaatt	gtgaagctgg	gaaattatgc	1380
atgtatagt	atgttgggca	caaactcatg	ttatgttcag	gtaacaagca	gagcattttg	1440
ggttcctttc	tttaggagtt	gtctaatttt	acttcatttg	ttatttctga	ttatattggt	1500
taccagctcg	tgccgaattc	gatatcaagc	ttatcgatac	c		1541

<210> 100

<211> 881

<212> DNA

<213> Homo sapiens

<400> 100

ggcagagccc	agctgtttct	tacacagtga	catcccaggt	gccctgggggt	ttgggactcc	60
tggcaggaga	gaagagaatg	aaccaacca	ttctgagaag	tcaggcattg	ttatggccct	120
ggagatgggt	ggtgaaggca	aaaccatgtg	tgtgtgtgtc	tatggatgcc	tggattcctg	180
accgcagtca	gcattgccc	tcaattccag	gtcaaaagaa	ggaaagggct	gggtcccatg	240
gtcaccaagc	ccttgccgyt	ctgctcttct	tgtgaaagt	accttcagag	tttcacagc	300
tgatagcaga	cagccagatt	gccctctagt	aaagcctagt	ttaagatcag	tttcaggaat	360
tttctgggtg	tttttatctt	tgcagtgaag	ccataaaagc	cctcatagca	gctgggcarg	420
ctgcaaagtt	gatgcttccg	agtggctcat	tagagtgtaa	tggcattcct	gagagtagat	480
gttgctgcaa	gcagcaggca	ggaagctcat	gcggcagcag	cctgatggat	actccgcgtc	540
aaggggggtc	tgggtgatgc	gtggcaggca	ggcagcggct	acactccatg	ggcgggtgctg	600
ggttgcgaaa	ggggctgaca	gtgctgctct	cggcagagg	gggtgggggca	gggtgcagca	660
cattgcagat	gaaaagggtga	ggggtctcag	tgggtgtgat	gggagttaga	caacagcact	720
gtttaggagg	tgggttaaag	ctttttctct	aatttaaatt	gggtagtaaa	ttctgaccct	780
aacatgagta	ttgtatctga	tgggattcag	tttggggaaa	aattcagctc	tgaagtctag	840
caaattccag	tttcggtctc	aacctttggg	tcattctcgta	g		881

<210> 101

<211> 947

<212> DNA

<213> Homo sapiens

<400> 101

ggcacgagca	gattgtttct	ggcttagtag	catcatctaa	tggcgagcag	gggcactgca	60
gcccctggac	gcaccttctt	ggcgatgatg	gtcacgtcct	ttttcttttg	tatgagatgg	120
ggatcatggg	ctgagcagat	gccccaaagg	tgtctgccat	gctgtatgca	ggaatgctga	180
ggtttcatgc	aaaaacgtat	gttgggatca	cttgacgggg	tgcattggaaa	gtatcctagc	240
agagttttgc	tagtagagaa	tgtgccctcc	gctagtactc	tagagtcaca	gccagatccc	300
tagccttcgg	gtggaggtgc	cgccttatga	acggccatca	gactgtccta	ttgttcagag	360
actccttttc	gtcccttact	acttcactct	aggcctgtga	ggatctctac	caaagcagag	420
caagcccata	aataaatagc	tttgtgtcca	ccaatgcata	agtggcgctg	aaataactgt	480
gcagtggaca	ccactctccc	tccctctggg	cctttcagct	gctctgtgtg	tccacgctgt	540
tgtgtgccct	gaaagaactt	tgagatttgc	tgaaaggtct	ccagctgcaa	tttcagacca	600
tgtgggtaac	tcaaattgtct	aacatttaaa	tatggctgaa	aggggtcgac	tgtgtggctg	660
gatggtggca	gaagtagcag	gctttccctt	ggatctcttt	acagggcaga	gatgactcct	720

52

tacaaatggc	accagtaatc	cctctgcctc	ctgataccca	tctaacagaa	aagtgcaccc	780
tgctcataag	aaaaccccag	gccacatttg	tcttatgtga	ccctggcttc	tggtctacagc	840
tgacttgaca	caaactaaga	caatcagatt	tttttcttct	ggaacttgct	gtgagacaca	900
gagacacagt	ggactggatg	acaatgctgg	gatgtgaagg	atgtgtgc		947

<210> 102

<211> 1369

<212> DNA

<213> Homo sapiens

<400> 102

cccacgcgtc	cgcccacgcg	tccggctggc	aagatggcgg	gaggggtgcg	cccgtgmgg	60
ggcctccgcg	ccttgtgtcg	cgtgtgtctc	ttcctctcgc	agttctgcat	tctgtcgggc	120
ggtgaaagta	ctgaaatccc	accttatgtg	atgaagtgtc	cgagcaatgg	tttgtgtagc	180
aggcttctctg	cagactgtat	agactgcaca	acaaatttct	cctgtacctt	tggaagcct	240
gtcacttttg	actgtgcagt	gaaaccatct	gttacctgtg	ttgatcaaga	cttcaaattc	300
caaaagaact	tcatcattaa	catgacttgc	agattttgct	ggcagcttcc	tgaaacagat	360
tacgagtgtg	ccaactccac	cagctgcatg	acggtgtcct	gtcctcggca	gcgctaccct	420
gccaaactgca	cgggtgcggga	ccacgtccac	tgcttggtga	accgtacttt	tcccaaaatg	480
ctatatgtca	attggactgg	aggctataag	tggctctacg	ctctggctct	aagcatcacc	540
ctcgggtgggt	ttggagcaga	ccgtttctac	ctggggccagt	gsggggaagg	cctcggcaag	600
ctcttcagct	tccgtggcct	gggaatatgg	acgctgatag	acgtcctgct	cattggagtt	660
ggctatgttg	gaccagcaga	tggctctttg	tacatttagc	tgtgggtgtg	gcttcagaaa	720
ggagcagggc	ttagaaaaag	cccttttgtc	cgtagagttg	atgtggtgtg	agtgatatat	780
ttctatgttt	ttaatgtaca	gcactctgtac	tttgtttgcc	ttgataaagg	taagataaat	840
gaaacgctga	actatgctaa	tctggaattt	gtttttattt	gcctgaaata	tatttttttc	900
tgtgaaaaaa	ttaaaacgta	cttaagccag	gagaatgaat	tatacagtga	ttgaaaaatc	960
atttaattcc	tatgactttt	gttttgtatt	gcccaagtca	aactacatca	cttgatcttc	1020
cagcccaaat	gtagtctgcc	ttgaaaagtc	tttcagctgt	gactgcagga	agtgaggagt	1080
tttttattgt	tagctaattg	ctgtgactgc	aggaagtggg	agtgtttctg	ttgttggtca	1140
attgaagtta	ttaggctcag	cttcagtcac	gtgtaagttt	tgcagtgtaa	tacatatgta	1200
gtctggctctg	tatatatgaa	aatttgaaat	aaactgcaga	atgtttatgt	ctagttatgg	1260
tttaaatttt	cttagtagta	tataaaagg	aagagtactg	aaaaattaat	aaaattgcaa	1320
gttaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaag	aggggggggc		1369

<210> 103

<211> 1231

<212> DNA

<213> Homo sapiens

<400> 103

ggaggaagga	agcaattcta	aaaacaaatt	ttcaaagcta	tttttacatt	aacattattt	60
tttctaaatg	tgcatccatt	atagtagagt	tatcttttcc	ttccttaagc	tcagaattaa	120
gtcaatttcc	tggtatgcaa	tgtggcttta	ttttttttgt	tatttaaatt	gtttagccaa	180
agtgagatca	gcaactactt	atttatatgt	ttgagggcac	tgtgagctga	aataacaaat	240
tttgtataac	ttatctaact	aaaaatat	ataatacttc	tcaataatga	ttctaataga	300
gatgtattga	gtaatttatt	accatctttt	atgttttaag	agttctgata	gtacaagtta	360
aaaaaaaaata	gcctacttct	actcttttac	tgaaatcaaa	tagatggcat	gtgatgatta	420
acataaaaaa	gaacacattc	aacataatat	attgcgatag	tggcagtcta	atagaaaaat	480
agcaactatt	tatttacttt	tatttattcg	ttttatttat	aatagatatg	tacaaaatgt	540
aaaattattt	ctaagaagat	ttttctttta	acaccaagca	ttacagtaga	tgttgtctga	600
atgtgtgtaa	ttgagaataa	taatattgta	taataatat	aatatgattg	tggttagaag	660
gaatttattt	gatcacataa	gtacagcaaa	ttaagaataa	gatctcagca	tatctggct	720
actacataaa	tgtagcaaag	ttattttgct	accttgaaaa	aatacaagtg	gacaaaaaat	780
tgcctaattt	catggaaaaa	ctataatttt	gccagttaat	acatgagatt	ctcaaattca	840
cgaattttaa	atgttttaaat	aattatactt	ttgctgcctg	aaaatatatt	ggatcaaaat	900

53

ttcaagtaaa	tatcatcagg	ggatttggat	ggggaagagc	aagaggggaa	gaaggaggag	960
aggagattta	gtctagcact	atttccagct	gatccccacc	cagcattgcc	aagctacagg	1020
acaccaattc	aataggatga	agtcagtcaa	atcctctttt	aagtaatggt	tgaataaaat	1080
aaaatgaggg	gaagaaaatc	aagtactcta	aatgattaaa	gcgtatgaat	aagtgaaaaa	1140
tgtttgcaaa	attgagaagg	aaggagaatc	gggaagtggg	tatttatggg	gtctcaatat	1200
gtaagggtaa	aaaaaaaaaa	aaaaaactcg	a			1231

<210> 104

<211> 1242

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (288)

<223> n equals a,t,g, or c

<400> 104

ttcgtatcca	ctaggatggc	tctaataaat	aacaaagtat	tgtcaaggat	gtagaaaaat	60
tggagccctc	ctgccttggg	gggagtgtaa	tatggtgcca	gatacaacct	ccatcctgaa	120
gtcatctctg	atgcttccctg	tttgtgtttt	taaactttta	ctatatcttt	atgtcctcat	180
aagaatatgt	actatcattt	gggtgtttta	agtgtacata	aatgctgtca	tcctgaacaa	240
atcctctcgc	taactgcata	tttaactcta	tactatattt	tcaagatntg	tccatgttga	300
tccacgtagc	tccctagttc	cctttaactg	ctataagata	ttctgttgcg	tcaatatatg	360
acaatttatg	catgctttgt	tgacaggtaa	ttggattttt	agtgttttgc	ctttacaaaa	420
atcactgcat	cttttgcaca	tgtctacttg	tgcatatgaa	ctgaggtaaa	attgctgggc	480
cttactgtaa	atatgttggt	ttaattcact	ttgcgctgct	gtaacagaat	accatagact	540
gggtgcttat	aaagaaaaga	aattttattc	tcatagttct	ggagaatggg	aattccaaga	600
tccattcaca	ggttcgggtg	tctggggaar	actttcttca	cacatcctca	cttggcagaa	660
cgaaggggcc	tgggttgatg	ctgtgtgaam	cctcttttat	aagggcctta	gtctcattcc	720
caaggaggag	ctctcataac	ctaatacact	cttaaaggcc	ccccactcaa	tactatgaca	780
ttgaatttca	acatctgaat	tttagagggg	acactgcaaa	cctgtcatat	gtcttttatct	840
ttactatcac	ttaaattgtcc	aaagtgattg	caacagtgat	ttatatactc	aaccacaga	900
gtataagaat	ttctcctttc	tagctgggca	cgggtggctca	cgccagtagt	cccagcactc	960
tgggaggccg	agatgggagg	atcacttgag	gccaggagtt	caagaccagc	ctggccaaca	1020
cagtgaiaacc	ccatctctgc	taaaaataga	aaaagttagc	tagctatggg	ggcgacaccc	1080
tgtaatccta	gctatttggg	gggctgaggg	aagagaattg	cttggacctg	ggaggctgag	1140
gtagcagtga	actgagatcg	taccattgca	ctccagcctg	ggtgacagag	cgagactctg	1200
tctcagaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaactcg	ta		1242

<210> 105

<211> 1151

<212> DNA

<213> Homo sapiens

<400> 105

gcagggacag	ccccacgcat	gggagggtgg	ggccagccat	caccccaggg	tgaattttctc	60
taagagcagt	cctaggccag	ggggtgagtg	tgggaaagtc	agggccggtt	ccatcagtta	120
acgtttcaga	gcgggttccc	gctggaagtg	ctggaaatac	ccagatgcca	tctgtagcat	180
tttatgaact	gtggcggatc	cactctctgt	gtcttgagct	tctgctctgt	ggtctgctca	240
gtggaagcat	cctgtcagtc	cactgtacag	tgggggtggag	cgcagccag	ggtgggtgtg	300
ccgtttgatt	ggtcacggaa	tgaacagggc	aaaggctcact	aagtagatat	gatactgcaa	360
gtaggattgt	ggtcataggt	atttttatga	atttgtgtta	tgtgaagttg	aggatttgaa	420
tgttgtgatt	attatatcca	gataaagttc	tagcctggca	cagtgcaggc	acgtgcctgt	480
ggtcccagct	gcttgaagtg	ggaggagagc	ttcagctcag	gatttccagg	ctatagggag	540
ctgtggtccc	accattgcac	cccagcctgg	gtgacagagt	gagaccccat	ctcaaaaaga	600

54

aaagaaaaga	ggctagggcgc	agtgggtcag	gcctgtaatt	caagcacttt	gggaggctga	660
ggcaggcgga	tcacttgaag	tctggagttc	gggaccggcc	tggccaacat	ggtgaagccc	720
ccccatctct	attaaaaata	caaaaatcat	acttaccggg	caggggagat	accatgatcg	780
cgaagggtgt	tttcccagg	caaggcttag	ccattgcact	ccggatgtgc	tgagcctgcg	840
ttttcccaa	atgtgggaaa	cttgatgcgt	aattttgtgt	agtggaggga	ctgtgttcac	900
actgtccccc	ccgccccaaa	aaaataaaaa	tacaaaaatc	agtcgggcgt	ggtggctcac	960
gcctgtcatc	ccagcacttt	gggaagccga	ggtgggtgga	tcacctgagg	tcaggagttt	1020
gagaccagcc	tggccaacat	ggtgaaaatt	gggaggccga	ggcgggcgga	tcccaaggtc	1080
aggagatcga	tacatcctgg	ctaacacagt	gaaacccgt	ctctactaaa	aaaaaaaaaa	1140
aaaaactcgt	a					1151

<210> 106

<211> 1628

<212> DNA

<213> Homo sapiens

<400> 106

gaggaaatat	cctctccatg	aggcatacac	caagtaaattg	actttgtaac	tttacttcat	60
cctcttcatt	tacacagggc	atacatgaag	taaccaatgg	aatcctctag	ggggtattta	120
aactcccaaa	aattctgtaa	cggggccctt	gagcccctat	gcttgggtcc	attcccaaac	180
tgtggagtgt	actttcattt	tcaataaatt	tctgcttttg	ttgcttcatt	ctttccttgc	240
tttgtttgag	cgttttgtcc	aattatttgt	tcaagacgcc	aagaacctgg	sacacctcc	300
accggttaaca	ctcttacaat	tatgcagttg	tgcagtgcat	agcccgtgcc	actatatgtg	360
gcagtgttgc	ccacttagca	atgaggagcg	catattttcc	tgcattatca	ccaaaacaat	420
gttatcatct	tttattttatt	attatttttt	tgagtcaggg	ccttgccctg	tcaccagggc	480
tggagtgcag	tgacgcagtc	tcagctcact	gcaacctctg	cctgtcaggc	tgagggtggga	540
tgatcacttg	agtccaggag	tttgagacca	gcctgggcaa	catggcaaaa	ccccatctct	600
acagaaaata	attagctgga	tgtggtgatg	catgcctgta	gtcccagcta	yticaggagac	660
tgagatggga	agatcacttg	agcccaggag	ttagargctg	cagtgaagta	tgatcatgcc	720
accgcaatcc	agcttgggca	acagaatgaa	aytctgtcaa	aaaaaaaaaa	gaagagagaa	780
aagaaaaagg	aaagtaaaaa	ttgctaataa	cytccaaagt	ctatgtagga	aatatgtaaa	840
tgggtggcca	ttccaagat	gaagtgcctc	aagtaggcct	gggtctgcca	gctacagaag	900
gacagaatat	cytaggccct	tgcttcaata	gctggagcct	gcttgttggg	gtgcccttag	960
ttgctctcat	cgaaccta	gagtttagtc	tagaatgaaa	atttactagc	ctgcaaaata	1020
gctcacttta	tctattcttt	tatcagcttg	cctgactacc	taggtcatag	gtcaaatact	1080
taaaaagccc	ttgagcagac	tataattgca	atgcattatg	ggctgcaaca	aaatgcaccg	1140
agacaaccct	aaagaaaaca	cccaaaaccc	ctacctggcc	aggcgcggtg	gctcatgcst	1200
gtaatcccag	cactttcgga	ggccgaagcg	ggtggatcac	ttgtcaggag	tttgagacca	1260
gcctggccaa	cgtgggtctc	cataatactc	agcctatgag	gaaccaggag	agggacctgc	1320
acactagagg	ataaattgct	tgttgtaact	gtgcggggta	tgctgcca	ccagacacct	1380
gatcttgcaa	gactgtatta	aaagtctcac	ttccgctatt	ctccgtgtct	ctgagtccat	1440
tctttgggtt	tggacgggtg	agtttgtttc	tcacagtcta	gactctagat	gtgaagaact	1500
ttgatcatat	caccaaagga	gatggtggta	tgcaatttta	taagtaaaaa	tacactagtg	1560
tcagtttttt	ttacaaggga	aacctgattt	gcactttttt	aattaaaaaa	aaaaaaaaaa	1620
actcgtag						1628

<210> 107

<211> 1465

<212> DNA

<213> Homo sapiens

<400> 107

ggcacgagcg	agccaagttt	gcaccactgc	actccagcct	gggcgacaga	gcaagactca	60
gtctcgaaaa	aaaaaaaaagtt	ggaagcagaa	gtaaaaaaca	tggtaaagaa	tgagaactaa	120
ataaatataa	taattgagag	gtctgcatta	gatgtggcag	ggagaacaag	caaaaagaga	180
tttcagagaa	gatcactgga	attggcagag	gccttgaaag	gcagagtcta	gcatacagaa	240

55

gatgtaaagc	cacattctgt	gaaggtaagt	agatgtggtt	acctcttttg	cactgtactg	300
gtgcattatg	gggtaaatrt	gtattacttt	tcctgtattg	cttagcacag	agttttgcct	360
atagcaggca	ccagactgtg	ggcttggtag	tacatgacta	ttggtgatta	cagatcaaaa	420
aggacttgaa	atgatcagtt	taaggctctg	atgggtattg	aagactcaaa	ggatgatggc	480
accctgggag	tgatccacag	aaggacagat	tatttgaa	tgtaataaac	taaagacaac	540
atggatgtta	aatgatgaaa	aaaagttgga	tggaaaataa	accattggat	ctgcytctgg	600
agtccaagaa	gaatattatt	cttcctacct	cccccttact	ctggctcttc	ctattgtagc	660
cacatgggtc	agtaatgcca	ttgaaaaaca	aaatttttaga	ctaagtgggg	tcgcagaaat	720
tttgggtctat	cttaaatgga	tgacatctta	ttaaagaaty	tattgtataa	agtgtgctta	780
ttctggcatt	tttttaatga	agaaaaagtg	taattcagtg	cacatttatg	aatttcaaa	840
atcaataaaa	atgggcaaa	tatatgaacg	cataatccat	agaagaagat	atctgacaaa	900
tgacgttcaa	taaatatatt	tttaataaaa	aattagcctg	tggtagaat	tgaaatggag	960
aaagaaatag	aaggtagcca	ggttacctac	agctttgtat	atgacactat	agagctagga	1020
ctttattcca	tagatgaggg	aagtctgcta	aagtgtctac	attgaacca	ttattttgct	1080
agcattgtag	ttgatgtacc	taaaacagac	ttgagccggt	agagtaagta	ggcagacttg	1140
tccaagttag	aaaaagatga	aactgtgggtg	aggataaaga	gagaaaagga	gcagatttaa	1200
gaaatattaa	aacttgaaag	tactaagact	tgatgatgaa	ctagatgtgt	tagataagag	1260
atagcatgga	gtctagttaa	agttctgttt	ttctcactcg	tgtgactgcc	tcaataacac	1320
aaagcttgat	aggaaataaa	catgagatag	cacatggatc	tattacaagt	ttttgaaatt	1380
gagcttgaaa	agctacttca	aaaaataaat	tctaggccag	gtgtgagycc	atgcgcttga	1440
ttaaaaaaa	aaaaaaaaac	tcgta				1465

<210> 108

<211> 1265

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (766)

<223> n equals a,t,g, or c

<400> 108

ggggcagatg	gaaatgtctc	ggattttgat	aatgaagaag	aggacagtc	agtccctccc	60
aaagtggatg	agaatgacac	ccgtccagat	gtggagccac	cactgccatt	gcagatccaa	120
atagccatgg	acgtgatgga	acgctgcac	cacttggtgt	cagataaaaa	tctgcaaatc	180
cgctgaagg	tcttggtatg	gctggatctg	tgtgtggttg	ttcttcagtc	ccacaaaaac	240
cagctgcttc	ccttggtctc	tcaggcctgg	ccctcgctcg	ttcaccgact	cacacgggac	300
gcccccttg	cagtgccttag	agccttcaag	ttttacgtac	cctgggaagc	aagtgtgggtg	360
actttcttcg	cagccggttc	tgcaaagatg	tcctgccaaa	gctggctggc	tccctagtca	420
cccaggcccc	catcagtgcc	agggtgggac	cagtttactc	gcacacgctg	gccttcaagt	480
tgcagctggc	tgtcttacag	ggcctgggac	ccctctgtga	gagactggac	ctaggtgagg	540
gtgacctgaa	taaagtggct	gatgcctgct	tgatttacct	cagtgtcaaa	cagcccgtga	600
aattacaaga	ggctgccagg	agcgtcttcc	tccacttgat	gaagggtggac	ccagactcca	660
cctggttcct	cctgaacgag	ctttactgcc	ccgtgcagtt	cacacctccc	caccccagcc	720
tccaccctgt	gcagctgcas	ggggccagcg	ggcagcagaa	cccctnacac	gaccaacgtg	780
ctccagctgc	tcaaggagct	gcagtgaccc	tgctccccca	ccacagaggc	caccgatccc	840
tcccctactg	ccagccagaa	gctgggctga	ccccaccccg	gccataggcg	gtggcagcgg	900
cagcagagaa	ggtgaattag	ttagccaatc	gatttataaa	ttgatcgatc	acacaactgc	960
ttagaaatgg	attgaaggaa	agtagctgac	tattatttat	atttcatacc	ttgtgttttc	1020
aagtgcatt	gtctgggtgc	tctaagggtt	taacccttta	gcctaccatc	tctatagccc	1080
cagctccctc	acaggccaca	cacacacaca	cacaagaggt	cagttccctc	ccatctgcat	1140
acacctccct	gtcttcaaat	aatgagatgg	aactaatttg	ttttacctaa	cctgatcttt	1200
gggaaacaaa	cggaaataaa	gacacttctt	ggatgaaaag	taaaaaaaaa	aaaaaaaaac	1260
tcgag						1265

56

<210> 109
 <211> 1006
 <212> DNA
 <213> Homo sapiens

<400> 109
 ccacgcgtcc ggcaataatg acccattgtg gtttttaact tatctcatga aaagacttag 60
 gtttgttctc agggatatttc agatgactgc ctttataact ggggcacata cgattactaa 120
 ctatagtgat aggcgtttat acatttcccc tttagagccat ttctttatga acagtgggtc 180
 ttctgctcaa agtgttctgt ctcattctta tgtttctcaa atcttcttta aaaatgtaag 240
 caaatatttt taaagaattt ttatgttttc caaaattagg atttttagact ttagggattt 300
 tgatcttttg ggatttcaac attcgggatt atgggtgttca gtgtgtattt tggggggatt 360
 atgatcagca tcccatcacg tggaatatca tttggcaata aaaaggaatt aaatattgat 420
 tcatgctaca acatggtgaa cctaaaaaac attatgttca gtgaaagaag ccaaaccata 480
 aaggcctacg tactgtgtgg ttaaatggta aaaatgggtga atttatcaca atcaaaacca 540
 ataaacctct acaggaaaaa ataaggacaa agaaaggctg ttccctatta atggagacta 600
 aagagacatg acaactaaat gcagtttatg attgtggatt acatccttga tcaggggtgaa 660
 aatagctata aaggaaatta ttgggacaat ttgctaaaat ttgaaaatgg actggatttg 720
 gctggctcga gtggcttaca tctgtaatcc cagaactttg ggaggccaag gtgggtggat 780
 tacctgagggc caggagtcca agaccagcct ggccaacatg gcaaaacccg tctctactaa 840
 aaatacaaaa attagatggg cgtgggtgtg tgcattctga atctcagcta ctcgaggaggc 900
 tgaggcttga acccgggagg cagagggtgc agtgagctga gatcacgcta ctgcattcca 960
 gcctgggtga cagagcgaga ctccatctca aaaaaaaaaa aaaaaa 1006

<210> 110
 <211> 1258
 <212> DNA
 <213> Homo sapiens

<400> 110
 aaccctcact aaaggaaca aaagctggag ctccaccgcg gtggcgccg ctggctgacc 60
 ggctaaaaac taaaatgaca ttattccct agctacaaac atcagcggtta ttatgttaat 120
 tataccttgc cctctatcat tataaatggt tgccatggtg tttctaaaaa taagtgtttt 180
 accattaatg tgtagagggc aaacaaagca taaagtacta agggatcatg cttatcctag 240
 ggtctcacag aagagaggac atattttaatt aatcttgtga attacagaac aggttgtggg 300
 ccagacacca agaatcatag ggggtttttt ttaaaaaacc taatagaagt aggggtgacct 360
 ctctcttttg tctaagagtt ctaaaggaag gtaggcattc gtttaattag ttgggttcacc 420
 ctggctttac ctctggttta tgcttgtgtt aataggaagg aaaaatcact ttatcttttc 480
 ttccaagccc ctccctgcct gaactaccca gactgggatt accagatacc aggtgattta 540
 tgtggagatg atttttcacc tttaaactct aagccaagtg taagaaactc ttgatagcta 600
 tgtctatttt atatcagtea ctgagacttt tttttaagtt tttatttatt attaagacaa 660
 ctttgccaaa aaagtccct aagcacaact atttacattt ctttatagcc tcttctgac 720
 tctaacacat atgcagtttt aactgttatt ttcatagtaa ctgatctttt gtctaaggat 780
 ttttacctga aagcacaatg tattgagtct cttgaaaatc atctttcaga tctttttaca 840
 gaatgaactt atgcactgct actgtagtat tctcaaggaa tatatgtaaa cacaaatgta 900
 tgcctgaggt tgggttttgc agaaaacagt ctctgcttct aaaaacttct atgtctagtc 960
 ttccatagga aatcctcact gtttaaccat gtgaggagcc taagtcatta aacggatcat 1020
 gtctgtacat tgtgtaatga atgaaaagca cataaatgta atctactttg aactttgtaa 1080
 aaatgatgtg tggaggctat tcttgtttct ccatctcaag tctgtgtgt gcacgtgtgt 1140
 gcaagtgcac atgtgtgtgt gtaataacac attgtaaaga acagaaatta ctttaaaaaa 1200
 taaacagaaa tggagacctg aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaa 1258

<210> 111
 <211> 1453
 <212> DNA
 <213> Homo sapiens

57

<220>

<221> SITE

<222> (946)

<223> n equals a,t,g, or c

<400> 111

ggtcacatctt	ctcttgctcg	tacagagagg	agacacccgt	agaaatggag	atttctacta	60
cagatgaaaa	tttcttttat	aaaagggtaa	cttctctgta	ttatcctgtg	tttgctattt	120
ctgaaaataa	taagctgaaa	atattcctct	ttggcataag	gattatttgg	tgtggcatgt	180
tctgaacctc	cactgttggc	atcctttctt	gattagcaga	aacctaggaa	cattgtttgta	240
ataatgacta	aattattgtc	actgtcacat	ttgttagtaa	ctttttttaa	tataattgcc	300
attaaatgta	aaaagcagca	tctaagacat	tcaaaatgta	atttkgatac	tactttttaa	360
aataagatgc	taaattaata	gataagggtg	gtttcctcag	tatattttca	ttctaaacca	420
tcactaaaag	tagggctaaa	gaggaattta	gagtaggaag	acttaggttt	tgtattctgc	480
ctttgttcag	tatcagtgtg	actttggcca	agttacctga	cttctgaact	gcattttgct	540
tttctctaaa	taagtggggg	taatacctat	attagaggat	tatgataaaa	agatgtgaac	600
atattataaa	attattttat	aaactagaag	acatttcaaa	gaagttaagc	tgccactggt	660
agtttcacag	acttgggtgt	attagatgaa	cagcttttca	gttattgctt	ctatagtgtt	720
cctcttgccc	tttcttggtg	tatcagtttc	tgctgtctca	cctagtcatt	cccatcagtg	780
taaaacattt	ataytggtat	ttcttccaag	ttcagaaaaa	accctctyty	gaytcccccc	840
atccccattcc	agcacttttg	gaggccaagg	cgggcagatc	atgaggtcag	gagatcgagm	900
ccatytctgg	taacatgggt	acccccatct	ctactaaaaa	tacaanacaa	attagccggg	960
cttggtggtg	ggcgctgtga	atcccagcta	ccggggaggc	tgaggcagga	gaaaggcatg	1020
aaccaggag	gcagagcttg	cagtgcagcca	agattgcgcc	attgcactcc	agcctgggag	1080
acagagtgc	actccatctc	aaaaaaamga	awaaaaaaa	caacttattt	taaattattt	1140
tcctagaaat	tatgatgtca	gcagaggtag	ctaggtggtg	ttatgggtga	cttttggtat	1200
ttttaagaca	gcttccgtat	ttcttaggag	ttttgctgaa	gaacatggta	tggggagaaac	1260
atataatatt	ctcatacact	tcttaggatg	ggatagatcc	ctgtaacaga	atattgggtta	1320
acaagagaaa	aacaagtttt	aagacatgta	tacctcatat	atacatggga	gatactcggg	1380
ggaagtgcgt	aaatctctca	gaggtggctt	aaataccatc	atgtcctgaa	aaaaaaaaaa	1440
aaagggcggc	cgc					1453

<210> 112

<211> 1552

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (1035)

<223> n equals a,t,g, or c

<400> 112

ttgcctaagg	cccactgtgc	caaattagat	aatacaagaa	gttcattttac	actgtagacc	60
agtgcagtc	atgactgttt	gctctgtgat	accgtttcaa	aatccaaaa	tcgagacttt	120
tctctgtgcc	atgcaggatg	cagctgtgtg	tgatatgggt	tacagtaata	tttctttctc	180
aaagtagcag	gcttggttaa	gaaaagataa	gcaacacatc	tggggaaaag	ggcaggtggc	240
cagcaatcga	tgtggtagct	ctttgcccc	ctcggacagc	aggaattagc	ttccccaggc	300
attttctgta	tgtgagttgt	attgtgggat	gtacaaatat	catctgttcc	tttgggtttc	360
caggccagta	gctctctatt	ttgggttcaa	acatgggttc	tcaggccggg	cgcggtggct	420
cacgcgtgta	atccccggc	tttgggaggc	caaggcgggc	ggatcacgag	gtcgggagat	480
ggagaccatc	ctggctaaca	tgggtgaaacc	ccaactctac	taaaaataca	aaaaattagg	540
caggcatggt	ggcgggtgcc	tgtgktcccc	gctactcagg	aggctgaggc	aggagaatgg	600
tgtggaccgg	ggaggttggg	ggttgacagta	agccgagatt	gcaccactgc	mctccagcct	660
gggcaacaga	gcgagactcc	atctcaaaaa	acaaacaaac	aaacaaacaa	aaacatgggt	720
tctcaaaagg	catgcccact	gtctcccatg	gagcttgaca	gcccattgcca	ttagctctca	780

58

ctgttaggtt	tctggggaag	gttctttctac	ttgattggaa	aatttccaaa	taaatctttc	840
cagaagatac	tatgcacaca	gctaagtggc	ctgtctgtgg	agtaaccctt	ttgtaaacia	900
acagaaacct	aaagcttgat	gttttggggg	gctgcctgtc	atctataggt	tcatttaggt	960
gtatttagga	agaggatcca	tgaaccact	ggtttctgt	tacataataa	tcattaataa	1020
tgatttaaaa	tgtgnacatt	gatttttttr	aattccraaa	tacaagcgta	tatggtawat	1080
taagtcaaat	ggtatgttca	gtgagcgaga	tggggcttgg	ggcaaaacia	tactttgctt	1140
ccaaagagga	tacaactctc	aaggagattc	tttcatcttg	cctttaaggt	catttaaact	1200
aattcacata	atcttcagaa	aactaattca	catcatctat	tcagtgtgaa	aatcaaaagg	1260
aagactgttt	tcttagtctc	tcgttgccca	actggccatt	tataactacta	ggttgattaa	1320
gggatttgcc	tttttctgct	gatatgggaa	caaaaagtct	taagcatttt	taaaggcaat	1380
ggaaaattca	gccacatggg	ggaaaattga	tattgtcacc	attgagttgc	tctgtttctt	1440
ggtgaagagt	gaatctaate	tgatttctct	cttcatcaga	tatgcctctt	taacaacaaa	1500
aaaaaaaaaa	aaggaattcg	atatcaagct	tatcgatacc	gtcgacctcg	ta	1552

<210> 113

<211> 1489

<212> DNA

<213> Homo sapiens

<400> 113

gaattcgcca	cgagtgccca	gtcctgtctg	taattagctc	cacgtgtacc	cccttcattc	60
cctccctccc	accgagccat	ccctgaccca	ggaactttcc	gcagactcgc	cgccatctgg	120
gagtgaagca	acatggatgc	agtcagccaa	gtcccatggg	aagtcgtgct	tccaagcac	180
atcctggata	tctgggttat	tgtcctcctc	atcctggcca	ccattgtcat	catgacctcg	240
ttgttgctgt	gcccagccac	tgcagtaate	atctatcgca	tgcggactca	tccgatcctt	300
agtggggctg	tttgagagcc	tccaagagg	gccgggtgag	ggatgaggac	aggcacctta	360
tccccagcct	cttctgtctt	tcagaaaagc	agcaggaggg	actttggggc	atggacctga	420
gttctggttt	tgattctgcc	acgagccagc	tgtgtgaatt	tggtaagggg	acctaactct	480
ctgagttcca	ggttccttat	ctttcaaatg	gggatggtga	tccctgccct	ttctacctca	540
tagggatgtg	agaaccacct	gacttagtgg	atgtgaaagc	tgtttgtgat	cagtaaagct	600
accacagata	taagggtgtt	atgctgaate	ctgagaagct	ttcaagaacc	agagaacctg	660
attgctgatg	atggccttaa	aggtggtgag	ggagatactg	ggggcagagc	agactttgcc	720
agtgcacctc	aggtcaaacc	aagccaagag	caccctgtcc	ccattccaag	gggccagcag	780
cactttggcc	caaagtattt	tctttaaggt	gccattcctt	catgttttct	cagtttggag	840
ggtgatgggt	agagctttcc	agaaccttct	ccattccaga	atctctgccc	ctgtgtaate	900
tgaaggaagg	ctgtgccatc	tttgggcact	gccaaggagg	ttgggggtgat	gggcttcttt	960
ctgcactgga	gtctcacatc	tgtagctttt	gacactcaag	caatgttgga	aaatgcaggg	1020
tgactgagtt	ccctgcccag	ctttcgggat	ctctggcccc	catccccctg	tgtgtgtccc	1080
tctgcccagc	tctgtctgta	attagctcca	cgtgtacccc	cttcaactcc	tcccaccagc	1140
tctgcagcca	gcctatggca	attatatttt	aagaggtgtt	cccaggactt	ttggggacct	1200
ctaaaacaat	gatggttatt	ttagatgtga	tgatttatat	ttatgtagag	atatttctgg	1260
accactcaag	ctcttcgata	ccaaaatcag	gagcatcttg	ggatttatta	aattatgtaa	1320
gaagatagca	cagatatcgg	gatattattg	tgtgaaaatg	ctgctttttac	tttgatgtga	1380
tctcattgat	gtacacaacc	aagttccaat	aaagtgtctg	aatgtgaaaa	aaaaaaaaaa	1440
aaaactgcga	ggggggggacc	cgtaacccta	atcgacctta	atgagtgtga		1489

<210> 114

<211> 607

<212> DNA

<213> Homo sapiens

<400> 114

ggcacgagtt	tcaacttgag	atttggaggg	gacagacatc	caaaccgtat	cattaaattt	60
aatagtttta	tgcagttttt	ttggctctag	atctgttttag	actcctgcag	tcagggtgtc	120
gtaactagcc	tctggctcct	tttgagagtt	cacagtttgg	tgcaaaccct	ttggatgtat	180
tatttgggaa	aatgggatat	ctggcagcct	gtgtccctgc	tttacattat	ccttttttgc	240

59

gcctgcccc	gcctcctcat	tagcatccct	gccaaaggcca	gtggagaagg	atggagatgc	300
ggtgacattc	agctgacagt	tgtcacagat	tgataatagc	taacagcaca	tctctcccc	360
ggctccttcc	ctagtgcacc	aattagccca	gcctcatctg	cacctgggac	tcaagttgcc	420
taaacatatt	tcatttccca	tagcagaaga	tgccatccat	ctagagtgag	actgaaaata	480
caaacaattc	agaagttgtg	actttccatg	ctctgcacac	agaggctacc	aaatgctaag	540
ggcgcttcct	ccccagcacc	aggettattg	ttctaagctc	cagaaaaata	tcaaataaac	600
cctgccc						607

<210> 115

<211> 1498

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (791)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (895)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (915)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (936)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (1017)

<223> n equals a,t,g, or c

<400> 115

gcatccccat	ttctgttcc	ggaccctctt	ctgtcacccc	atcgacaagt	aaagaaatca	60
agaaatccaa	actgattcga	agccagtctt	ttaataatca	agcttttcat	gcaaaatgatg	120
gcaacttaga	gaaatgtgct	agtaaaagtt	ctacaacagg	atatacaact	tctgtctcag	180
ggttaggaaa	gacttctgtg	ctttcactag	ctgatgattc	attccggact	cgtaatgcc	240
gtagcgttcc	atcttccttt	tctcctaata	ctcccttacc	gagtacttcc	cgtgggacag	300
gtaactcagt	tgaccccaag	agcagtggaa	gtaaagatac	acaaccacgg	aaggctacct	360
taaaatccag	aaaatccaat	ccttaaatca	actgcttgat	gaaggaggca	aaacaaaggc	420
agcaggagat	aatgtgattg	gtacacaaaa	gctaaagcag	tggctggggc	tttgttttta	480
aattttgggt	tttttttttg	ttgttggtta	tgagcagaaa	gagagacata	atgacagctg	540
atgtttaaact	tttcatattt	caaattagat	tccctaggag	gtataatata	tatttcttga	600
gtaataatgt	ggttacggaa	ttccaatggt	atagtgaagt	gtaatgaaaa	acatctctag	660
gaatgtgctt	taaccactgc	tgcaaaagar	acaagtctgc	atttatttgt	gcaggaaacc	720
accatttaat	tgttctagar	tttttagcatt	taaaaatcgt	atgaaagtct	acatcagctg	780
aattgtccta	ncttgataag	cacttggagg	gggacttgga	aggtgagaaa	gatactgcat	840
tttctcatga	gtctccaagc	ccacttgaaa	agtcacactg	aaaggatgca	aatancggc	900
tgcgattttg	gcggnccctc	cggattgtgg	tgacantatg	ttgtaccggg	acattcccaa	960
atttaaattt	ggcagtgraa	attccacaaa	gatttctggg	aacttkgagc	tataatncct	1020
taaaaataat	mgattgatga	ttttctttat	ttccagagaa	tttattttat	aaatactttg	1080

60

tttacatttt	agattgtact	tgtcttttatt	taaatgatga	atctagtctg	aatcagctgt	1140
tattccagct	atcatgetta	agctatgtca	acagcattta	tttgtactaa	tgctaataatt	1200
tccatcaaaa	tttgectgtg	gaatatatac	agttcttaaat	tttaaactgt	cagttcttga	1260
gatataccgt	gtgaagctat	tgtcagcttc	tctatttcaa	aatgcaatca	gcatataact	1320
atattgatat	tagaagatat	ttgtttttta	aaatatattg	atgtatgata	aagatgatct	1380
aatttgtgaa	tgcatatgta	tgtgtgggta	ctttttataa	tgtgaaataa	tgaataatga	1440
atttactcaa	aataaaacat	aggttaatga	gaaaaaaaaa	aaaaaaaaaa	aactcgaa	1498

<210> 116

<211> 1797

<212> DNA

<213> Homo sapiens

<400> 116

ggaacacaaa	aaaaacactg	ggaaaaagaa	atgaaatctt	agcagaagct	ggaacttttt	60
atacttataa	aaatagtatg	ttcttatggt	tctccttatg	catttatgtg	ttcacttaaa	120
aatgttttta	aatctaaggt	tttctttgtt	tatgttcagg	taaggaaactg	ttgtcatgat	180
ctggaaatgt	ttaaaacaat	gtttgttagc	attctgtgag	cagcaaaaact	tatagtgata	240
aaaatcgatt	gttgtaata	tgatgtttac	catgtgcaca	tagtaatgaa	aaggaacata	300
aaagcccagc	aggctcgtac	caaagtcaca	gcagtaatgc	tatgtactgc	agagtctgat	360
gagctggcct	ttgtgcacac	ttttattttc	atgggattgc	atcttagctg	ttaaaacttc	420
tagattgaaa	tttgacagcc	agggttacat	attggggact	tttaaagtgt	ctttccaaag	480
agatttcatt	aaccgttttag	attagaatat	ctttcccaat	tggtacagtg	acatatatgc	540
tgcaatattt	aacaactgga	gtattagcca	catgggttat	tttttcaatc	tgtgttttga	600
atttttttat	tgtgtgttat	ttaaaatatt	acatatgcag	ctggggagaac	tacacctttg	660
tgcacataga	tttatatatt	aatttgtaga	aaatattttc	tttatatatt	tccttaccat	720
acaaggtgcc	ttgttcatca	ggaaaacttt	tgttttgtat	tttgacaaga	aaggcacctt	780
cagagtttct	ttttaagtat	agttgacaag	tgtataaatg	ttacacttac	tttcagagtt	840
cttttttagat	ctaaagaagt	cagttcaaaa	atggaaatca	acaatgttag	gagaaatctg	900
aattctgtta	agttagttaag	tattatgtat	agcatctgtt	tttaaccatt	tccattctta	960
tccttagtgt	atcagttgat	cacactaaga	aagcttaaaag	attgagcatt	tgaataaaat	1020
gctctttata	aatgattaga	tttttgaagg	gatattgaaa	tcattgcgct	gtgatttcat	1080
ctgtgatgtg	aaaaatcaat	ttattatcct	tgggtgcttc	ccccccacca	atgcacaaat	1140
aattgtgaac	agcttgaaat	gacttaaaact	gtaatccaaa	tgggacaatc	tgataaagaat	1200
ttcatgcatt	ggtagttaaa	taacttaaat	tgctaaagct	ttagcttgaa	atttatgttt	1260
agaaaaacta	ttgatttccc	atgggtatgaa	tataactatt	gtaattcttc	aaatgagact	1320
cttctcacct	taaatagtca	tatattaatt	aacttatagg	aaataagcat	actatatgtt	1380
agctgtttta	aaaggtacca	gatgtaagag	tcataaatat	atgcaattaa	agaagttcat	1440
agatttcaca	tgaatgtaaa	tgtgttatat	ggagacatgt	cttgtaaaca	gttgaatgta	1500
tgtaagtttt	ctgtttgtga	aaatgtagtt	aatgtactca	ctgtggaggt	cataaggaag	1560
ctactttttt	tttaaagtgg	aacctaatta	aaatatttcc	agaatcaaag	agacttaaat	1620
ggtaaatttt	taaaattttc	ttatctctta	cttttttagtt	ttcaaagtag	aaaaaatcag	1680
gaattttttt	attaactagt	acttacatat	taaataaaat	ttattattgg	ctaaaaaaa	1740
aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aactagttct	agatcgcggg	cggccgc	1797

<210> 117

<211> 952

<212> DNA

<213> Homo sapiens

<400> 117

ggcacgagcc	tggcacatag	aaggtgccca	ttaattggat	gaacgaatgg	gggagttaga	60
gtcagtacta	gaaaaagcat	gggattatac	ttttaagagg	ccaaatgaaa	atgtggggaa	120
tagccatttt	ccaggaattt	aaaaggagcc	attttttaaa	tgtcaataat	aatttattgg	180
ttattatttt	ttaagcactt	attatgggtg	cttattatag	gcatggtaaa	agcacttcat	240
ccacattatt	taaatctcag	aatctatgag	tttggtgaga	tcactgcagt	tttacagagg	300

H

aaaaaacagg	gcagagagaa	cggttaatttc	ctcaagttct	cactcttgtc	acttaataga	360
tctagaattc	caaccagat	ctgatggtga	agtcagtata	agcttcctgg	agaaaggagt	420
agagttaagg	tgggggatgg	ggcttgaaga	cttgatagga	ttaggggttg	gagtgtcaac	480
tcggagatcc	acagtcaggc	ggaaggaacc	cacaaaggca	ggaatgcaca	cagcatgctc	540
agaggagatg	gaacctgaaa	atagagagaa	ttagcaagta	gtgcccattg	gatggagtaa	600
agagccatac	ctagaagggt	atcactgggt	aaccatgata	agagttttgg	cctgggtgctg	660
tggctcacgc	ctgtaatccc	agcactttgg	gaggccaagg	caggaggatc	acctgagggtc	720
aggagttcga	gaccagcctg	gccaacatga	tgaaccacca	tctgtactaa	aactacgaaa	780
attatctggg	tgtggcgga	ggcacctgta	atcccaccta	ctcggagggtg	acgcaggggg	840
aattgcttga	accggggagg	cagaggtggc	agtgagccac	gatggtgcca	ctgcactcta	900
gcctgggcga	cagagcgaga	tccgtctcaa	aaaaaaaaa	aaaaaactcg	ag	952

<210> 118

<211> 1185

<212> DNA

<213> Homo sapiens

<400> 118

ggcacgagct	tatgtctatg	ttttgagata	gaatccttct	attttatagc	agggatgggc	60
aaaccacaag	caaggggcca	aatccagcct	gctgcctggt	tttgctaaaa	aagttttatt	120
ggaacacagc	catggccatt	cacttccata	tcattccaatg	gctgcttttg	tgctacaatt	180
gccaccatgc	ccagtggggc	ctgtggcaca	caactgcaga	agtgagtggg	tgtggcagaa	240
atcacttagc	cttcaaagcc	taaagcactt	actatctgac	cctttccaga	aaaagtgtgc	300
tgacctctgt	tttagagata	caaactaaca	tacttacaga	taagtgatct	gaatatgaag	360
caaattttca	acataatctg	ggttgatggg	gaaccgtggt	acataataac	aatccagag	420
ctaatacttg	ccaggtgagg	gctgtgcaga	cgatcacgac	actattctct	tttgcataatg	480
ttttccataa	taaaaagtct	tttgaaaaaa	tcaatgattt	aggggtttta	atatactgta	540
gtcacttttg	aaaacagttt	ggcagttcct	caaaagttag	acatagaatt	accctatgac	600
ccagcagttc	cctttctaga	tataccccc	agagaattaa	aaacacacat	tcacacaaaa	660
acctgcacac	aaatgttcac	aggagcattg	ccttaatagc	aaaacaagcg	aaacaaccca	720
aatgcccata	aggtgacggg	tggataaaca	cagtgcgggtc	tgtccataca	tggaaatgtga	780
ctcagccaca	ggcakgaatg	aagcgtgac	acacgcagca	acacggatga	accgtgaggt	840
cacggagtgg	agtgaagat	accagtcag	aagtcacaca	ccgtatgatg	ccattgacat	900
gaagtgttca	gagcaagtaa	atccttacag	atggaaggca	gagcgggtgac	tgccagggac	960
taggagtggg	ggggcagggg	gtgactgcta	atgggtatgg	gatttcattt	cggggctggg	1020
ggaaacgttc	cggagccaga	gagtagtgat	agctgcacaa	ctctatgtat	atgctatgaa	1080
tcaccaccga	atggtatatt	tttaaaggac	gaatttatgg	tatgtaaatt	gtgtctcaat	1140
aaagctgcta	tcttaaaatt	caaaaaaaaa	aaaaaaaact	cgtag		1185

<210> 119

<211> 1098

<212> DNA

<213> Homo sapiens

<400> 119

tcgacccacg	cgtccgcaga	actctagatc	tgggaataac	cctgtatcta	tttctttaca	60
tttttctttc	attaatgtat	ttatcactat	tttttttttg	tttttctttg	caggcctcag	120
ctgttgaaga	acgtcttgca	gagagcagta	gagagggggc	agttagaaca	gataactggc	180
aaaggtgctt	cggggacatt	ccaggaactc	cgaccacact	ggtgcagagg	agtcttgtct	240
tgacctgctt	tgggagagtg	ttgtcttgat	gcttttctct	tcctccttgg	aaaacagctg	300
aagaaatcag	gggagaaacc	cctgcttggt	ggaagcctga	tggaaatgac	aatctgtct	360
gccattgctg	ccatgaatga	gccgaagacc	tgctctacca	ctgctctgaa	gaagtatgtc	420
ctagagaatc	accaggaac	caattctaac	tatcaaatgc	atttgctgaa	aaaaaccctg	480
cagaaatgcg	aaaagaatgg	gtggatggaa	cagatctctg	ggaaagggtt	cagtggcacc	540
ttccagctct	gttttcccta	ttatcccagc	ccaggagtgc	tgtttccgaa	gaaagagcca	600
gatgattcta	gagatgagga	tgaagatgaa	gatgagtcac	cagaagaaga	ctctgaggat	660

62

gaagagccgc	cacctaagag	aagggtgcag	aagaaaacc	cagccaagtc	cccaggggaag	720
gccgcatctg	tgaagcagag	aggggtccaaa	cctgcaccta	aagtctcagc	tgcccagcgg	780
gggaaagcta	ggcccttgcc	taagaaagca	cctcctaagg	ccaaaacgcc	tgccaagaag	840
accagaccct	catccacagt	catcaagaaa	cctagtgggtg	gtcctcctcaa	gaagcctgca	900
accagtgcaa	gaaaggaagt	aaaattgccc	ggcaagggca	aatccaccat	gaagaagtct	960
ttcagagtga	aaaagtaaat	tttataggaa	aaaaggggat	catgatgaaa	ttcaaaatct	1020
tattttctaa	gcacttttga	tatcaagcaa	gtggcttctc	ttttgagata	ttaaaaaaa	1080
aaaaaaaaag	gcggccgc					1098

<210> 120

<211> 805

<212> DNA

<213> Homo sapiens

<400> 120

ggcagcaggt	ccctaattgt	cttgtaccta	gccctagggg	gaccagggca	ggggaatcat	60
ggcgagaagc	gtaagggcct	gatgaagaag	gtgtgctggg	tgtgggctct	agcccacttg	120
gttttgtgtg	agaggtggct	gacagcaggt	tgtttctgtg	atgtaggagt	tatccagccc	180
tgcaagggca	gtccctccag	tgtctgcaaa	gcccgaagat	gtctgcatcc	aaaatacaga	240
ataaaaagat	atggttacta	caagtactca	gtaagactga	taatctgtca	tcatcatcct	300
catgccctta	aagcagagct	aactgatgat	taatatatgc	ttctatgtta	acagtcttgg	360
actttattaa	tggtgggtgg	aagttaactt	aatgtatgta	tgcaaactaa	aaagtggcat	420
ccttttcatt	aatgacccaa	ccattattca	agagctatgt	ctagttaggg	acttcagact	480
tttgaaagaa	atgaagaaat	aatgccagat	acatgggctc	gcacttgga	tcccagctac	540
ttggggggacc	gaggtgggag	gaccgcttga	gcccaggagt	tcgagaccag	cctgggcaac	600
atagcgaaac	cctgcctcag	ttttaaaaaa	gaaaaaaaga	agtagtgaag	aaattggaaa	660
ggattctgag	aagaaatatg	caaggtggaa	aagagcctag	aaagaaaggt	gacagatgct	720
gggatttggg	cgtcagaaga	gatatctagg	aaatagcatg	gcagccctca	agtactagct	780
ccacttaaaa	aaaaaaaaaa	aaaaa				805

<210> 121

<211> 1598

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (1067)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (1069)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (1577)

<223> n equals a,t,g, or c

<400> 121

aggaaagaac	aaaggttatt	tcctggagaa	aagacaattt	attcaacacc	aacragggac	60
tcatcatatg	ggcacaactc	tggtgtcctt	ctatggagaa	aacctcaagt	aaagttttat	120
tctgcctttr	aaaatgcttc	caaaagtaga	ccctgtcccc	acacaggtca	agactacaga	180
gaaggctttg	tagaaatgtg	tcacctatgt	acacctgcta	cttacacatt	tcctcttttg	240
gaaaaatgag	atacttagaa	taacargaaa	attaagacat	actggcctgg	tgccagcaga	300

63

tggtcttttct	atagacaaac	taggttagtg	tggaagatat	aggttaaaat	aaactatgct	360
gttttatttta	tcttcccaac	ctgattggca	gctagacttt	tttagggctt	catttaatgg	420
ccctgttttt	ttcattatta	tatttaatga	tagggcagga	tttcgtatgc	aagctcttgt	480
ttctcaggct	gcctgcagaa	gaagtcgcta	taaattatct	gttgtctaca	tgggtacaagg	540
cccattgact	catctgatgc	ttgttttgtt	aattttctta	atatttttat	cacggggcag	600
tgggagggct	tgggctttta	gccacagctg	ttttaagact	tctgatctcc	tgccttgcag	660
gaataggttg	gaagtcattg	aattttttaca	ctatagtaat	ttgcattccc	acataagttt	720
gagtgttacg	aaaacattcc	tttaaaggga	tctgtgctac	acaaaatatg	ccaggacctc	780
acagacaaaag	ccattgctag	aaatgtcatt	ccaatgatca	gatctggaaa	caggctgcc	840
taaccacttt	tccttcttgt	agactcagct	cacctgtata	tttaaactgt	tcttggcatc	900
ttgaaacacc	tattttctact	cagggtactca	ttgtcctgtt	actgattcac	ctttctgac	960
cttttcaacc	agttttcccc	caagggggga	aatttttactt	aacctctagt	atttgaacaa	1020
ctcaatattt	gaattgttgc	cccatttgct	tttacctgta	ctgtatncnt	ggatcatctca	1080
aatggcgctc	aaaccagct	actttgcatt	ccagaagttt	ccattccctc	caattccacc	1140
taatttttca	tctgtcctag	ttactggctc	tttcttcatg	tcttatttct	cttgcttttg	1200
gagcttaaaa	gatttttaca	gacctaat	tggtttcctt	ccttggagcc	atagttaccc	1260
tgccaagaag	agtagaaaat	gggttcaact	cctgtttcgc	tccaccaaca	cctctgtgag	1320
tctcatcatc	agctgagcga	tgatgcctta	cagggtgcat	agcactggaa	ctttcctaga	1380
gtaacggctc	tgctgccagg	gtttctctgg	gctcattctt	ccactgactt	aattatgatc	1440
tatgcctaac	agagcccag	tacaactatt	ttgcagaatg	gctgttacc	tagaattact	1500
atagcacata	ttgagatata	gttgactctc	ctagtagata	ggaactgacc	ccaacaataa	1560
actttgataa	taaaganaaa	aaaaaaaaaa	actcgtag			1598

<210> 122

<211> 1020

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (3)

<223> n equals a,t,g, or c

<400> 122

cgncacgagg	tgaagttcaa	cccaatgcaa	ctttccttca	gtctttccgt	gaaagcgctt	60
gtgaaaaatg	aggatcatatt	tccctttctc	agtctgcccc	ttcccgtttt	gctctccggt	120
tttcttcttt	gtcttcacag	atgtttacct	atgttttttc	tttgtttttg	ctgttggaag	180
acatctaagt	gacccctttc	ccattctctt	tttactcat	aaatgtcctg	atgttttagca	240
aaaggcagtt	ctcttttgcta	cttgagcttg	taaactgttg	ttaaatgagt	aaccaaaagg	300
aaagtccctg	cgaagttggg	taccatttca	gatacaagaa	ccgtttatct	tcccacgctg	360
acgaattttg	cgagtggagt	gattattttt	ccttggtgtt	gtaatttatt	taagtaaatt	420
ccttgtttgt	ttttcttttc	agtacaccag	gggtatatat	tttcaatatg	acatgtacct	480
ttggttcagg	gctaagttag	agtctgaaaa	atgaagcctg	taggattcat	ggcagtgatc	540
taattgtgat	tcatcttact	gattgtaggg	caagaagagt	ggactaactc	aagacacaag	600
gcaccttcag	cgaggacagc	aaagggcgct	tacagagacc	agccatatgg	cagatactga	660
ttgtactgtc	tgatgttgtg	aaatagccaa	tctccaccag	tctgtatatac	tgttcaaagt	720
aatttttttc	tatgaacaat	ccctttttta	ataaatcaaa	atgcttaaaa	tctgaatgga	780
tggaaactta	aactactttg	ttgaaacatc	aacctgggca	gaaaaaaaaa	aaaaaaagac	840
atgtaaaatt	ttgttatttc	cagtctgtat	atgaaaaaat	aggatcatca	aaggaaaaaa	900
aataactttg	attaactagt	gttaaacaaa	aaatagggtt	actaaatctc	gtgccgaatt	960
cgatatcaag	cttatcgata	ccgtcgacct	cgtagggggg	gcccgtaccc	aatcgccctgt	1020

<210> 123

<211> 1378

<212> DNA

<213> Homo sapiens

64

<400> 123

atcttagcaa	tctccttggc	ccaaaacttc	accccatctt	ggaagggagg	ggagagagaa	60
tgttctgac	tatatctgat	gagggcgtgt	ggttgggacc	tgagcatcct	cctgggtggg	120
ctagtgatgg	ggagagaggg	ctgttactca	cgactccctc	caacagaata	ccagaaacag	180
gcaggcagct	caggtgtatg	taaggatgtg	aggccaagaa	accagccctc	accaagttac	240
ccctgtaaat	ccttgtctcc	ccatgcacct	ctacttttag	tcagaaatgg	attcattgca	300
ggctcagttg	tttgtattat	gtgaatgaac	tgaacgtaac	caagcaccaa	gagagcccta	360
aagacacagt	agacctcctg	tagaagggct	ctgatggacc	ttcaaacatt	gctyctccaa	420
ctttatgggtg	cacacaaatc	acctgtgcat	gttaaaatgc	agacggtgac	ttacagatct	480
tggggggaggc	caagtgtctg	cattttccaac	aagctcccag	atgatggcca	cactgctgat	540
ctgaggacca	cattttgaac	agcaatcctt	aaaacacaca	aggctgtggg	accgcactcc	600
tgagaagaga	cctcaytctt	ggaagggatt	ytaggaggct	gcattcaaac	catcccagag	660
acagctcaca	attttgttag	ggacctgata	ctaaaaatta	ccaaatcaca	atcgtgtgct	720
gagggccttac	tttgtgcttt	tcagatgaca	tttcatttga	tgctctcaat	aaccctgaag	780
ggatttcaga	gatggggatt	atgatggagc	ttcagaatta	aactgattaa	aactggccca	840
gagtcacaca	actatgaaga	ggctgagttg	ggattttctt	taatttttat	tttttgaaac	900
agggctctcac	tatgttgccc	agactggtct	cgaactcttg	agcttgagca	atcctcctgc	960
ctctgcccc	gagtagctgg	gactacagat	gtgagacagc	atgccgagcc	tgctgagttg	1020
ggatttgaac	cccaggtttg	tctggctgta	agtcctccag	tgtggaagta	cgatagctac	1080
ccatctcagc	cacggaaaac	agcagagttt	tggttcaggc	aagtagcact	aagtagaaag	1140
cccaggatg	aaccagatct	gcaggagccc	aaagtcattg	ctcagaacca	ccccctaccc	1200
gctgagaagt	tcttctctgc	cctggaagtt	ctctctgaga	tccacccaaa	tgtgtcctca	1260
ctaaggcgaa	atctggcttc	tctgttttgg	gcctggggag	aaacataaac	ctagcccttt	1320
gtacacaccg	cctcgtgccg	aattcgatat	caagcttata	gataccgtcg	acctcgta	1378

<210> 124

<211> 1146

<212> DNA

<213> Homo sapiens

<400> 124

tcttttctgt	aacttcattt	agcactcaaa	acaatatcac	tatttaatgt	tacactttga	60
ctacaaactt	atctttctcc	ttacaagctt	tttatgtatt	taatattatc	ttggctattt	120
ctgtgcaaac	tagttaaatg	ttactttgaa	attcttcttt	tctctacatc	acctcagtta	180
ctccagtgga	cagtaattgt	tacttactgt	ggctctctgt	tacggttttg	aatttcttca	240
aatgcctttt	gaatttaatg	aaaatactac	atgaaataat	actggtggct	acataatttt	300
cttccacttt	ttcttaagtc	tctgcaatga	aacagctgac	agtaagggtg	gcgtgagtg	360
tgagatatgt	taactcctat	acttcacttt	agcttatgtt	gtcagggagt	gatctccaaa	420
tgcaaaaata	taaagtactc	tattcttggg	gaacaatgtg	agcaatggaa	ggcttggcct	480
ttcttaggca	gatccctcag	attcaaaagg	aacacatctt	gatttaatat	caatctctat	540
acaggggcct	gagaatggga	tgaggagagc	tgctggctct	tgtactgcaa	tttaattcac	600
tctgattact	ttctatcctc	ccacccccac	tcttctctca	gtctctttct	gattagttat	660
catagtcttc	ccaagactc	tatatatgct	ctctgggttc	ctgtcaagag	gaaatcctgt	720
acacaccccc	tactctctta	tactgtcttc	ctggctctgg	cactctcact	gtgcagcttc	780
ttctgtctctg	gtttagccaa	caatataatc	catagatttt	ctgaattcca	tcatagcatc	840
tggtctttgt	gtgagtcttc	tttcattccg	tgtctattac	gggtattttt	aaatatcttt	900
actgttgata	cagtggagatt	tgtgggaaag	aggatagaaa	cataagccaa	tctccccctc	960
tgaaatggac	atctattatt	aattaacact	catcaaaaata	aatgttctta	ttgttatatc	1020
ttagttggat	aaagatgaaa	agataagacc	aaaatcctgc	tctctaatac	cgatttattg	1080
taacaataat	attattgtat	tattgggggg	ggcccgtagc	caatcgcttc	acatgcatcg	1140
tataca						1146

<210> 125

<211> 1675

<212> DNA

<213> Homo sapiens

65

<400> 125

cttaaagacg	ccaggtagag	acacacagaa	cgtatgtatt	aagaatatcc	tctctgggct	60
ctgaaatfff	aggagtgtt	cttatccact	ccaagttgta	agtatttgta	gaaatttggt	120
caaacaacaa	aaaactatca	aatgaaaaga	aaatgtactc	aacctactt	atagtttagca	180
gctggaattc	tcaactcttc	cctgccagca	ctataccaca	gtgtggaaga	aattagtcaa	240
atgcttgfff	tectgtctct	cttttcaact	gttactgtgc	tttgtttgaa	agtagttttc	300
tctctcaaa	ccgttgctta	tatcgtttaag	aatgaagggt	tgtgtttaaa	atttattgca	360
ttgcaaagg	tagtttctact	gaagtcctgc	accattaaat	aagatgaaat	atttgtatff	420
attgtcctac	ttcctaagcc	gtaacttctt	ttcctctgtg	aatttgcatf	gagtcactca	480
tgctacacta	catcgcttta	gtatttgaga	tggcatttat	gtttcctctc	gtttatcatg	540
aaatgggggc	agattccatc	agattccacc	tctgtcagg	ggactcttgt	ctgccttcca	600
tgatgagatt	ttttttctcc	ttcccctttc	tttaagagag	gctgacagat	ctaggtgtca	660
atcaattgga	aaccagtctc	tgattttttt	tcattagtta	ttttctatca	ttagtttcac	720
tgtgtaaatt	agatatcaac	tgcacttctt	taaaaaaaa	tacatctccc	tattacctcc	780
ttgaaagatt	tacttctgta	ggcctttttc	aataggctca	tgactgcaga	caaggaaaaa	840
aaaagtaaaa	acaaaaacag	tatgtgcctg	aaaatgacaa	aaaaaaaaat	tgtaacattt	900
aaaaaagaaa	cctgaatagc	ctttaattct	ttaataatac	acttaaattt	tatgtaaatc	960
ggttttcgcc	acgtgtgttt	gttcacattc	taaatgactt	aatgggattc	tcacggctctg	1020
tgtctttgtg	tcacgtgtat	aaaatgggct	tgtgatgtaa	gcgtttctatc	tggtcagtgg	1080
ttcctttgat	attgtactgc	tgctgggagt	gggctgtgga	acctgccttc	gggtaactgg	1140
gttcctcttg	ggtagattgg	agagatgggg	gtgggcgtgg	gcaaattctc	acacatgttt	1200
tcttaaccta	tttgagaaaa	ctttcaaaag	gcatttgatt	aaacctcttg	gcagtacagt	1260
attcttgtat	ttgttaacgt	ctgtgttttag	gtactggtac	ctttttgttt	taaaatgttc	1320
taagtgttgg	ctttaaagt	aatttatctt	tagtatgata	gttatatgaa	aattatagga	1380
tttgtgtgca	gagaattttt	ttataaagt	ctttgtaaaa	aaaaaaaaat	gtattctagc	1440
ttttgcggta	catatgtgtg	ataactttta	tacctatgac	agttaagtgc	aattatttca	1500
tcactctaaa	aatgctatff	ttgtgtcagt	tcctgcagg	gttttctatg	ctttgcaaa	1560
tgacacattt	tgatgccttc	ttgataaagt	ggtagacatt	ttgtagcttt	ctagaaactt	1620
tgtattcata	cggatatcaat	gaaaaataaa	gaaaatgaaa	gtgtgggtca	aaact	1675

<210> 126

<211> 1064

<212> DNA

<213> Homo sapiens

<400> 126

gcggaggggt	ttctgtgcag	gacgggagtc	tcagagagga	gacggatgtg	ggggagggag	60
gccggccacg	cgggtggacag	agcgaggggtg	ccagggtgac	ccgaagaccg	tcaccacccg	120
acagcaacgc	aagtgccttt	gaccttgatt	tggacttttc	tcctttttgc	atttggtgct	180
acagacttga	gacaccagca	gaagtgtgtg	tcagcccggc	cccgtgctgc	ctgtccgggc	240
cggggctggc	gccggttgtg	tttgtgtcca	ccttgccctc	tttgagcca	agcagttttt	300
gtggatggga	cttacctgca	cgccccagg	gtctttcagg	attcaggatg	acttttcttt	360
tacaatgggt	tcctctcggc	agagcccggg	ttgtggggga	tctgtgtggg	ttctcaacgc	420
agatccatcc	tggggctctcc	cgggcaggga	tggctgacct	cgagtcctct	cccttcccga	480
gaacctgctc	tgtcccagag	gcagctaaca	agggctgagc	cccaggatca	ggttgccctc	540
tccacggcag	gaattttttac	caaaaccaca	agcaaaaaac	aaaacagacc	accacgacca	600
acaacaaaga	tgggggggtar	ggttttgtaa	aggttctgtt	aggttcatat	ttttatatca	660
ttttgcccac	aaatgcggaa	tttgccgtgg	gaatttgaa	acaaatgatc	tatgttttta	720
tggttttcta	gggaargtgt	tctggggggc	gggctctctc	cagctgtggg	aggcctgctc	780
cctctggggg	gcaccctggg	caggggtggg	gggccttggg	aggcgtctct	tgccaaatgc	840
agacgagggg	tgagcctgcc	agcgtttgcg	acgtccccgc	acgacaggct	catactttct	900
gaggatcgtg	catagcatag	gacgtctgaa	cctttgtaca	aatgtgtaga	tgacatcttg	960
ctacagcttt	tatttgtgaa	ttaaagatgc	attgatgggt	aaaaaaaaaa	aaaaaaaaaa	1020
aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaagggcgg	ccgc		1064

<210> 127
 <211> 1607
 <212> DNA
 <213> Homo sapiens

66

<400> 127
 ggcacgagga aaaacttatt tatcgataga cctcatttat gcaaatgttt gttaaagtatc 60
 atgttattat ggtaattatt tttattttca tacttatcac atcagacaag catgggtgaaa 120
 taatatatat taaatacata gatagagtta tcataacaga gagaatctag taaaaaaaaa 180
 agacatgaaa aacatcccat taaagaaaaa gcctttatat gggacattat tcacatttct 240
 gcaaatggaa ttttctgatt aactgccata tattatatta tttgaataac tttaggraag 300
 rttaatttcc aagctctgtc cctggtagta ttttttctct gctgaaattg ggctttaaaa 360
 caaaacgtca gcaatgcaag ctgacgattc cagttacagg ttcagaaact tcgagtgcag 420
 gcagttaaaa gtgatttaag aaaggcaaaa tcattctttc acaaatgtgt atgttaaaact 480
 tgtaaattga tatatcaaaa ccacaaccgg aatatagcaa tttgtctctc tttgcctgct 540
 gtgggtgtctc ccgcagctgc ctgccagata atatactgat tgtaactttc ctaatttaca 600
 gagcctttta gtctaactctc ctgggccagc ctaaagggtc tcttcatcca ttgaaatgcc 660
 agaataattg aaattgtctt cagcagacag tcaggcaata atgaatattt tctctgctga 720
 ttgcatgcc aaggtgcaca tagcactgca aacagaaatg atcccaaacc gagcycccca 780
 gggaggggca gccgctaacc tttggcatga ggcacaatac cgacggcttc ctttctcacg 840
 ggctcctgaa rtcacggacg cccaccaagc ttctgcacaa cgaggtgcag cacagttgcc 900
 gcgtgaacaa tgatgccatt ggcttccctg actctccaga ggcagcttaa aggctgggaa 960
 aggttgtggt tatggatttt ctgatggggg cagcgggaga gatgagcaga ggaggctgct 1020
 gttggcatta agaggaaata taccaatagt taatcaaaaag aaaccggcg ttgaaggggc 1080
 atgagaggac gctggcttgt cagctctgga gcagcatttc ccataccttt gaatggtttt 1140
 tgtgaaagcc gggagttctt tcctgactca ggatcagtgc tgcttcattg gcgtcctaatt 1200
 grtgtgtcga ttgaaattaa agtggtttgg agccgaagcc aatcactaat atcttcaaaa 1260
 aatctaaaga ctagtttaac cttcatctac ggaaagggtg aagaagtttt gaataattaa 1320
 tataagcaac taatcttaga ttatggctac ttccatgata tgcgttagag gtggaaattg 1380
 ggggatgaga aaggaactac cctaagacaa tatggattta agccacatag gaagtgtgca 1440
 gattaaacag ccgggcgtgg tgatgcacct ctgtaatccc agcccagcta ctcatgaggc 1500
 tgaggcagga gaatcgcttg aaccctggag acggagggtt cagtaagccg agattgcac 1560
 actgcactcc agcctcaaaa aaaaaaaaaa aaaaaaaaaa aactcga 1607

<210> 128
 <211> 1037
 <212> DNA
 <213> Homo sapiens

<400> 128
 agctactccg tctggccccg ctttttctct gctctcctga acctttaggg ttgtctcggc 60
 ccatttgaag accaggaagt tgatcaatcc cgaggctgct gagagacggg ggcgcgattg 120
 ggacagtcgc cagggatggc tgagcgtgaa gatgcagcgg gtgtccgggc tgctctcctg 180
 gacgctgagc agagtcctgt ggctctccgg cctctctgag ccgggagctg cccggcagcc 240
 ccggatcatg gaagagaaag cgctagaggt ttatgatttg attagaacta tccgggaccc 300
 agaaaagccc aatacttttag aagaactgga agtggctctg gaaagtgtg tggaagttca 360
 ggagataaat gaagaagaat atctggttat tatcaggttc acgccaacag tacctcattg 420
 ctctttggcg actcttattg ggctgtgctt aagagtaaaa cttcagcgat gtttaccatt 480
 taaacataag ttggaaatct acattttctga aggaaccac tcaacagaag aagacatcaa 540
 taagcagata aatgacaaag agcgagtggc agctgcaatg gaaaacccca acttacggga 600
 aattgtggaa cagtgtgtcc ttgaacctga ctgatagctg ttttaagagc cactggcctg 660
 taattgtttg atatatattg ttaaaactct tgtataatgt cagagactca tgtttaatac 720
 ataggtgatt tgtacctcag agcatttttt aaaggattct ttccaagcga gatttaatta 780
 taaggtagta cctaatttgt tcaatgtata acattctcag gatttgtaac acttaaatga 840
 tcagacagaa taatattttc tagttattat gtgtaagatg agttgctatt tttctgatgc 900
 tcattctgat acaactattt ttcgtgtcaa atatctactg tgcccaaatg tactcaattt 960

b7

aaatcattac tctgtaaaat aaataagcag atgattctta aaaaaaaaaa ataaaaaaaaa	1020
aaaaaaaaaggg cgccgcg	1037

<210> 129
 <211> 1146
 <212> DNA
 <213> Homo sapiens

<400> 129	
ggcacgagat ttttggtcag gtttttatgt tgtttttcag tttgaaggag tcgttatata	60
tttttcatac tgctattttg ttagtagtat gctttgcatg tgtagtagta tgcagtagta	120
taatagtacg tgtagtgca gtagtatgtt gcttttcaaa atctcagtct ttgatctgaa	180
tttaaagaat agtatgtttt tgtgtaagag gttatgttta tcttttgcta tttaaaaaat	240
caagtgccat ttgtatatga tagtacttaa gtcttcagtt tgaattctaa aaatgtgtat	300
acctatataa tccacacacc tgtcatccca tcctctatc aacacccctg aaaattccct	360
tgtgcctctt ctagtcaac cccaccactg cctcaagtaa gtactcttct gatttctgtt	420
ctcatagatt agtttcaact gctctagaac atcatataat ggggctgggc gcgatggctc	480
acgccagtaa tcccagcact tgggaggctg aggtgggcag atcacttgag gtcaggagtt	540
tgagaccagc ctggcaaaca tgggtgaaacc ctgtctctac taaaaatgca aaaattaacc	600
tgggtgtgtg gcatgccttt tgtaatcccc agctactcca ggagtccaag gcccgagaat	660
cgcttgaacc tgggagacag aggttgcaact gagctgctgc actccagcct gggcaacaga	720
gtgagactgt ctaagaaaaa agaagtttat atgatggaat tatacagtaa ataactctga	780
tctatgttca tttagcattg tatctatgaa attcatgcat gttttatcta ttaatatgtt	840
ttttggccgg gctcgggtgc tcacgcctgt aatcccagca cttttggagg ctgaggcagg	900
tggatcacta ggtagcagg tcaagaccag cctaggcaaa gatgggtgaaa ccccatctct	960
actaaaaata caaaaattag ctgggcatgg tggcaggcgc ctgtaatccc agctactcag	1020
gaggctgagg cagagaattg cttgaacctg ggaggcagag gttgcagcaa gctgagatcg	1080
tgccactgca ctccagcctg ggcgacagag cgagactcca tctcaaaaaa aaaaaaaaaa	1140
aaaaaa	1146

<210> 130
 <211> 1172
 <212> DNA
 <213> Homo sapiens

<400> 130	
ccacgcgtcc gaacaccttg cagctttacc tgtgaatgtc aagattggca agatgcttat	60
ttttggtgcc atatttggtt gccttgaccc agtggcaaca ctagctgcag ttatgacaga	120
gaagtctcct tttaccacac caattggctg aaaagatgaa gcagatcttg caaaatcagc	180
tttggtcatg gcggattcag accacctgac gatctacaat gcatacttag gatggaagaa	240
agcagacaaa gaaggagggt atcgttctga aatcacatac tgccggagga actttcttaa	300
tagaacatca ctgttaaccc tagaggatgt aaagcaggag ttaataaagt tgggtaaggc	360
agcaggattt tcatcttcca caacttctac cagctgggaa ggaaacagag cctcacagac	420
cctctcattc caagaaattg cccttcttaa agctgtactg gtggctggac tgtatgacaa	480
tgtggggaag ataacttata caaagtcagt ggatgttaca gaaaaattgg ctgcatgtgt	540
ggagacggcc caaggcaaag cacaagtaca cccatcctca gtaaatcgag atttgcaaac	600
tcatggatgg ctcttatacc aggagaagat aaggatgccc agagtgtatt tgagagaaac	660
taccctaata accccttttc cagttttact ttttggtggt gatatagaag ttcagcaccg	720
agaacgtctt ctttctattg atggctggat ctattttcag gccctgttaa agatagctgt	780
cattttcaag cagctgagag ttctcattga ttcagtttta agaaaaagc ttgaaaatcc	840
aaagatgtcc cttgaaatga caagattctg cagatcatta cggaattgat aaaaacagag	900
aataactgaa actgaaattc atggtcaact gctttaaaaa ttaagatgaa gatacagtca	960
tgaaattatc tgaaatggg tcatcacatt aagtatttca ttacttaaaa tgttggtact	1020
agccattaac ttaaagggtg tgggaaaaaa gcacatactt taaacatgta taattttcta	1080
gttccttttt aatgatgati attctgaatg tatttgccac tacattttaca ataattctt	1140
tgggtattatg aaaaaaaaaa aaaaaaaaaa aa	1172

68

<210> 131
 <211> 663
 <212> DNA
 <213> Homo sapiens

<400> 131
 ggcacgagaa accatgaaag tcctttcttg gatccacttt atcttgatta gtctgcattt 60
 tactagttca ctggatccct cctctagggg cctggggact ttcactgatg ctcttcctga 120
 ttctagagca aagggtgtggg aaggggaaat ggaggaatgc cctcctgtct gtgtcgttct 180
 ctgtgccaca gctacagatg cagaagggtt ctctggatag cacacctctg aatgtaaatac 240
 atgataaaat ggatattttg aaacttactc ctaagctgtg atttaggggtg tatttctact 300
 tctggactgc ctcaatatca agggctgaga cttttgaatt ttgaatattc gttgggtttc 360
 atgttaagaa gcctgtgggc taggagtgtt attcagtgtt tcttttcctg ataaacactt 420
 tgaatatttt ttttgtgttt ttgtttcctt ttctgaagct gttcctcctt ttaaataattt 480
 ttaatcacat tgataaaatc tatecttcac cacctctggg tctactatag ttgattttta 540
 ttttaaatgt ttaattgtat ttgattaaac acttaactgg attttggaat aataaaactc 600
 tcgtccaatt tggcttttaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 660
 aaa 663

<210> 132
 <211> 776
 <212> DNA
 <213> Homo sapiens

<400> 132
 ggcacgagct gatttctatt tttaggagct acttggattt gtatgtattt tttctacgtg 60
 aaaatatatg tactcttcac ttttgttcca gtactataat tgctcatgca ctctttctcc 120
 cttttgagaa cattcagtga aatacaactt catcaaagat ttgctcaaag gagaagaatc 180
 gcatgagtgt gaaaagtaga tgctcgtagc cagaacagaa aagggttacac atgatcatgg 240
 cacagaagat aggaggtttg acttgggtggg ccataatgtt tattatcctt tttgaaataa 300
 cagggaccag cagcagtttt ctcaggataa atgctctacc ccacttctct atgaacaggt 360
 gtggggaggc ttactttcca ttttcatatt tatacacctc tctacaaaag caatttttaa 420
 tgaagggttag tggaattgtt aaaaatctga gaggggaatga tgactggagg tgttttgggg 480
 tttttttctg tattcatttt ttaatgagaa aagtttttaa ttagtagacag gttagaccca 540
 actactacct tactattata ggacgattct atgtttctgt taaagtattc aagtagcttt 600
 ctctggggga aaaagtagca cttggacact taaaggaatt gggatttttg tctactttgg 660
 ataaggcagt tgacttctta agtaaaagca atagtgtaaa atgtcatttt gtttggaatg 720
 ttaagtgagc aaataaaaaa catgttgaaa ttgtaaaaaa aaaaaaaaaa aaaaaa 776

<210> 133
 <211> 1543
 <212> DNA
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (1055)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (1143)
 <223> n equals a,t,g, or c

<400> 133 69

cttcgcccgtc	atccgcttcg	aaagcatcat	ccacgagttc	gacccgtggt	ttactatag	60
atcaacacat	catcttgcat	ctcatgggtt	ctatgaattt	ttaaattggt	ttgatgaaag	120
agcatgggtat	ccactaggaa	gaatagtagg	tggtactggt	taccagggt	tgatgataac	180
cgctggcctt	attcattgga	ttttaaatat	attgaacata	actgttcaca	taagagacgt	240
atgtgtgttc	cttgaccaa	cttttagcgg	ccttacatct	atatctactt	tcctgcttac	300
aagagaactt	tggaaccaag	gagcaggact	tttagctgct	tgttttattg	ctattgtacc	360
aggctacata	tctcggtcag	tagctggatc	ctttgataat	gaaggcattg	ctatttttgc	420
acttcagttc	acatactatt	tatgggtaaa	atctgtaaaa	actgggtcag	ttttttggac	480
aatgtgctgc	tgcttatcct	atttctatat	ggctctgct	tgggggtggt	atgtatttat	540
catcaatctt	attccactgc	atgtatttgt	gttggtactg	atgcagagat	acagcaaaag	600
agtctacata	gcataatgca	ctttctacat	tgtgggttta	atattatcaa	tgcatatacc	660
ttttgtggga	ttccagccaa	tcagaacaag	tgaacacatg	gcagctgcag	gtgtctttgc	720
attgctgcaa	gcttatgctt	tcttgacgta	tctgagagac	cgattaacaa	aacaagagtt	780
ccagaccctt	ttcttttttg	gtgtatcact	agctgcagg	gctgtgttcc	ttagtgtcat	840
ctatttgact	tatacagggt	acattgcacc	atggagtggc	aggttttatt	cattgtggga	900
tactgggtat	gcaaaaatac	acattccaat	tattgcata	gtgtctgagc	atcaacctac	960
gacttgggtg	tctttcttct	ttgatctaca	tattcttgta	tgtaccttcc	cagcaggcct	1020
ttggttctgc	atcaaaaata	tcaacgatga	aagantattt	ggtaagarag	gtttttaatg	1080
actactttga	tatggaatag	ttatttttct	ttttgagatt	atttacttta	aatttttgtt	1140
tttctatggt	tgactctata	tattcaagat	aaattttctc	ctttattttg	cataggtgct	1200
taaccaagaa	aaattcactg	agaggctggg	catggtggca	cacgcctkta	atcccagcac	1260
tttgggaggc	cgaggcgggc	ggatcacctg	aggtcaggag	ttcgagacca	gcctggccaa	1320
catggtgaaa	ccttgctctc	actaaaaata	caaaaattag	ccgaacatgg	tggtgcatgc	1380
ctgtaatccc	agctactcag	gaggctgagg	caggataatt	gcttgaacct	gggaggcgga	1440
ggttgcatg	agtcaagatc	aagccactgc	actccaccct	gggaatcaga	gcgggactct	1500
gtctcaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaggcgggc	cgc		1543

<210> 134

<211> 2157

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (309)

<223> n equals a,t,g, or c

<400> 134

caaaaaggac	cgccattga	agatgccatt	gcttcttccg	atgttcttga	gactgcttct	60
aaatctgcta	atccaccca	cacgattcaa	gcatcagaag	agcagagttc	aacccagca	120
ccggtgaaaa	agtctggcaa	gctgaggcag	caaatagatg	tgaaggcgga	actggagaag	180
cggcaaggag	ggaagcagct	actcaactta	gtggtcattg	gtcatgttga	tgctgggaaa	240
agtactctga	tgggccatat	gctttatctt	ctgggtaata	taaacaaaag	aactatgcat	300
aagtatganc	aggagtctaa	aaaggctggc	aaagcttcgt	ttgcatatgc	atgggtcttg	360
gatgaaactg	gcgaagaaa	ggaaagggga	gtaaccatgg	atgttggtat	gacaaagttt	420
gaaaccacaa	ccaaagtatt	tacattaatg	gatgtctcag	gccataagga	cttcattcca	480
aatatgatta	caggagcagc	ccaggcggat	gtagctgttt	tagttgtaga	tgccagcagg	540
ggagagtttg	aagctggatt	tgagactgga	ggacaaacac	gagagcatgg	actcttggtc	600
cgttctctgg	gagtgcagca	gcttgagttt	gcagtttaata	aaatggatca	ggtaatttgg	660
caacaagaaa	ggtttcaaga	gattactgga	aaacttgggc	actttcttaa	gcaagcagg	720
tttaaggaga	gtgatgtagg	ttttattcct	acaagtggte	tcagtgggtga	aaatctaata	780
acaagatctc	agtcaagtga	actcacaata	tggtataaag	gactatgttt	attagaacaa	840
attgattcct	ttaagcctcc	ccagcgatct	attgacaaac	cttttagatt	atgtgtgtcc	900
gatgttttca	aagatcaagg	atctggattt	tgcataactg	gtaaaataga	agctgggtat	960
atccaaactg	gtgaccgact	actggcaatg	cctcctaata	aaacttgtac	cgtgaaagga	1020
atcactctgc	atgatgaacc	tgctcgactgg	gcggcagcag	gcgatcatgt	tagtcttact	1080

70

ttgggttgga	tggatatcat	caaaatcaat	gttggtgca	tattttgtgg	cccaaagta	1140
cccattaaag	cttgactcgc	tttcagagcc	cgaatcctca	tctttaatat	tgaaattcct	1200
atcactaaag	gatttcctgt	gctgttacac	taccaaactg	tcagtgaacc	cgccgttatt	1260
aaacgattga	ttagtgtctt	aaacaaaagc	acgggtgaag	tcacaaagaa	aaagcctaag	1320
tttttgacta	aaggccagaa	tgcattggta	gagctacaga	cacaaagacc	aatagctctt	1380
gagctatata	aagactttta	agagctgggg	aggttcatgc	tacgttacgg	tggttctaca	1440
atagctgctg	gtgttgtcac	tgagataaaa	gaatgatggg	tcmgaatttc	taccacgttt	1500
ctggatacag	tgaaatagct	aacctctgty	tcaagaatgc	agttattaag	tcaaaggaac	1560
aatgtgcaat	tgatatgttt	ttagatgaga	gagaaaaatt	aaagctaaaa	ttagctgcaa	1620
agaagtatta	ataatcacct	ctgcaaaaat	tctaagttgc	caactggcaa	agraagtcta	1680
atgttaaaaa	caactttgcc	tttgaamcgt	taataaatgg	atttactttg	ctaagattta	1740
tggaagtgt	caaaaatagt	atctgaagat	actgaatcat	catgaaatga	actctacttc	1800
tggaacaaagc	acaatgtatt	tgcaagtttc	tcttttgatt	caattatact	gcacatgttt	1860
taaggaaaag	taacttaatt	gggtttttca	ggcagttgat	atttgacct	agcttttttt	1920
tttttttttt	ttccagttaa	tgctaagaaa	agatttgggg	aagggtataa	taaaagtatt	1980
ttgtggtgac	cataagaatg	tccctcccca	aacaagtaaa	cttgtagaaag	tttaattttg	2040
aattagtggg	agctgttcct	ttgaaagcca	agatattatt	taagttgtaa	agccagctaa	2100
taaaatgcct	tagtttgagc	ataaaaaaaaa	aaaaaaaaaaa	aaaaaaaaaaa	actcgag	2157

<210> 135

<211> 420

<212> DNA

<213> Homo sapiens

<400> 135

ggcacgagag	agagcagagc	tatacatagc	tatccaggtc	taacttcacg	aagaatagaa	60
tggtttcttt	tcattttcaa	tgtacatcat	actttgtcag	actttttttt	cagttgcagc	120
tcttcgttgg	actggtgata	gtattggctt	tattaatctc	tcattctctc	acttattcat	180
tccacaaaca	ttttagaag	gccaccaagc	tctagggaga	ggaaaatggt	tttataaatt	240
agtgtcttct	gggataaagg	aaatttataa	tctgtactac	ttaatagtag	ccactagcca	300
catgtgggtt	tcgaacaaga	tttccatcac	ctctccaacc	actttctcct	cattgggtcag	360
atctagaccc	cgagaaactg	ttcctttcat	tgttttctcc	gccttctaca	aactgagata	420

<210> 136

<211> 334

<212> PRT

<213> Homo sapiens

<400> 136

Met	Phe	Gln	Cys	Gly	Leu	Leu	Gln	Gln	Leu	Cys	Thr	Ile	Leu	Met	Ala
1					5				10					15	

Thr	Gly	Val	Pro	Ala	Asp	Ile	Leu	Thr	Glu	Thr	Ile	Asn	Thr	Val	Ser
			20					25					30		

Glu	Val	Ile	Arg	Gly	Cys	Gln	Val	Asn	Gln	Asp	Tyr	Phe	Ala	Ser	Val
		35					40					45			

Asn	Ala	Pro	Ser	Asn	Pro	Pro	Arg	Pro	Ala	Ile	Val	Val	Leu	Leu	Met
		50					55				60				

Ser	Met	Val	Asn	Glu	Arg	Gln	Pro	Phe	Val	Leu	Arg	Cys	Ala	Val	Leu
	65					70				75					80

Tyr	Cys	Phe	Gln	Cys	Phe	Leu	Tyr	Lys	Asn	Gln	Lys	Gly	Gln	Gly	Glu
						85				90					95

71

Ile Val Ser Thr Leu Leu Pro Ser Thr Ile Asp Ala Thr Gly Asn Ser
 100 105 110
 Val Ser Ala Gly Gln Leu Leu Cys Gly Gly Leu Phe Ser Thr Asp Ser
 115 120 125
 Leu Ser Asn Trp Cys Ala Ala Val Ala Leu Ala His Ala Leu Gln Glu
 130 135 140
 Asn Ala Thr Gln Lys Glu Gln Leu Leu Arg Val Gln Leu Ala Thr Ser
 145 150 155 160
 Ile Gly Asn Pro Pro Val Ser Leu Leu Gln Gln Cys Thr Asn Ile Leu
 165 170 175
 Ser Gln Gly Ser Lys Ile Gln Thr Arg Val Gly Leu Leu Met Leu Leu
 180 185 190
 Cys Thr Trp Leu Ser Asn Cys Pro Ile Ala Val Thr His Phe Leu His
 195 200 205
 Asn Ser Ala Asn Val Pro Phe Leu Thr Gly Gln Ile Ala Glu Asn Leu
 210 215 220
 Gly Glu Glu Glu Gln Leu Val Gln Gly Leu Cys Ala Leu Leu Leu Gly
 225 230 235 240
 Ile Ser Ile Tyr Phe Asn Asp Asn Ser Leu Glu Ser Tyr Met Lys Glu
 245 250 255
 Lys Leu Lys Gln Leu Ile Glu Lys Arg Ile Gly Lys Glu Asn Phe Ile
 260 265 270
 Glu Lys Leu Gly Phe Ile Ser Lys His Glu Leu Tyr Ser Arg Ala Ser
 275 280 285
 Gln Lys Pro Gln Pro Asn Phe Pro Ser Pro Glu Tyr Met Ile Phe Asp
 290 295 300
 His Glu Phe Thr Lys Leu Val Lys Glu Leu Glu Gly Val Ile Thr Lys
 305 310 315 320
 Ala Ile Tyr Lys Ser Ser Glu Glu Asp Lys Lys Lys Lys Lys
 325 330

<210> 137

<211> 43

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (43)

<223> Xaa equals stop translation

<400> 137

22

Met Thr Val Ala Ser Ile Arg His Ile Leu Val Glu Ile Trp Leu Pro
 1 5 10 15

Ile Ala Leu Ala Met Gly Thr Arg Gly Leu Thr Gln Ile Val Ala Val
 20 25 30

Ile Gln Ser Arg Ser Gln Trp Ala Leu Ser Xaa
 35 40

<210> 138

<211> 87

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (87)

<223> Xaa equals stop translation

<400> 138

Met Leu Phe Ile Phe Leu Leu Leu Ile Leu Ser Ile Thr Ala Ser Tyr
 1 5 10 15

Ser Leu Thr Cys Ile Leu Ser Gly Ala Gly Glu Pro Ser Ser Val Ser
 20 25 30

Ala Ser Val Val Ser Gly Pro Gly Phe Cys Leu Ala Ala Leu Leu Leu
 35 40 45

Met Arg Thr Gly Gly Phe Ala Ala Thr Leu Leu Pro Val Ala Pro Thr
 50 55 60

Glu Arg Phe Phe Ser Cys Cys Thr Val Leu Ser Ala Gln Arg Asn Val
 65 70 75 80

Ser Arg Thr Arg Ser Pro Xaa
 85

<210> 139

<211> 122

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (122)

<223> Xaa equals stop translation

<400> 139

Met Leu Ser Thr Arg Trp Met Gly Leu His Leu Val Gln Ile Leu Trp
 1 5 10 15

Arg Cys Trp Thr Ser Ser Ala Thr Ile Thr Ser Arg Lys Leu Ser Thr
 20 25 30

Ala Leu Arg Ser Pro Val Leu Ser Gly Thr Gln Thr Ser Arg Ser Ser

35 40 73 45
 Gly Asp Ser Gly Trp Ser Met Lys Thr Ser Val Lys Ala Thr Pro His
 50 55 60
 Gln Met Ser Leu Arg Ser Gly Lys Glu Thr Pro Ser Ala Asp Ile Pro
 65 70 75 80
 Arg Ile His His Gln Leu Val Arg Leu Arg His Gln Ala His Gly Gly
 85 90 95
 Trp Ser Pro His Gly Val Pro Glu Gln Gly Thr Met Pro Leu Val Leu
 100 105 110
 Pro Pro Val Ser Cys Asp Ile Gln Pro Xaa
 115 120

<210> 140

<211> 276

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (131)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (276)

<223> Xaa equals stop translation

<400> 140

Met Ala Asn Thr Gly Val Phe Gly Phe Ser Phe Leu Leu Leu Thr Val
 1 5 10 15
 Ala Leu Leu Ala Ser Tyr Ser Val His Leu Leu Leu Ser Met Cys Ile
 20 25 30
 Gln Thr Ala Val Thr Ser Tyr Glu Asp Leu Gly Leu Phe Ala Phe Gly
 35 40 45
 Leu Pro Gly Lys Leu Val Val Ala Gly Thr Ile Ile Ile Gln Asn Ile
 50 55 60
 Gly Ala Met Ser Ser Tyr Leu Leu Ile Ile Lys Thr Glu Leu Pro Ala
 65 70 75 80
 Ala Ile Ala Glu Phe Leu Thr Gly Asp Tyr Ser Arg Tyr Trp Tyr Leu
 85 90 95
 Asp Gly Gln Thr Leu Leu Ile Ile Ile Cys Val Gly Ile Val Phe Pro
 100 105 110
 Leu Ala Leu Leu Pro Lys Ile Gly Phe Leu Gly Tyr Thr Ser Ser Leu
 115 120 125

24

Ser Phe Xaa Phe Met Met Phe Phe Ala Leu Val Val Ile Ile Lys Lys
 130 135 140

Trp Ser Ile Pro Cys Pro Leu Thr Leu Asn Tyr Val Glu Lys Gly Phe
 145 150 155 160

Gln Ile Ser Asn Val Thr Asp Asp Cys Lys Pro Lys Leu Phe His Phe
 165 170 175

Ser Lys Glu Ser Ala Tyr Ala Leu Pro Thr Met Ala Phe Ser Phe Leu
 180 185 190

Cys His Thr Ser Ile Leu Pro Ile Tyr Cys Glu Leu Gln Ser Pro Ser
 195 200 205

Lys Lys Arg Met Gln Asn Val Thr Asn Thr Ala Ile Ala Leu Ser Phe
 210 215 220

Leu Ile Tyr Phe Ile Ser Ala Leu Phe Gly Tyr Leu Thr Phe Tyr Gly
 225 230 235 240

Ser His Ser Val Ala Gln Val Gly Val Gln Trp Cys Asp Leu Ser Ser
 245 250 255

Leu Gln Pro Leu Pro Pro Gly Leu Lys Gln Ser Ser His Leu Ser Leu
 260 265 270

Gln Ser Ser Xaa
 275

<210> 141

<211> 195

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (138)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (195)

<223> Xaa equals stop translation

<400> 141

Met Lys Leu Ala Ser Gly Phe Leu Val Leu Trp Leu Ser Leu Gly Gly
 1 5 10 15

Gly Leu Ala Gln Ser Asp Thr Ser Pro Asp Thr Glu Glu Ser Tyr Ser
 20 25 30

Asp Trp Gly Leu Arg His Leu Arg Gly Ser Phe Glu Ser Val Asn Ser
 35 40 45

Tyr Phe Asp Ser Phe Leu Glu Leu Leu Gly Gly Lys Asn Gly Val Cys
 50 55 60

Gln Tyr Arg Cys Arg Tyr Gly Lys Ala Pro ^XMet Pro Arg Pro Gly Tyr
 65 70 75 80
 Lys Pro Gln Glu Pro Asn Gly Cys Gly Ser Tyr Phe Leu Gly Leu Lys
 85 90 95
 Val Pro Glu Ser Met Asp Leu Gly Ile Pro Ala Met Thr Lys Cys Cys
 100 105 110
 Asn Gln Leu Asp Val Cys Tyr Asp Thr Cys Gly Ala Asn Lys Tyr Arg
 115 120 125
 Cys Asp Ala Lys Phe Arg Trp Cys Leu Xaa Ser Ile Cys Ser Asp Leu
 130 135 140
 Lys Arg Ser Leu Gly Phe Val Ser Lys Val Glu Ala Cys Asp Ser Leu
 145 150 155 160
 Val Asp Thr Val Phe Asn Thr Val Trp Thr Leu Gly Cys Arg Pro Phe
 165 170 175
 Met Asn Ser Gln Arg Ala Ala Cys Ile Cys Ala Glu Glu Glu Lys Glu
 180 185 190
 Glu Leu Xaa
 195

<210> 142

<211> 183

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (183)

<223> Xaa equals stop translation

<400> 142

Met Leu Leu Leu Cys His Ala Leu Ala Ile Ala Val Val Gln Ile Val
 1 5 10 15
 Ile Phe Ser Glu Ser Trp Ala Phe Ala Lys Asn Ile Asn Phe Tyr Asn
 20 25 30
 Val Arg Pro Pro Leu Asp Pro Thr Pro Phe Pro Asn Ser Phe Lys Cys
 35 40 45
 Phe Thr Cys Glu Asn Ala Gly Asp Asn Tyr Asn Cys Asn Arg Trp Ala
 50 55 60
 Glu Asp Lys Trp Cys Pro Gln Asn Thr Gln Tyr Cys Leu Thr Val His
 65 70 75 80
 His Phe Thr Ser His Gly Arg Ser Thr Ser Ile Thr Lys Lys Cys Ala
 85 90 95

76

Ser Arg Ser Glu Cys His Phe Val Gly Cys His His Ser Arg Asp Ser
 100 105 110

Glu His Thr Glu Cys Arg Ser Cys Cys Glu Gly Met Ile Cys Asn Val
 115 120 125

Glu Leu Pro Thr Asn His Thr Asn Ala Val Phe Ala Val Met His Ala
 130 135 140

Gln Arg Thr Ser Gly Ser Ser Ala Pro Thr Leu Tyr Leu Thr Ser Ala
 145 150 155 160

Cys Leu Gly Leu Cys Ala Ser Ile Ala Val Met Pro Pro Phe Leu Gly
 165 170 175

Glu Ala Glu Thr Ser Leu Xaa
 180

<210> 143
 <211> 122
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (122)
 <223> Xaa equals stop translation

<400> 143
 Met Leu Arg Gly Thr Met Thr Ala Trp Arg Gly Met Arg Pro Glu Val
 1 5 10 15

Thr Leu Ala Cys Leu Leu Leu Ala Thr Ala Gly Cys Phe Ala Asp Leu
 20 25 30

Asn Glu Val Pro Gln Val Thr Val Gln Pro Ala Ser Thr Val Gln Lys
 35 40 45

Pro Gly Gly Thr Val Ile Leu Gly Cys Val Val Glu Pro Pro Arg Met
 50 55 60

Asn Val Thr Trp Arg Leu Asn Gly Lys Glu Leu Asn Gly Ser Asp Asp
 65 70 75 80

Ala Leu Gly Val Leu Ile Thr His Gly Thr Leu Val Ile Thr Ala Leu
 85 90 95

Asn Asn His Thr Val Gly Arg Tyr Gln Cys Val Ala Arg Met Pro Ala
 100 105 110

Gly Ala Val Ala Thr Cys Gln Pro Leu Xaa
 115 120

<210> 144
 <211> 267
 <212> PRT

<213> Homo sapiens

77

<220>

<221> SITE

<222> (267)

<223> Xaa equals stop translation

<400> 144

Met Trp Trp Phe Gln Gln Gly Leu Ser Phe Leu Pro Ser Ala Leu Val
 1 5 10 15

Ile Trp Thr Ser Ala Ala Phe Ile Phe Ser Tyr Ile Thr Ala Val Thr
 20 25 30

Leu His His Ile Asp Pro Ala Leu Pro Tyr Ile Ser Asp Thr Gly Thr
 35 40 45

Val Ala Pro Glu Lys Cys Leu Phe Gly Ala Met Leu Asn Ile Ala Ala
 50 55 60

Val Leu Cys Ile Ala Thr Ile Tyr Val Arg Tyr Lys Gln Val His Ala
 65 70 75 80

Leu Ser Pro Glu Glu Asn Val Ile Ile Lys Leu Asn Lys Ala Gly Leu
 85 90 95

Val Leu Gly Ile Leu Ser Cys Leu Gly Leu Ser Ile Val Ala Asn Phe
 100 105 110

Gln Lys Thr Thr Leu Phe Ala Ala His Val Ser Gly Ala Val Leu Thr
 115 120 125

Phe Gly Met Gly Ser Leu Tyr Met Phe Val Gln Thr Ile Leu Ser Tyr
 130 135 140

Gln Met Gln Pro Lys Ile His Gly Lys Gln Val Phe Trp Ile Arg Leu
 145 150 155 160

Leu Leu Val Ile Trp Cys Gly Val Ser Ala Leu Ser Met Leu Thr Cys
 165 170 175

Ser Ser Val Leu His Ser Gly Asn Phe Gly Thr Asp Leu Glu Gln Lys
 180 185 190

Leu His Trp Asn Pro Glu Asp Lys Gly Tyr Val Leu His Met Ile Thr
 195 200 205

Thr Ala Ala Glu Trp Ser Met Ser Phe Ser Phe Phe Gly Phe Phe Leu
 210 215 220

Thr Tyr Ile Arg Asp Phe Gln Lys Ile Ser Leu Arg Val Glu Ala Asn
 225 230 235 240

Leu His Gly Leu Thr Leu Tyr Asp Thr Ala Pro Cys Pro Ile Asn Asn
 245 250 255

Glu Arg Thr Arg Leu Leu Ser Arg Asp Ile Xaa
 260 265

78

<210> 145
 <211> 92
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (92)
 <223> Xaa equals stop translation

<400> 145
 Met Leu Cys His Pro His Val His His His Leu Val Cys Leu Leu Ala
 1 5 10 15
 Thr Leu Thr Phe Ser Leu Asn Ala Ser Cys Ala Glu Gln Thr Phe His
 20 25 30
 Ser Gln Gln Ser Asn Gly Glu Phe Met Ala Thr Leu Pro Ser Ile Ser
 35 40 45
 Lys Gln Phe Gly Val Ile Val Trp Lys Pro Gln Arg Lys Asp Val Ile
 50 55 60
 Arg Leu Pro Val Ala Leu Ser Phe Ser Met Gly Leu Gly Leu Leu Ser
 65 70 75 80
 Pro Ala Leu Gly Arg Phe Leu Ala Ser Glu Leu Xaa
 85 90

<210> 146
 <211> 109
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (109)
 <223> Xaa equals stop translation

<400> 146
 Met Ala Ile Leu Leu Ala Cys Phe Thr Ala Val Leu Ala Phe Ile Cys
 1 5 10 15
 Leu Gln Phe Trp Cys Val Arg Cys His Glu Pro Arg Trp Ser Tyr Arg
 20 25 30
 Ala Gly His Met Glu Glu Ala Asn Gly Leu Val Arg Trp Pro Glu Glu
 35 40 45
 Ala Pro Asp Leu Gly Gln Arg Glu Glu Asp Leu Gln Gly Leu Pro Leu
 50 55 60
 Val Glu Met Pro Arg Lys Asn Ser Arg Asp Gly Ala Glu Leu Asp Pro
 65 70 75 80

<400> 148

Met	Ser	Arg	Val	Val	Ser	Leu	Phe	Phe	Phe	Ile	Leu	Phe	Ser	Phe	Phe	
1				5					10					15		
Phe	Phe	Ala	Phe	Ser	Leu	Ser	Ser	Ser	Leu	Ser	Phe	Val	His	Tyr	Glu	
			20					25					30			
Lys	Leu	Val	Gln	Val	Lys	Glu	Cys	Leu	Asp	Ser	Phe	Leu	Lys	Lys	Ile	
		35					40					45				
Lys	Ile	Lys	Glu	Tyr	Lys	Thr	Arg	Gln	Cys	Tyr	His	Leu	Ile	Arg	Trp	

50 55 80 60
 Glu Asn Asn Gly Ala Lys Leu Gln Ser Xaa
 65 70

 <210> 149
 <211> 72
 <212> PRT
 <213> Homo sapiens

 <220>
 <221> SITE
 <222> (72)
 <223> Xaa equals stop translation

 <400> 149
 Met Ser Ala Ser Leu Lys Asn His Leu Thr His Cys Phe Leu Leu Leu
 1 5 10 15
 Leu Leu Lys Glu Leu Val Ser Pro Thr Met Ile Ser Phe Val Pro Thr
 20 25 30
 Leu Arg His Ser Tyr Arg Phe Phe Asn Leu Phe Ser Cys Asp Ala Glu
 35 40 45
 Ser Thr Lys Glu Ser Pro Gly Arg Thr Val Gln Phe Ser Lys Thr Pro
 50 55 60
 Arg Gly Val Thr Met Phe Ile Xaa
 65 70

 <210> 150
 <211> 152
 <212> PRT
 <213> Homo sapiens

 <220>
 <221> SITE
 <222> (152)
 <223> Xaa equals stop translation

 <400> 150
 Met Lys Tyr Gly Leu Thr Gly Pro Trp Ile Lys Arg Leu Leu Pro Val
 1 5 10 15
 Ile Phe Leu Val Gln Ala Ser Gly Met Asn Val Tyr Met Ser Arg Ser
 20 25 30
 Leu Glu Ser Ile Lys Gly Ile Ala Val Met Asp Lys Glu Gly Asn Val
 35 40 45
 Leu Gly His Ser Arg Ile Ala Gly Thr Lys Ala Val Arg Glu Thr Leu
 50 55 60
 Ala Ser Arg Ile Val Leu Phe Gly Thr Ser Ala Leu Ile Pro Glu Val
 65 70 75 80

<220>
<221> SITE

<222> (94) ⁹²
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (97)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (98)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (123)
 <223> Xaa equals stop translation

<400> 152
 Met His Arg Ser Glu Pro Phe Leu Lys Met Ser Leu Leu Ile Leu Leu
 1 5 10 15
 Phe Leu Gly Leu Ala Glu Ala Cys Thr Pro Arg Glu Val Asn Leu Leu
 20 25 30
 Lys Gly Ile Ile Gly Leu Met Ser Arg Leu Ser Pro Asp Glu Ile Leu
 35 40 45
 Gly Leu Leu Ser Leu Gln Val Leu His Glu Glu Thr Ser Gly Cys Lys
 50 55 60
 Glu Glu Val Lys Pro Phe Ser Gly Thr Thr Pro Ser Arg Lys Pro Leu
 65 70 75 80
 Pro Lys Arg Glu Glu His Val Glu Xaa Pro Xaa Asn Ala Xaa Thr Trp
 85 90 95
 Xaa Xaa Thr Tyr Leu Phe Val Ser Tyr Asn Lys Gly Asp Trp Phe Thr
 100 105 110
 Phe Ser Ser Gln Val Leu Leu Pro Leu Leu Xaa
 115 120

<210> 153
 <211> 55
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (55)
 <223> Xaa equals stop translation

<400> 153
 Met Ser Pro Cys Ala His Ile Cys Leu Tyr Val Leu Val Phe Leu Cys
 1 5 10 15

93

Asn	Val	Thr	Arg	Cys	Lys	Cys	Val	Arg	Ala	Phe	Thr	Thr	Trp	Asp	Thr
			20					25					30		
Glu	Lys	Val	Lys	Tyr	Phe	Met	Ala	His	Trp	Ser	Lys	Leu	Lys	Arg	Val
		35					40					45			
Arg	Gly	Thr	Arg	Val	Glu	Xaa									
		50				55									

<210> 154

<211> 111

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (93)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (111)

<223> Xaa equals stop translation

<400> 154

Met	Phe	Leu	Ala	Ser	Trp	Leu	Leu	Phe	Cys	Ile	Val	Ala	Pro	Lys	Asp
1				5					10					15	

Asp	Ala	His	Leu	Ser	Phe	Ile	Gln	Cys	Lys	Asp	Ile	Trp	Lys	Asp	Asn
			20					25					30		

Arg	Lys	Tyr	Ser	Cys	Phe	His	Phe	Lys	Ser	Asp	Gln	Leu	Leu	Glu	Leu
		35					40					45			

Ala	Ser	Lys	Ala	Cys	Thr	Ser	Phe	Gln	Ala	Gln	Ser	Arg	Ser	Phe	Thr
	50					55					60				

Ala	Gly	Ala	Val	Pro	Ser	Glu	His	Pro	Glu	Leu	Pro	Cys	Gly	Ser	Gln
	65				70					75					80

Gln	Leu	Cys	Cys	Gly	Cys	Thr	Ala	Arg	Leu	Gly	Gly	Xaa	Trp	Ile	Gly
				85					90					95	

Ala	Ser	Arg	Cys	Gly	Ser	Gly	Ser	Ala	Phe	Leu	Ala	Ser	Pro	Xaa	
			100					105					110		

<210> 155

<211> 48

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (48)

<223> Xaa equals stop translation

<400> 155

Met Ser Leu Gln Ala Ile Asp Leu Leu Trp⁸⁴ Ser Leu Cys Thr Gln Thr
 1 5 10 15

Ser Leu Leu Thr Leu Ile Cys Ile Cys Ser His Ser Gln Ala Leu Ser
 20 25 30

Ser Ser Pro Gln Leu His Leu Arg Ser Ser Ser Ile Arg Phe Ser Xaa
 35 40 45

<210> 156

<211> 82

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (82)

<223> Xaa equals stop translation

<400> 156

Met Phe His Phe Gly Leu Trp Asp Leu His Phe Phe Leu Ile Val Met
 1 5 10 15

Ala His Arg Asp Asp Cys Ser Phe Lys Gly Gly Cys Gly Leu Leu Glu
 20 25 30

Arg Phe Gln Cys Pro His Thr Ser Phe Ser Ser Ala Ser Gln Lys Arg
 35 40 45

Leu Ala Asp Gly Met Glu Cys Leu Cys Glu Ile Glu Arg Thr Gln Thr
 50 55 60

Arg Ile Arg Lys Ile Cys Leu Pro Thr Leu His Gly His Leu Leu Ala
 65 70 75 80

Val Xaa

<210> 157

<211> 156

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (108)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (113)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (156)
 <223> Xaa equals stop translation

<400> 157
 Met Met Ala Arg Gln Thr Gly Val Phe Tyr Leu Thr Leu Val Leu Ile
 1 5 10 15
 Leu Val Thr Ser Gly Leu Phe Phe Ala Phe Asp Cys Pro Tyr Leu Ala
 20 25 30
 Val Lys Ile Thr Pro Ala Ile Pro Ala Val Ala Gly Ile Leu Phe Phe
 35 40 45
 Phe Val Met Gly Thr Leu Leu Arg Thr Ser Phe Ser Asp Pro Gly Val
 50 55 60
 Leu Pro Arg Ala Thr Pro Asp Glu Ala Ala Asp Leu Glu Arg Gln Ile
 65 70 75 80
 Gly Asn Thr Glu Ser Leu Pro Met Ala Ser Gly His Phe Pro Pro Gly
 85 90 95
 Pro Ser Tyr Ser Gly Glu Gly Arg Pro Arg Ala Xaa Gln Glu Glu Leu
 100 105 110
 Xaa Ala Gly Lys Glu Gly Gly Gln Lys Ser Ala Phe Leu Ser Ser Leu
 115 120 125
 Gly Gly Gln Asp Glu Leu Lys Lys Arg Cys Asp Ile Arg Leu Glu Gly
 130 135 140
 Gln Val Ser Trp Arg Gln Asp Cys Arg Pro Thr Xaa
 145 150 155

<210> 158
 <211> 295
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (295)
 <223> Xaa equals stop translation

<400> 158
 Met Arg Leu Asp Lys Pro Ile Gly Thr Trp Leu Leu Tyr Leu Pro Cys
 1 5 10 15
 Thr Trp Ser Ile Gly Leu Ala Ala Glu Pro Gly Cys Phe Pro Asp Trp
 20 25 30
 Tyr Met Leu Ser Leu Phe Gly Thr Gly Ala Ile Leu Met Arg Gly Ala
 35 40 45

56

Gly Cys Thr Ile Asn Asp Met Trp Asp Gln Asp Tyr Asp Lys Lys Val
 50 55 60

Thr Arg Thr Ala Asn Arg Pro Ile Ala Ala Gly Asp Ile Ser Thr Phe
 65 70 75 80

Gln Ser Phe Val Phe Leu Gly Gly Gln Leu Thr Leu Ala Leu Gly Val
 85 90 95

Leu Leu Cys Leu Asn Tyr Tyr Ser Ile Ala Leu Gly Ala Gly Ser Leu
 100 105 110

Leu Leu Val Ile Thr Tyr Pro Leu Met Lys Arg Ile Ser Tyr Trp Pro
 115 120 125

Gln Leu Ala Leu Gly Leu Thr Phe Asn Trp Gly Ala Leu Leu Gly Trp
 130 135 140

Ser Ala Ile Lys Gly Ser Cys Asp Pro Ser Val Cys Leu Pro Leu Tyr
 145 150 155 160

Phe Ser Gly Val Met Trp Thr Leu Ile Tyr Asp Thr Ile Tyr Ala His
 165 170 175

Gln Asp Lys Arg Asp Asp Val Leu Ile Gly Leu Lys Ser Thr Ala Leu
 180 185 190

Arg Phe Gly Glu Asn Thr Lys Pro Trp Leu Ser Gly Phe Ser Val Ala
 195 200 205

Met Leu Gly Ala Leu Ser Leu Val Gly Val Asn Ser Gly Gln Thr Ala
 210 215 220

Pro Tyr Tyr Ala Ala Leu Gly Ala Val Gly Ala His Leu Thr His Gln
 225 230 235 240

Ile Tyr Thr Leu Asp Ile His Arg Pro Glu Asp Cys Trp Asn Lys Phe
 245 250 255

Ile Ser Asn Arg Thr Leu Gly Leu Ile Val Phe Leu Gly Ile Val Leu
 260 265 270

Gly Asn Leu Trp Lys Glu Lys Lys Thr Asp Lys Thr Lys Lys Gly Ile
 275 280 285

Glu Asn Lys Ile Glu Asn Xaa
 290 295

<210> 159

<211> 60

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (60)

<223> Xaa equals stop translation

27

<400> 159

Met Gly Pro Phe Leu Leu Val Phe Leu Phe Pro Ile Leu Arg Val Cys
 1 5 10 15

Gly Ile Ile Arg Glu Pro Thr Gln Asp Trp Ser Val Leu Leu Glu Arg
 20 25 30

Ala Arg Leu Thr Ala Pro Gly Gln Pro Pro Ala Leu Phe Pro Leu Glu
 35 40 45

Ser Gly Pro Met Ala Thr Ala Gln Asn Thr Ser Xaa
 50 55 60

<210> 160

<211> 122

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (30)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (32)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (87)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (101)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (115)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (122)

<223> Xaa equals stop translation

<400> 160

Met Cys Ser His Ser Thr Leu Ile His Leu Tyr Leu Val Leu Pro Phe
 1 5 10 15

Phe Phe Leu Phe Leu Pro Ser Ser Phe Pro Phe Pro Ser Xaa Ser Xaa
 20 25 30

Ser Ser Ile Leu Pro Ser Leu Arg Leu Pro Pro Phe Phe Pro Pro Ser

35 40 38 45
 Leu Phe Leu His Ser Ser Leu Pro Pro Ser Leu Ser His Pro Leu Gly
 50 55 60
 Leu Ser Ile Thr Ser Ser Arg Gln Ser Phe Leu Asp Tyr His His Leu
 65 70 75 80
 Cys Thr Lys His Leu Ser Xaa Thr Leu Cys Gly Leu Ile Tyr His Cys
 85 90 95
 Leu Asn Ile Phe Xaa Thr Arg Ala Val Met Trp His Met Gln Val Ser
 100 105 110
 Phe Leu Xaa Ile His Trp Leu Leu Pro Xaa
 115 120

<210> 161
 <211> 73
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (73)
 <223> Xaa equals stop translation

<400> 161
 Met Ser Ile Tyr His Val Cys Leu Ile Leu Leu Leu Tyr Ile Thr Ser
 1 5 10 15
 His Ser His Gln Asn Met Ser Ser Cys Leu Gln Val Pro Leu Ser Leu
 20 25 30
 Leu Ser Cys Pro Leu Lys Gly Glu His Leu Ser Gln Phe Ala Gly Asp
 35 40 45
 His Ser Leu Pro Glu Val Arg Asp Arg Asn His His Cys Ile Leu Phe
 50 55 60
 Lys Glu Ser His Gln Lys Arg Lys Xaa
 65 70

<210> 162
 <211> 123
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (123)
 <223> Xaa equals stop translation

<400> 162
 Met Leu Ala Asn Phe Thr Leu Phe Ile Leu Thr Leu Ile Ser Phe Leu
 1 5 10 15

89

Leu Leu Val Cys Ser Pro Cys Lys His Leu Lys Met Met Gln Leu His
 20 25 30

Gly Lys Gly Ser Gln Asp Leu Ser Thr Lys Val His Ile Lys Pro Leu
 35 40 45

Gln Thr Val Ile Ser Phe Leu Met Leu Phe Ala Ile Tyr Phe Leu Cys
 50 55 60

Ile Ile Thr Ser Thr Trp Asn Pro Arg Thr Gln Gln Ser Asn Leu Val
 65 70 75 80

Phe Leu Leu Tyr Gln Thr Leu Ala Ile Met Tyr Pro Ser Phe His Ser
 85 90 95

Phe Ile Leu Ile Met Arg Ser Arg Lys Leu Lys Gln Thr Ser Leu Ser
 100 105 110

Val Leu Cys Gln Val Thr Cys Trp Val Lys Xaa
 115 120

<210> 163

<211> 143

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (143)

<223> Xaa equals stop translation

<400> 163

Met Pro Gly Pro Cys Leu Ser Gln Gln His Pro Phe Leu Ser Leu Ser
 1 5 10 15

Leu Phe Pro Phe Cys Leu Trp Ile Cys Leu Ala Arg Val Pro Gly Val
 20 25 30

Arg Asn Ile Cys Lys Thr Gln Pro Ala Pro Ser Gln Pro Ser Leu Leu
 35 40 45

Gly Leu Gly Leu Ser His Pro Ala Ala Gly Thr Thr Asp Ala Gly Thr
 50 55 60

Gln Ser Leu Pro Arg Ser Gln His Lys Cys Thr Ser Ala Leu Trp Gly
 65 70 75 80

Leu Cys Pro Ala Gln Arg Pro Leu Leu Leu Pro Ala His Ile His Ser
 85 90 95

Ser Gly His Gly Ala Pro Gln Glu Leu Gln Ser His Leu Ser His Arg
 100 105 110

Leu Pro Ala Ser Ala Ser Leu Ser Met Met Ser Pro Phe Ser Glu Ala
 115 120 125

90

Trp	Thr	His	Pro	Ser	Leu	Ser	Leu	Gly	Pro	Ala	Pro	Ser	His	Xaa
130						135					140			

<210> 164

<211> 117

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (46)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (117)

<223> Xaa equals stop translation

<400> 164

Met	Pro	Gly	Gly	Thr	Arg	Cys	Arg	Val	Leu	Leu	Leu	Ser	Leu	Thr	Phe
1				5					10					15	

Gly	Thr	Ser	Met	Ala	Cys	Gly	Asn	Val	Gly	Leu	Arg	Leu	Cys	Pro	Trp
			20					25					30		

Thr	Trp	His	Asn	Trp	Leu	Leu	Pro	Pro	His	Leu	Cys	Ser	Xaa	Trp	Pro
		35					40					45			

Cys	Arg	Arg	Cys	Cys	Trp	Ala	Ala	Ala	Thr	Thr	His	Phe	Ser	Trp	Pro
	50					55					60				

Pro	Trp	Val	Arg	Ser	Ala	Trp	Gly	Pro	Pro	Ala	Ala	Trp	Leu	Glu	Ser
65					70					75				80	

Ser	Gly	His	Pro	Leu	Pro	Ala	Val	Ala	Ser	Cys	Ser	Gln	Pro	Pro	Ala
				85					90					95	

Ser	Ala	Asp	Ser	Ser	Arg	Phe	Ser	Lys	Val	Pro	Cys	Cys	Arg	Arg	Arg
			100					105					110		

Gly	Trp	Thr	Arg	Xaa
				115

<210> 165

<211> 59

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (59)

<223> Xaa equals stop translation

<400> 165

Met	Ser	Val	Cys	Leu	Pro	Leu	His	Leu	Pro	Phe	Leu	Met	Leu	Ala	Lys
1				5					10					15	

Val Ala Thr Ser Phe Cys Arg Trp Gln Leu⁹ Thr Leu Phe Val Ser Thr
 20 25 30

Phe Tyr Lys Asp Ala Leu Val His Thr Val Asn Asp Arg Asn Gln Glu
 35 40 45

Ala Glu Leu Glu Ala Leu Lys Lys Ser Cys Xaa
 50 55

<210> 166
 <211> 126
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (126)
 <223> Xaa equals stop translation

<400> 166
 Met Lys Ala Leu Met Leu Leu Thr Leu Ser Val Leu Leu Cys Trp Val
 1 5 10 15

Ser Ala Asp Ile Arg Cys His Ser Cys Tyr Lys Val Pro Val Leu Gly
 20 25 30

Cys Val Asp Arg Gln Ser Cys Arg Leu Glu Pro Gly Gln Gln Cys Leu
 35 40 45

Thr Thr His Ala Tyr Leu Gly Lys Met Trp Val Phe Ser Asn Leu Arg
 50 55 60

Cys Gly Thr Pro Glu Glu Pro Cys Gln Glu Ala Phe Asn Gln Thr Asn
 65 70 75 80

Arg Lys Leu Gly Leu Thr Tyr Asn Thr Thr Cys Cys Asn Lys Asp Asn
 85 90 95

Cys Asn Ser Ala Gly Pro Arg Pro Thr Pro Ala Leu Gly Leu Val Phe
 100 105 110

Leu Thr Ser Leu Ala Gly Leu Gly Leu Trp Leu Leu His Xaa
 115 120 125

<210> 167
 <211> 87
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (87)
 <223> Xaa equals stop translation

<400> 167

12

Met Phe Leu Val Ala Val Trp Trp Arg Phe Gly Ile Leu Ser Ile Cys
 1 5 10 15

Met Leu Cys Val Gly Leu Val Leu Gly Phe Leu Ile Ser Ser Val Thr
 20 25 30

Phe Phe Thr Pro Leu Gly Asn Leu Lys Ile Phe His Asp Asp Gly Val
 35 40 45

Phe Trp Val Thr Phe Ser Cys Ile Ala Ile Leu Ile Pro Val Val Phe
 50 55 60

Met Gly Cys Leu Arg Ile Leu Asn Ile Leu Thr Cys Gly Ser His Trp
 65 70 75 80

Ala Pro Ile Arg Trp Phe Xaa
 85

<210> 168
 <211> 63
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (16)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (54)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (63)
 <223> Xaa equals stop translation

<400> 168
 Met Val Thr Gly Phe Phe Phe Ile Leu Met Thr Val Leu Trp Phe Xaa
 1 5 10 15

Arg Glu Pro Gly Phe Val Pro Gly Trp Asp Ser Phe Phe Glu Lys Lys
 20 25 30

Gly Tyr Arg Thr Asp Ala Thr Val Ser Val Phe Leu Gly Phe Leu Leu
 35 40 45

Phe Leu Ile Pro Ala Xaa Glu Ala Leu Leu Trp Glu Lys Glu Xaa
 50 55 60

<210> 169
 <211> 48
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (48)
 <223> Xaa equals stop translation

93

<400> 169
 Met Ser Gln Leu Cys Phe Ser Leu Leu Leu Ser Ser Thr Cys His Gly
 1 5 10 15
 Gly Val Ala Ser Leu Leu Thr Ser Asp Leu Ser Ser Gln Ser His Arg
 20 25 30
 Phe Ser Ile Cys Thr Asn Val Asn His Ser Lys Tyr Ser Ser Leu Xaa
 35 40 45

<210> 170
 <211> 137
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (84)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (137)
 <223> Xaa equals stop translation

<400> 170
 Met Leu Phe Ser Leu Arg Glu Leu Val Gln Trp Leu Gly Phe Ala Thr
 1 5 10 15
 Phe Glu Ile Phe Val His Leu Leu Ala Leu Leu Val Phe Ser Val Leu
 20 25 30
 Leu Ala Leu Arg Val Asp Gly Leu Val Pro Gly Leu Ser Trp Trp Asn
 35 40 45
 Val Phe Val Pro Phe Phe Ala Ala Asp Gly Leu Ser Thr Tyr Phe Thr
 50 55 60
 Thr Ile Val Ser Val Arg Leu Phe Gln Asp Gly Glu Lys Arg Leu Ala
 65 70 75 80
 Val Leu Arg Xaa Phe Trp Val Leu Thr Val Leu Ser Leu Lys Phe Val
 85 90 95
 Phe Glu Met Leu Leu Cys Gln Lys Leu Ala Glu Gln Thr Arg Glu Leu
 100 105 110
 Trp Phe Gly Leu Ile Thr Ser Pro Leu Phe Ile Leu Leu Gln Leu Leu
 115 120 125

94

Met Ile Arg Ala Cys Arg Val Asn Xaa
 130 135

<210> 171
 <211> 89
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (40)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (89)
 <223> Xaa equals stop translation

<400> 171
 Met Glu Leu Ser Phe Val Arg Arg Leu Leu Leu Phe Thr Phe Phe Phe
 1 5 10 15
 Ser Thr Phe Ser Pro Pro Pro Pro Thr Pro Cys Leu Glu Gly Leu Met
 20 25 30
 Ser Cys Leu Pro Ser Pro Leu Xaa Lys Asn Thr Ala Gly Ser Gln Thr
 35 40 45
 Lys Ser Leu Arg Glu Ile Gly Thr Gly Ile Ser Asp Thr His Val Ser
 50 55 60
 Pro Ser Pro Ala Gln Ala Pro Leu Cys Ser Arg Ser Pro Thr Trp Asp
 65 70 75 80
 Ser Ser Asp Pro Asn Ser Met Asp Xaa
 85

<210> 172
 <211> 58
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (58)
 <223> Xaa equals stop translation

<400> 172
 Met Thr Met Val Met Glu Gln Val Tyr Leu Met Ser Phe Leu Leu Leu
 1 5 10 15
 Leu Leu Arg Thr Met Met Lys Ala His Trp Thr Tyr Thr Leu Gly Trp
 20 25 30
 Thr Val Leu Phe Leu Thr Ala Leu Pro Asn Pro Val Tyr His Gln Glu

35

40

95

45

Ile Val Trp Thr Tyr Met Lys Arg Ser Xaa
 50 55

<210> 173

<211> 64

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (64)

<223> Xaa equals stop translation

<400> 173

Met Asp Thr Asp Asn Gly Gly Arg His Phe Lys Pro Phe Lys Leu Val
 1 5 10 15

Leu Phe Val Val Leu Leu Ile Lys Ile Leu Leu Ile Leu Ala Lys Thr
 20 25 30

Asn Cys Cys Asp Lys Leu Val Phe Phe Gly Cys Phe Lys His Thr Leu
 35 40 45

Thr Asn Phe Leu Ile Pro Leu Leu Val Pro Pro Ile Val Leu Lys Xaa
 50 55 60

<210> 174

<211> 61

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (61)

<223> Xaa equals stop translation

<400> 174

Met Cys Leu Trp Gly Gln Ala Asn Leu Gly Leu Ile Leu Phe Gln His
 1 5 10 15

Cys Leu Thr Lys Phe Met Gly Gly Tyr Cys Phe Gly Leu Gly Ser Cys
 20 25 30

Thr Arg Pro Leu Arg Asp Gln Thr Lys Met Glu Ser Leu Ile Leu Lys
 35 40 45

Leu Gln Val Thr Glu Pro Lys Leu Ser Cys Phe Ile Xaa
 50 55 60

<210> 175

96

```

<400> 176
Met Leu Arg Val Leu Cys Leu Leu Arg Pro Trp Arg Pro Leu Arg Ala
  1             5             10             15

Arg Gly Cys Ala Ser Asp Gly Ala Ala Gly Gly Ser Glu Ile Gln Val
          20             25             30

Arg Ala Leu Ala Gly Pro Asp Gln Gly Ile Thr Glu Ile Leu Met Asn
          35             40             45

Arg Pro Ser Ala Arg Asn Ala Leu Gly Asn Val Phe Val Ser Glu Leu
  50             55             60

Leu Glu Thr Leu Ala Gln Leu Arg Glu Asp Arg Gln Val Arg Val Leu
  65             70             75             80

```

97

Leu Phe Arg Ser Gly Val Lys Gly Val Phe Cys Ala Gly Ala Asp Leu
 85 90 95

Lys Glu Arg Glu Gln Met Ser Glu Ala Glu Val Gly Val Phe Val Gln
 100 105 110

Arg Leu Arg Gly Leu Met Asn Asp Ile Ala Ala Phe Pro Ala Pro Thr
 115 120 125

Ile Ala Ala Met Asp Gly Phe Ala Leu Gly Gly Gly Leu Glu Leu Ala
 130 135 140

Leu Ala Cys Asp Leu Arg Val Ala Ala Ser Ser Ala Val Met Gly Leu
 145 150 155 160

Ile Glu Thr Thr Arg Gly Leu Leu Pro Gly Ala Gly Gly Thr Gln Arg
 165 170 175

Leu Pro Arg Cys Leu Gly Val Ala Leu Ala Lys Glu Leu Ile Phe Thr
 180 185 190

Gly Arg Arg Leu Ser Gly Thr Glu Ala His Val Leu Gly Leu Val Asn
 195 200 205

His Ala Val Ala Gln Asn Glu Glu Gly Asp Ala Ala Tyr Gln Arg Ala
 210 215 220

Arg Ala Leu Ala Gln Glu Ile Leu Pro Gln Ala Pro Ile Ala Val Arg
 225 230 235 240

Leu Gly Lys Val Ala Ile Asp Arg Gly Thr Glu Val Asp Ile Ala Ser
 245 250 255

Gly Met Ala Ile Glu Gly Met Cys Tyr Ala Gln Asn Ile Pro Thr Arg
 260 265 270

Asp Arg Leu Glu Gly Met Ala Ala Phe Arg Glu Lys Arg Thr Pro Lys
 275 280 285

Phe Val Gly Lys Xaa
 290

<210> 177

<211> 46

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (46)

<223> Xaa equals stop translation

<400> 177

Met Leu Ser Ser Leu Tyr Leu Leu Leu Met Pro Pro Tyr Lys Phe Thr
 1 5 10 15

Gly Glu Leu His Pro Pro Val Ala Ala Thr Cys Leu Leu Thr Val Leu

20

25

98

30

Leu Gly Cys Leu Ile Gly Val Ser Ser Asp Gly Trp Ile Xaa
 35 40 45

<210> 178

<211> 47

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (47)

<223> Xaa equals stop translation

<400> 178

Met Cys Ile Pro Glu Ala Leu Gly Lys Asn Ser Leu Phe Leu Ser Ser
 1 5 10 15

Thr Phe Leu Trp Leu Leu Ala Phe Phe Gly Leu Trp Ser His His Ser
 20 25 30

Tyr Leu Glu Gly Gln His Leu Gln Ile Cys Phe Phe Phe Thr Xaa
 35 40 45

<210> 179

<211> 55

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (55)

<223> Xaa equals stop translation

<400> 179

Met Thr Thr Ser Leu Phe Gly Leu Val Cys Val Val Cys Gln Gly Ala
 1 5 10 15

Gly Val Ser Ala Phe Thr Gln Val Asn Leu Phe Ser Phe Ser Leu Val
 20 25 30

Ile Val Lys Lys Gln Asn Lys Thr Ser Cys Glu Pro Phe Gly Thr Ser
 35 40 45

Gly Lys Val Pro Leu Leu Xaa
 50 55

<210> 180

<211> 67

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

99

<222> (67)

<223> Xaa equals stop translation

<400> 180

Met	Leu	Ile	Tyr	Trp	Leu	Gln	Ser	Ser	Phe	Ile	Leu	Ser	Ala	Phe	Val
1				5					10					15	

Leu	Ile	Asn	Ser	Pro	Val	Thr	Thr	Gly	Ile	Gln	Lys	Ser	Cys	Cys	Lys
			20					25					30		

Phe	Phe	Pro	Val	Ser	Ile	Asn	Leu	Cys	Phe	Ala	Ser	Leu	His	Arg	Met
		35					40					45			

Lys	Val	Val	Thr	Leu	Val	Ala	Leu	Gln	Trp	Leu	Asn	Ile	Ala	Leu	Arg
	50						55				60				

Ser	Ser	Xaa
65		

<210> 181

<211> 51

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (51)

<223> Xaa equals stop translation

<400> 181

Met	Val	Cys	Cys	Gly	Phe	Phe	Leu	Leu	Trp	Ser	Arg	Val	Arg	Ser	Tyr
1				5					10					15	

Met	Lys	Leu	Ser	Gly	His	Arg	Trp	Ser	Ser	Ser	Cys	Pro	His	His	Cys
			20					25					30		

Tyr	Ser	Lys	Cys	Gly	Leu	His	Thr	Ser	Asn	Gly	Lys	Ser	Ser	Val	His
		35					40					45			

Thr	Val	Xaa
50		

<210> 182

<211> 91

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (29)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (30)

<223> Xaa equals any of the naturally occurring L-amino acids

100

<220>

<221> SITE

<222> (65)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (91)

<223> Xaa equals stop translation

<400> 182

Met	Leu	Arg	Cys	Ser	Phe	Ser	Ser	Phe	Leu	Leu	Cys	His	Thr	Ile	Leu
1				5					10					15	

Leu	Phe	Leu	Gly	Ser	Ser	Ala	His	Leu	Leu	Val	Glu	Xaa	Xaa	Val	Trp
			20					25						30	

Gly	Leu	Tyr	Glu	Tyr	Arg	Ile	Gly	Asp	Met	Val	Asp	Gln	Lys	Ala	Thr
		35					40					45			

Phe	Cys	Val	Gln	Lys	Gln	Glu	Cys	Leu	Phe	Pro	Leu	Gly	Ser	Trp	Val
	50					55					60				

Xaa	Arg	Val	Glu	Gly	Gly	Ala	Phe	Ala	Arg	Glu	Pro	Pro	Ser	Ser	Thr
65					70					75					80

Gln	Tyr	Phe	Pro	Val	Ser	Cys	Leu	Tyr	Gln	Xaa
				85					90	

<210> 183

<211> 55

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (55)

<223> Xaa equals stop translation

<400> 183

Met	Ser	Ala	Leu	Leu	Ser	His	His	Val	Pro	Leu	Phe	Tyr	Leu	Thr	Gly
1				5					10					15	

Cys	Leu	Phe	Ser	Leu	Leu	Ala	Ser	Trp	Asp	Cys	Asn	Gly	Lys	Glu	Gly
			20					25					30		

Ala	Gly	Arg	Ala	Ile	Lys	Gly	Lys	Asn	Asn	Thr	Trp	Asn	Cys	Met	Ile
		35					40					45			

Leu	Ser	Lys	Val	Lys	Phe	Xaa
	50					55

<210> 184

<211> 64

<212> PRT

<213> Homo sapiens

101

<220>

<221> SITE

<222> (26)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (41)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (64)

<223> Xaa equals stop translation

<400> 184

Met	Val	His	Lys	Ala	Ile	Leu	Ala	Leu	Leu	Pro	Trp	Gly	Phe	Ser	Ala
1				5					10					15	

Asp	Glu	Leu	Leu	Ala	Ser	Leu	Met	Met	Xaa	Leu	Thr	Glu	Lys	Tyr	Gln
		20						25					30		

Asn	Cys	Ser	Ser	Thr	Thr	Asp	Ile	Xaa	Asn	Gln	Gln	Leu	Arg	Ser	Leu
		35					40					45			

Gly	Gln	Asn	Phe	Met	Phe	Gln	Gln	Asn	Leu	Gln	Leu	Ile	Leu	Met	Xaa
	50					55					60				

<210> 185

<211> 113

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (113)

<223> Xaa equals stop translation

<400> 185

Met	Met	Thr	Ser	Ser	Leu	Gly	Leu	Ser	Phe	Leu	Leu	Asn	Leu	Ile	Leu
1				5					10					15	

Gly	Met	Lys	Phe	Thr	Tyr	Leu	Ile	Pro	Gln	Asn	Lys	Tyr	Ile	Gln	Leu
			20					25					30		

Phe	Thr	Thr	Ile	Leu	Ser	Phe	Phe	Ser	Gly	Val	Leu	Ser	Leu	Leu	Glu
		35				40						45			

Cys	Lys	Leu	Ser	Thr	Ser	Ser	Cys	Thr	Cys	Leu	Asn	Ile	His	Lys	Ser
	50					55					60				

Asp	Asn	Glu	Cys	Lys	Glu	Ser	Glu	Asn	Ser	Ile	Glu	Asp	Ile	Ser	Leu
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

Xaa

```

<400> 186
Met Leu His Leu Thr Leu Tyr Leu His Phe Ile Leu Phe Val Phe Pro
  1             5             10             15

Ile Thr Ser Asn Phe Ser Ser Leu His Pro Phe Leu Phe Ile Ser Ser
      20             25             30

Gln Phe Thr Ser Cys Cys Gln Ile Asn Phe Pro Asn Ala Gln Ala Leu
      35             40             45

Ser Tyr His Glu Phe Leu Ile Ala Thr Tyr Asp Xaa
      50             55             60

```

```

<400> 187
Met Pro Cys Ile Arg Gly Val Phe His Cys Phe Ile Leu Ile Ile Leu
  1             5             10             15

Ile Leu Leu Ala Ser His Ala Phe Ser Gly Ser Gly Asn Gln Arg Leu
      20             25             30

Lys Glu Ala Leu Thr Leu Ile Val Ser Val Asn Val Asp Ile Ala Arg
      35             40             45

His Arg Pro Phe Leu Glu Arg Ile His Val Lys Lys Gly Asn Thr Xaa
  50             55             60

```

103

<210> 188
 <211> 71
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (71)
 <223> Xaa equals stop translation

<400> 188
 Met Phe Ser Arg Leu His Phe Leu Thr His Ser Leu Ser Leu Leu His
 1 5 10 15
 Leu Pro Ser Gln Val Phe Gly Glu Val His Ser Ser Cys Val Ser Ser
 20 25 30
 Leu Pro Cys Pro Asp Thr Pro Ala Leu Pro Tyr Cys Pro Ser Phe Leu
 35 40 45
 Arg Tyr Asp Asp His Ile Glu Ala Gln Pro Leu Lys His Ile Asn Thr
 50 55 60
 Asn Asp His Ile Ser Ile Xaa
 65 70

<210> 189
 <211> 63
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (63)
 <223> Xaa equals stop translation

<400> 189
 Met Cys Val Phe Ser Ser Asp Ser Ile Pro Ser Leu Leu Ile Leu Leu
 1 5 10 15
 Val Leu Cys His Ser Val Cys Cys Leu Lys Leu Phe Phe Lys Leu Ile
 20 25 30
 Phe Pro Asn Ser Phe Ile Ile Tyr Glu Lys Leu Gly Leu Asn His Phe
 35 40 45
 Ala Tyr His Leu Ser Gly Trp Phe Glu Leu Ser Leu Asp Thr Xaa
 50 55 60

<210> 190
 <211> 193

<212> PRT

104

<213> Homo sapiens

<220>

<221> SITE

<222> (193)

<223> Xaa equals stop translation

<400> 190

Met Glu Ala Leu Leu Gln Ser Leu Val Ile Val Leu Leu Gly Phe Lys
 1 5 10 15

Ser Phe Leu Ser Glu Glu Leu Gly Ser Glu Val Leu Asn Leu Leu Thr
 20 25 30

Asn Lys Gln Tyr Glu Leu Leu Ser Lys Asn Leu Arg Lys Thr Arg Glu
 35 40 45

Leu Phe Val His Gly Leu Pro Gly Ser Gly Lys Thr Ile Leu Ala Leu
 50 55 60

Arg Ile Met Glu Lys Ile Arg Asn Val Phe His Cys Glu Pro Ala Asn
 65 70 75 80

Ile Leu Tyr Ile Cys Glu Asn Gln Pro Leu Lys Lys Leu Val Ser Phe
 85 90 95

Ser Lys Lys Asn Ile Cys Gln Pro Val Thr Arg Lys Thr Phe Met Lys
 100 105 110

Asn Asn Phe Glu His Ile Gln His Ile Ile Ile Asp Asp Ala Gln Asn
 115 120 125

Phe Arg Thr Glu Asp Gly Asp Trp Tyr Gly Lys Ala Lys Phe Ile Thr
 130 135 140

Gln Thr Ala Arg Asp Gly Pro Gly Val Leu Trp Ile Phe Leu Asp Tyr
 145 150 155 160

Phe Gln Thr Tyr His Leu Ser Cys Ser Ala Ser Pro Leu Pro Gln Thr
 165 170 175

Ser Ile Gln Glu Lys Arg Ser Thr Glu Trp Ser Ala Met Gln Val Gln
 180 185 190

Xaa

<210> 191

<211> 112

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (112)

<223> Xaa equals stop translation

105

<400> 191

Met Gln Phe Ser Leu Cys Leu Thr Ala Val Phe Leu Leu Gln Leu Ala
 1 5 10 15

Ala Gly Ile Leu Gly Phe Val Phe Ser Asp Lys Ala Arg Gly Lys Val
 20 25 30

Ser Glu Ile Ile Asn Asn Ala Ile Val His Tyr Arg Asp Asp Leu Asp
 35 40 45

Leu Gln Asn Leu Ile Asp Phe Gly Gln Lys Lys Val Trp Val Ser Gln
 50 55 60

Trp Ser Gly Gly Leu Trp Val Lys Val Asn Val Ile Pro Arg Asp Ala
 65 70 75 80

Ser Pro Ser Met Pro Val Gly Leu Phe Ile Thr Cys Gln Val Met Ala
 85 90 95

Ser Gly Lys Gly Phe Gly Lys Lys Ser Thr Arg Ser Arg Val Leu Xaa
 100 105 110

<210> 192
 <211> 80
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (80)
 <223> Xaa equals stop translation

<400> 192

Met Cys Arg Pro Leu Leu Pro Leu Leu Phe Pro Trp Gly His Cys Leu
 1 5 10 15

Ser Ile Pro Leu Cys Lys Trp Pro Gln Ile Met Ser Gln Pro Pro Arg
 20 25 30

Leu His Arg Leu Leu Ala Ser Gly Pro Ser Thr Lys Lys His Ser Lys
 35 40 45

Leu Gln Thr His Ser Trp Glu Asn Ser Asn Gly Leu Thr Leu Pro Phe
 50 55 60

Glu Pro Ala Arg Ser His Gly Leu Trp Arg Ala Ala Phe Glu Ser Xaa
 65 70 75 80

<210> 193

<211> 88
 <212> PRT
 <213> Homo sapiens

106

<220>
 <221> SITE
 <222> (88)
 <223> Xaa equals stop translation

<400> 193

Met Leu Ser Ile Ile Asp Leu Leu Phe Leu Leu Ser Pro Thr Phe Gly
 1 5 10 15

Leu Ile Thr Glu Leu Leu Phe Ser Pro Glu Val Pro Lys Ala Leu Ser
 20 25 30

Cys Pro Leu Lys Ala Leu Gly Gly Gly Ser His Ser His Glu Pro Leu
 35 40 45

Gly Met Phe Ala Pro Val Pro Pro Gly Cys Glu Ser Ser Thr Pro Phe
 50 55 60

Pro Lys Gly Leu Gly Ala Ser Lys Ile Leu Thr Leu Gly Ala Gln Ala
 65 70 75 80

Glu Phe Arg Arg Arg Ser His Xaa
 85

<210> 194
 <211> 42
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (42)
 <223> Xaa equals stop translation

<400> 194

Met Glu Asp His Phe Leu Ile Gly His Phe Pro Phe Phe Phe Leu Phe
 1 5 10 15

Ser Phe Pro Cys Phe Cys Thr Lys Pro Leu Cys Arg Glu Tyr Phe Leu
 20 25 30

Ile Cys Ser Ile Gln Asp Glu Ser Lys Xaa
 35 40

<210> 195
 <211> 69
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (69)

<223> Xaa equals stop translation 107

<400> 195

Met Phe Asn Leu Pro Lys Pro Val Phe Leu Ser Trp Trp Arg Trp Lys
1 5 10 15

Thr Ile Val Ile Phe Leu Ala Cys Leu Ala Ser Ala Ala Ile Lys Glu
20 25 30

Thr Ala Val Ser Met Lys Thr Val Phe Pro Ile Phe Val Gln Ile Thr
35 40 45

Leu Ile Leu Leu Leu Glu Ser Arg Val Leu Lys Ile Gly Asp Phe Ser
50 55 60

Asn Phe Phe Cys Xaa
65

<210> 196

<211> 153

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (66)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (77)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (81)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (84)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (86)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (87)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (93)

<223> Xaa equals any of the naturally occurring L-amino acids

108

<220>
 <221> SITE
 <222> (103)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (110)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (153)
 <223> Xaa equals stop translation

<400> 196
 Met Asp His Ser Pro Thr Thr Gly Val Val Thr Val Ile Val Ile Leu
 1 5 10 15
 Ile Ala Ile Ala Ala Leu Gly Ala Phe Asp Pro Gly Leu Leu Val Leu
 20 25 30
 Pro Ala Ala Ala Ala His Gln Pro Val Arg Gly Arg Gly Glu His Arg
 35 40 45
 Gly Gly Trp Gly Asp Gln Gly Thr Leu Pro Ala Gly Ala Val Phe Gly
 50 55 60
 Gln Xaa Thr Val Arg Gly Glu Lys Gly Gln Ala Asp Xaa Ser Gln Thr
 65 70 75 80
 Xaa Arg Lys Xaa Thr Xaa Xaa Pro Gly Cys Lys Gly Xaa Leu Val Pro
 85 90 95
 Val Cys Lys Pro Ala Lys Xaa Gly Leu Gly Gly Ala Lys Xaa Ile Arg
 100 105 110
 Met Arg Cys Cys Leu Arg Gly Arg Ala Asp Thr Cys Trp His Gly Leu
 115 120 125
 Cys Gly Phe Arg Pro Ser His Ala Leu Met Pro Gly Asp Leu Ala Val
 130 135 140
 Leu Gly Phe Pro Ser Ala Ser Arg Xaa
 145 150

<210> 197
 <211> 63
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (63)
 <223> Xaa equals stop translation

<400> 197

109

Met Lys Asn Ser Thr Ser Leu Leu Tyr Lys Leu Phe Ser Ser Leu Ser
 1 5 10 15

Val Phe Ile Phe Lys Phe Leu Leu Leu Phe Tyr Thr Leu His Ile Ala
 20 25 30

Leu Gly Val Lys Ile Gln Tyr Lys Pro Leu Ala His Phe Ile Asp His
 35 40 45

Ser Cys Ile Gln Gln Val Ser Gln Val Gln Trp Ser Ile Pro Xaa
 50 55 60

<210> 198

<211> 64

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (64)

<223> Xaa equals stop translation

<400> 198

Met Gln Glu Pro His Gly Lys Phe Leu Ser Trp Gly Arg Trp Leu Trp
 1 5 10 15

Trp Trp Ser Leu Ala Ala Pro Ala Leu Val Gln Ala Val Asn Met Pro
 20 25 30

Pro Ala Tyr Ile Gln Ile Glu Asn Trp Tyr Met Met Leu Leu Met Gly
 35 40 45

Trp Glu Thr Lys Cys Cys His Val Arg Ser Leu Trp Val Gly Thr Xaa
 50 55 60

<210> 199

<211> 43

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (43)

<223> Xaa equals stop translation

<400> 199

Met Leu Ile Asn Cys Ile Phe Ser Leu Leu Leu Leu Ser His Ala
 1 5 10 15

Asp Gly Met His Leu Phe Ile Ser Ser Gly Asp Arg Ile Leu Phe Cys
 20 25 30

110

Leu Tyr Phe Leu His Ser Arg Val Cys Ala Xaa
 35 40

<210> 200
 <211> 41
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (41)
 <223> Xaa equals stop translation

<400> 200
 Met Ser Val Tyr Val Asn Ile Met His Ile Val Ile Tyr Ile Tyr Leu
 1 5 10 15

Cys Val Tyr Met Cys Val Ala Gln Ser His Thr His Thr Gln Ile Cys
 20 25 30

Ile Gln Met Leu Pro Gly Leu Gln Xaa
 35 40

<210> 201
 <211> 44
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (44)
 <223> Xaa equals stop translation

<400> 201
 Met Ile Leu Ser Phe Leu Met Leu Phe Leu Ile Val Lys Thr Ile Pro
 1 5 10 15

Leu Ile Leu Ala Tyr Cys Tyr Asn Ser Ile Ser Phe Phe Ser Asn Asn
 20 25 30

Leu Val Leu Val Lys Met Gly Tyr Asn Asn Lys Xaa
 35 40

<210> 202
 <211> 42
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (42)
 <223> Xaa equals stop translation

<400> 202
 Met Arg Leu Leu Ser Thr Leu Leu Ser Phe Tyr Pro Phe Ser Asn Cys

<220>
<221> SITE
<222> (113)

<223> Xaa equals stop translation

112

<400> 204

Met Ala Pro Trp Leu Pro Leu Leu Ser Leu Leu Gly Leu Leu Leu Gly
 1 5 10 15

Xaa Ala Pro Ala Pro Pro Arg Arg Ala Ala Asp Ala Gln Ala Arg Glu
 20 25 30

Ala Ala Tyr Pro Glu Leu Leu Gly Pro Ala Arg Phe Ala Leu Glu Met
 35 40 45

Tyr Asn Arg Gly Arg Ala Ala Gly Xaa Arg Ala Thr Leu Gly Ala Val
 50 55 60

Arg Gly Arg Val Arg Arg Ala Gly Glu Gly Ser Leu Tyr Ser Leu Arg
 65 70 75 80

Ala Thr Leu Glu Glu Pro Pro Cys Asn Xaa Xaa Thr Val Cys Gln Leu
 85 90 95

Pro Val Ser Lys Arg Pro Cys Ser Ala Ala Leu Lys Ser Trp Thr Ser
 100 105 110

Xaa

<210> 205

<211> 45

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (45)

<223> Xaa equals stop translation

<400> 205

Met Pro Thr Trp Pro Leu Leu Gln Leu Leu Ser Cys Ser Phe Pro Ser
 1 5 10 15

Leu Leu Cys Glu Thr Phe Thr Phe Cys Ser Lys Asp Glu Val Ser Arg
 20 25 30

Trp Lys Ala Gly Cys Phe Val Pro Leu Pro Ala Ser Xaa
 35 40 45

<210> 206

<211> 71

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (71)

<223> Xaa equals stop translation

<400> 206 113
 Met Thr His Trp Ser Gly Cys Ala Ala Leu Tyr Leu Ile Phe Leu Ser
 1 5 10 15
 Leu Lys Leu Ala Phe Gln Ala Gly Ala Gly Arg Gly Ala Gln Val Gly
 20 25 30
 Ser Val Leu Pro Pro Ser Gly Gly Ala Val Val Val Asp Gln Tyr Cys
 35 40 45
 Cys Arg Leu Ser Ala Gln Thr Tyr Phe Ser Leu Pro Ala Leu Gln Lys
 50 55 60
 Cys Ile Gly Ile Cys Arg Xaa
 65 70

<210> 207
 <211> 42
 <212> PRT
 <213> Homo sapiens

 <220>
 <221> SITE
 <222> (42)
 <223> Xaa equals stop translation

<400> 207
 Met Ile Asn Phe Trp Pro Val Thr His Val Cys Ile Trp Leu Leu Trp
 1 5 10 15
 Leu Gln Ala Leu Glu Ala Arg Gly Gln Gly Ser Asn Ile Asp Cys Thr
 20 25 30
 Arg Asn Ser Lys Thr Val Phe Thr Ser Xaa
 35 40

<210> 208
 <211> 51
 <212> PRT
 <213> Homo sapiens

 <220>
 <221> SITE
 <222> (51)
 <223> Xaa equals stop translation

<400> 208
 Met Tyr Ile Tyr Leu Ile His Leu Cys Met Cys Val Tyr Ile Tyr Ile
 1 5 10 15
 Tyr Ile Leu Leu Ile Ile Tyr Thr Leu Asp Pro Glu Pro Pro Ser Trp
 20 25 30
 Ser Pro Lys Leu Asp Ser His Leu Ser Leu Arg Gln Pro Ser Asn Asp
 35 40 45

Arg Phe Xaa
50

114

<210> 209
<211> 65
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (65)
<223> Xaa equals stop translation

<400> 209
Met Phe Val Leu Cys Thr Arg Ala Val Arg Thr Arg Leu Phe Ser Leu
1 5 10 15
Cys Cys Cys Cys Cys Ser Ser Gln Pro Pro Thr Lys Ser Pro Ala Gly
20 25 30
Thr Pro Lys Ala Pro Ala Pro Ser Lys Pro Gly Glu Ser Gln Glu Ser
35 40 45
Gln Gly Thr Pro Gly Glu Leu Pro Ser Thr Trp Ser Phe Cys Pro Phe
50 55 60
Xaa
65

<210> 210
<211> 77
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (77)
<223> Xaa equals stop translation

<400> 210
Met Leu Ala Leu Leu Val Gly Gly Leu Val Ala Ala Leu Ala Cys His
1 5 10 15
Gly Ile Leu Ala Ala Ile Leu Ala Val Cys Gly Glu Leu Val Ser Gly
20 25 30
Lys Gly Thr Arg Ser Ser Asp Glu Asp Asp Gly Gly Asp Gly Asp Arg
35 40 45
Gly His Arg Gly Leu Ser Leu Leu Asn Ser Ala Phe Gly His Met Gly
50 55 60
Asp Gly Asp Arg Lys Asp Asp Asn Ser Gly Thr Leu Xaa
65 70 75

<210> 211
 <211> 45
 <212> PRT
 <213> Homo sapiens

115

<220>
 <221> SITE
 <222> (45)
 <223> Xaa equals stop translation

<400> 211
 Met Phe Val Gly Thr Arg Val Leu Leu Val Pro Leu Pro Phe Phe Ser
 1 5 10 15
 Ile Ser Gly Met Leu Ala Ile Asp Lys Tyr Leu His Lys Lys Leu Leu
 20 25 30
 Leu Asn Glu Ile Ile Thr Thr Ser Thr Trp Ala Leu Xaa
 35 40 45

<210> 212
 <211> 66
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (27)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (66)
 <223> Xaa equals stop translation

<400> 212
 Met Gly Lys Gly His Gln Arg Pro Trp Trp Lys Val Leu Pro Leu Ser
 1 5 10 15
 Cys Phe Leu Val Ala Leu Ile Ile Trp Cys Xaa Leu Arg Glu Glu Ser
 20 25 30
 Glu Ala Asp Gln Trp Leu Arg Gln Val Trp Gly Glu Val Pro Glu Pro
 35 40 45
 Ser Asp Arg Ser Glu Glu Pro Glu Thr Pro Ala Ala Tyr Arg Ala Arg
 50 55 60
 Thr Xaa
 65

<210> 213
 <211> 62
 <212> PRT
 <213> Homo sapiens

116

<220>
 <221> SITE
 <222> (62)
 <223> Xaa equals stop translation

<400> 213
 Met Arg Leu Cys Thr Thr Trp Met Ala Val Lys Phe Leu Trp Trp Gly
 1 5 10 15
 Met Thr Trp Ile Pro Ser Gly Lys Ala Cys Ser Trp Thr Gln Pro Leu
 20 25 30
 Cys Ser Ser Gly Gly Trp Ser Ser Pro Thr His Leu Pro Thr Ser Leu
 35 40 45
 Leu Leu Gly Trp Arg Ala Ser Leu Cys Met Lys Arg Ser Xaa
 50 55 60

<210> 214
 <211> 56
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (56)
 <223> Xaa equals stop translation

<400> 214
 Met Phe Ala Ser Tyr His Ile Gln Phe Phe Thr Trp Leu Ile Gln Lys
 1 5 10 15
 Leu Ser Leu Val Trp Lys Ser Val Val Ala Ile Arg Glu Gln Gly Lys
 20 25 30
 Glu Leu Val Trp Lys Gln His Leu Pro Leu Arg Ser Tyr Ser Pro Asn
 35 40 45
 Asn Ala Lys Ser Leu Gly Leu Xaa
 50 55

<210> 215
 <211> 213
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (88)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (213)
 <223> Xaa equals stop translation

<400> 215

117

Met Leu Ser Phe Asn Phe Thr Trp Met Val Trp Val Ser Leu Val Leu
 1 5 10 15

Lys Ser Gln Arg Ala Lys Leu Ala Leu His Ser Leu His Leu His Gln
 20 25 30

Glu Val Arg Leu Arg Met Ser Arg Arg Glu Ser Pro Gly Arg Pro Leu
 35 40 45

Arg Cys Gly Val Arg Gly Asn Met Gly Ala Arg Thr Pro Val Pro Thr
 50 55 60

Ala Asp Tyr Pro Ser Pro Tyr Arg Thr Leu Pro Arg Met Ala Ala Pro
 65 70 75 80

Pro Pro Gln Lys Ser Ser Cys Xaa Arg Leu His Arg Pro His Trp Trp
 85 90 95

Arg Pro Arg Thr Pro Ser Ser Glu Lys Thr Gly Gly Gln Ser Gln Ser
 100 105 110

Thr Leu Asp Arg Cys Ala His Leu Val His Met Leu Leu Arg Asp Gln
 115 120 125

Arg Ala Thr Ser Gln Trp Lys Ala Gly Gly Arg Leu Cys Arg Ala Leu
 130 135 140

Ser Lys Thr Pro Leu Gln His Gln Leu His Ser Thr Ser Tyr Arg Lys
 145 150 155 160

Ala Leu Pro Ile Leu Arg Pro Ser Ser Arg Arg Glu Ala Gly Pro Leu
 165 170 175

His His Ile Asp Leu Arg Arg Cys Phe Ser Arg Leu Gly Arg Gly Ala
 180 185 190

Asp Phe Ala Val Cys Ala Lys Glu Pro Val Ser Asp Asn Pro Ile Phe
 195 200 205

Leu Leu Ile Thr Xaa
 210

<210> 216

<211> 41

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (41)

<223> Xaa equals stop translation

<400> 216

Met Asn Met Phe Gln Thr Ile Leu Val Cys Val Leu Phe Val Phe Val
 1 5 10 15

118

Arg Trp Phe Phe Leu Leu Leu Gln Ile Glu Ser Ile Gln Thr Lys Phe
 20 25 30

His Cys Ile Ser Ser Gln Phe Trp Xaa
 35 40

<210> 217
 <211> 60
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (60)
 <223> Xaa equals stop translation

<400> 217
 Met Glu Leu Val Trp Phe Arg Phe Leu His Leu Asn Leu Leu Pro Arg
 1 5 10 15

Gly Val Cys Cys Gly Ile Cys Val Cys Val Arg Arg Gly Met Val Leu
 20 25 30

Ser Glu Pro Thr Ser Cys Gly Gln Arg Ala Leu Ser Cys Glu Gly Gly
 35 40 45

Cys His Ser Gly Arg Val Gln Phe Arg Arg Pro Xaa
 50 55 60

<210> 218
 <211> 59
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (59)
 <223> Xaa equals stop translation

<400> 218
 Met Arg Arg Met Arg Met Lys Ser Leu Ser Pro Arg Arg Ser Trp Trp
 1 5 10 15

Thr Leu Trp Leu Gly Gln Gly Val Leu Gly Ala Ala Leu Lys Ala Asn
 20 25 30

Thr Leu Trp Ile Ala Met Arg Arg Arg Met Met Met Met Gly Gly Pro
 35 40 45

Ala Asn Met Thr Ser Trp Pro Gln Arg Met Xaa
 50 55

<210> 219
 <211> 46

<212> PRT
 <213> Homo sapiens

119

<220>
 <221> SITE
 <222> (46)
 <223> Xaa equals stop translation

<400> 219
 Met Pro Phe Phe Leu Leu Thr Phe Pro Leu Val Leu Tyr Pro His Leu
 1 5 10 15
 Ser Arg Gly Ser Asp Pro Val Leu Pro Cys Val Met Gly Ile His Val
 20 25 30
 Phe Gly Leu Ser His His Ser Arg Lys Val Ala Pro Pro Xaa
 35 40 45

<210> 220
 <211> 62
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (62)
 <223> Xaa equals stop translation

<400> 220
 Met Asp Arg Val Arg Phe Arg Ser Trp Leu Leu Tyr Pro Cys Cys Val
 1 5 10 15
 Ala Leu Gly Gln Glu Leu Gly Leu Ser Ala Pro Gln Trp Leu Ile Thr
 20 25 30
 Glu Asn Gly Met Pro Ala Leu Ala Leu Val Gly Cys Phe Glu Pro Thr
 35 40 45
 Ala Gly Ser Gly Ser Ser Trp His Asp Val Phe Leu Pro Xaa
 50 55 60

<210> 221
 <211> 52
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (52)
 <223> Xaa equals stop translation

<400> 221
 Met Lys Leu Asn Val His Phe Leu Trp Cys Thr Phe Ile Phe Gln Thr
 1 5 10 15
 Ser Gly Ser His Ile Glu Leu Leu Ile Ser Gly Gln Val Ser Ser Tyr

20 25 30
 Ile Pro Ser Leu Asp Phe Cys Thr His Lys Val Val Ser Arg Glu Lys
 35 40 45

Phe Glu Glu Xaa
 50

<210> 222
 <211> 51
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (51)
 <223> Xaa equals stop translation

<400> 222
 Met Ala Ser Pro Val Phe Lys Thr Phe Trp Arg Leu Glu Leu Ser Val
 1 5 10 15

Pro Leu Ser Leu Leu Phe Ile Leu Gln Ile Val Thr Ser Leu Ser Ser
 20 25 30

Asp Glu Ile Cys Tyr Ser Thr Arg Lys Val Phe Ile Ile Arg Arg Gln
 35 40 45

Leu Tyr Xaa
 50

<210> 223
 <211> 47
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (47)
 <223> Xaa equals stop translation

<400> 223
 Met Cys Met Cys Val Gly Val Cys Leu Ile Thr Leu Leu Asp Arg Phe
 1 5 10 15

Leu Trp Phe Gly Thr Ala Gly Ala Lys Phe Ile Gln Lys Ser Thr Phe
 20 25 30

Leu Ser Lys Leu Pro Met Thr Leu Val Ser Phe His Ser Ile Xaa
 35 40 45

<210> 224
 <211> 52
 <212> PRT
 <213> Homo sapiens

121

<220>
 <221> SITE
 <222> (52)
 <223> Xaa equals stop translation

<400> 224
 Met Cys Pro Phe His Lys Ala Tyr Leu Asp Cys Phe Phe Gln Ile Ser
 1 5 10 15
 Leu Leu Leu Leu Ile Phe Leu Thr Tyr Leu Asp Ile Gly Lys Cys Gly
 20 25 30
 Leu Trp Ser His Glu Trp Arg Ile Arg Glu Leu Gly Lys His Glu Arg
 35 40 45
 Trp Trp Asn Xaa
 50

<210> 225
 <211> 66
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (61)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (66)
 <223> Xaa equals stop translation

<400> 225
 Met Asn Gln Pro Ile Leu Arg Ser Gln Ala Leu Leu Trp Pro Trp Arg
 1 5 10 15
 Trp Val Val Lys Ala Lys Pro Cys Val Cys Val Ser Met Asp Ala Trp
 20 25 30
 Ile Pro Asp Arg Ser Gln His Cys Pro Ser Ile Pro Gly Gln Lys Lys
 35 40 45
 Glu Arg Ala Gly Ser His Gly His Gln Ala Leu Ala Xaa Leu Leu Phe
 50 55 60
 Leu Xaa
 65

<210> 226
 <211> 47
 <212> PRT
 <213> Homo sapiens

<220>

<221> SITE
 <222> (47)
 <223> Xaa equals stop translation

122

<400> 226
 Met Ala Ser Arg Gly Thr Ala Ala Pro Gly Arg Thr Phe Leu Ala Met
 1 5 10 15
 Met Val Thr Ser Phe Phe Phe Cys Met Arg Trp Gly Ser Trp Ala Glu
 20 25 30
 Gln Met Pro Gln Arg Cys Leu Pro Cys Cys Met Gln Glu Cys Xaa
 35 40 45

<210> 227
 <211> 222
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (184)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (222)
 <223> Xaa equals stop translation

<400> 227
 Met Ala Gly Gly Val Arg Pro Leu Arg Gly Leu Arg Ala Leu Cys Arg
 1 5 10 15
 Val Leu Leu Phe Leu Ser Gln Phe Cys Ile Leu Ser Gly Gly Glu Ser
 20 25 30
 Thr Glu Ile Pro Pro Tyr Val Met Lys Cys Pro Ser Asn Gly Leu Cys
 35 40 45
 Ser Arg Leu Pro Ala Asp Cys Ile Asp Cys Thr Thr Asn Phe Ser Cys
 50 55 60
 Thr Tyr Gly Lys Pro Val Thr Phe Asp Cys Ala Val Lys Pro Ser Val
 65 70 75 80
 Thr Cys Val Asp Gln Asp Phe Lys Ser Gln Lys Asn Phe Ile Ile Asn
 85 90 95
 Met Thr Cys Arg Phe Cys Trp Gln Leu Pro Glu Thr Asp Tyr Glu Cys
 100 105 110
 Thr Asn Ser Thr Ser Cys Met Thr Val Ser Cys Pro Arg Gln Arg Tyr
 115 120 125
 Pro Ala Asn Cys Thr Val Arg Asp His Val His Cys Leu Gly Asn Arg
 130 135 140

123

Thr	Phe	Pro	Lys	Met	Leu	Tyr	Cys	Asn	Trp	Thr	Gly	Gly	Tyr	Lys	Trp
145					150					155					160
Ser	Thr	Ala	Leu	Ala	Leu	Ser	Ile	Thr	Leu	Gly	Gly	Phe	Gly	Ala	Asp
			165						170					175	
Arg	Phe	Tyr	Leu	Gly	Gln	Trp	Xaa	Glu	Gly	Leu	Gly	Lys	Leu	Phe	Ser
			180					185					190		
Phe	Gly	Gly	Leu	Gly	Ile	Trp	Thr	Leu	Ile	Asp	Val	Leu	Leu	Ile	Gly
			195				200					205			
Val	Gly	Tyr	Val	Gly	Pro	Ala	Asp	Gly	Ser	Leu	Tyr	Ile	Xaa		
			210				215					220			

<210> 228
 <211> 49
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (49)
 <223> Xaa equals stop translation

<400> 228

Met	Cys	Ile	His	Tyr	Ser	Arg	Val	Ile	Phe	Ser	Phe	Leu	Lys	Leu	Arg
1					5				10					15	
Ile	Lys	Ser	Ile	Ser	Trp	Tyr	Ala	Met	Trp	Leu	Tyr	Phe	Phe	Cys	Tyr
			20					25						30	
Leu	Asn	Cys	Leu	Ala	Lys	Val	Arg	Ser	Ala	Thr	Thr	Tyr	Leu	Tyr	Val
		35					40					45			

Xaa

<210> 229
 <211> 41
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (41)
 <223> Xaa equals stop translation

<400> 229

Met	Leu	Pro	Val	Cys	Val	Phe	Lys	Leu	Leu	Leu	Tyr	Leu	Tyr	Val	Leu
1				5				10						15	
Ile	Arg	Ile	Cys	Thr	Ile	Ile	Trp	Cys	Phe	Lys	Val	Tyr	Ile	Asn	Ala
			20				25						30		
Val	Ile	Leu	Asn	Lys	Ser	Ser	Arg	Xaa							

35

40

124

<210> 230
<211> 53
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (53)
<223> Xaa equals stop translation

<400> 230
Met Asn Cys Gly Gly Ser Thr Leu Cys Val Leu Ser Phe Cys Ser Val
1 5 10 15
Val Cys Ser Val Glu Ala Ser Cys Gln Ser Thr Val Gln Trp Gly Gly
20 25 30
Ala Ala Ala Arg Val Gly Val Pro Phe Asp Trp Ser Arg Asn Glu Gln
35 40 45
Gly Lys Gly His Xaa
50

<210> 231
<211> 50
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (45)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (50)
<223> Xaa equals stop translation

<400> 231
Met Leu Gly Ser Ile Pro Lys Leu Trp Ser Val Leu Ser Phe Ser Ile
1 5 10 15
Asn Phe Cys Phe Cys Cys Phe Ile Leu Ser Leu Leu Cys Leu Ser Val
20 25 30
Leu Ser Asn Tyr Leu Phe Lys Thr Pro Arg Thr Trp Xaa Thr Leu His
35 40 45
Arg Xaa
50

<210> 232
<211> 45

<212> PRT
<213> Homo sapiens

125

<220>
<221> SITE
<222> (16)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (45)
<223> Xaa equals stop translation

<400> 232
Met Cys Leu Pro Leu Leu His Cys Thr Gly Ala Leu Trp Gly Lys Xaa
1 5 10 15
Val Leu Leu Phe Leu Tyr Cys Leu Ala Gln Ser Phe Ala Tyr Ser Arg
20 25 30
His Gln Thr Val Gly Leu Val Val His Asp Tyr Trp Xaa
35 40 45

<210> 233
<211> 55
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (55)
<223> Xaa equals stop translation

<400> 233
Met Cys Trp Ile Cys Val Trp Leu Phe Phe Ser Pro Thr Lys Thr Ser
1 5 10 15
Cys Phe Pro Trp Leu Ile Arg Pro Gly Pro Arg Ser Phe Thr Asp Ser
20 25 30
His Gly Thr Pro Pro Trp Gln Cys Leu Glu Pro Ser Ser Phe Thr Tyr
35 40 45
Pro Gly Lys Gln Val Trp Xaa
50 55

<210> 234
<211> 69
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (69)
<223> Xaa equals stop translation

<400> 236
Met Thr Lys Leu Leu Ser Leu Ser His Leu Leu Val Thr Phe Phe Asn
1 5 10 15
Ile Ile Ala Ile Lys Cys Lys Lys Gln His Leu Arg His Ser Lys Cys

20

25

177

30

Asn Xaa Asp Thr Thr Phe Lys Asn Lys Met Leu Asn Xaa
 35 40 45

<210> 237

<211> 78

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (78)

<223> Xaa equals stop translation

<400> 237

Met Gln Leu Cys Val Ile Trp Phe Thr Val Ile Phe Leu Ser Gln Ser
 1 5 10 15

Ser Arg Leu Val Lys Glu Lys Ile Ser Asn Thr Ser Gly Glu Lys Gly
 20 25 30

Arg Trp Pro Ala Ile Asp Val Val Ala Leu Cys Pro Ser Arg Thr Ala
 35 40 45

Gly Ile Ser Phe Pro Arg His Phe Leu Tyr Val Ser Cys Ile Val Gly
 50 55 60

Cys Thr Asn Ile Ile Cys Ser Phe Gly Phe Pro Gly Gln Xaa
 65 70 75

<210> 238

<211> 53

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (53)

<223> Xaa equals stop translation

<400> 238

Met Glu Val Val Leu Pro Lys His Ile Leu Asp Ile Trp Val Ile Val
 1 5 10 15

Leu Ile Ile Leu Ala Thr Ile Val Ile Met Thr Ser Leu Leu Leu Cys
 20 25 30

Pro Ala Thr Ala Val Ile Ile Tyr Arg Met Arg Thr His Pro Ile Leu
 35 40 45

Ser Gly Ala Val Xaa
 50

<210> 239

128

<400> 241
Met Ser Trp Pro Leu Cys Thr Leu Leu Phe Ser Trp Asp Cys Ile Leu
1 5 10 15

129

Ala Val Lys Thr Ser Arg Leu Lys Phe Asp Ser Gln Gly Tyr Ile Leu
 20 25 30

Gly Thr Phe Lys Val Ser Phe Gln Arg Asp Phe Ile Asn Arg Leu Asp
 35 40 45

Xaa

<210> 242

<211> 75

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (75)

<223> Xaa equals stop translation

<400> 242

Met Ser Ile Ile Ile Tyr Trp Leu Leu Phe Phe Lys His Leu Leu Trp
 1 5 10 15

Val Leu Ile Ile Gly Met Val Lys Ala Leu His Pro His Tyr Leu Asn
 20 25 30

Leu Arg Ile Tyr Glu Phe Gly Glu Ile Thr Ala Val Leu Gln Arg Lys
 35 40 45

Lys Gln Gly Arg Glu Asn Gly Asn Phe Leu Lys Phe Ser Leu Leu Ser
 50 55 60

Leu Asn Arg Ser Arg Ile Pro Thr Gln Ile Xaa
 65 70 75

<210> 243

<211> 44

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (44)

<223> Xaa equals stop translation

<400> 243

Met Ala Ile His Phe His Ile Ile Gln Trp Leu Leu Leu Cys Tyr Asn
 1 5 10 15

Cys His His Ala Gln Trp Gly Leu Trp His Thr Thr Ala Glu Val Ser
 20 25 30

Gly Cys Gly Arg Asn His Leu Ala Phe Lys Ala Xaa
 35 40

<210> 244
 <211> 65
 <212> PRT
 <213> Homo sapiens

130

<220>
 <221> SITE
 <222> (65)
 <223> Xaa equals stop translation

<400> 244
 Met Tyr Leu Ser Leu Phe Phe Phe Cys Phe Ser Leu Gln Ala Ser Ala
 1 5 10 15
 Val Glu Glu Arg Ser Ala Glu Ser Ser Arg Glu Gly Pro Val Arg Thr
 20 25 30
 Asp Asn Trp Gln Arg Cys Phe Gly Asp Ile Pro Gly Thr Pro Thr His
 35 40 45
 Leu Val Gln Arg Ser Leu Val Leu Thr Cys Phe Gly Arg Val Leu Ser
 50 55 60
 Xaa
 65

<210> 245
 <211> 84
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (84)
 <223> Xaa equals stop translation

<400> 245
 Met Lys Lys Val Cys Trp Val Trp Ala Leu Ala His Leu Val Leu Cys
 1 5 10 15
 Glu Arg Trp Leu Thr Ala Gly Cys Leu Leu Tyr Val Gly Val Ile Gln
 20 25 30
 Pro Cys Lys Gly Ser Pro Ser Ser Val Cys Lys Ala Arg Arg Cys Leu
 35 40 45
 His Pro Lys Tyr Arg Ile Lys Arg Tyr Gly Tyr Tyr Lys Tyr Ser Val
 50 55 60
 Arg Leu Ile Ile Cys His His His Pro His Ala Leu Lys Ala Glu Leu
 65 70 75 80
 Thr Asp Asp Xaa

<210> 246

<211> 72
<212> PRT
<213> Homo sapiens

131

<220>
<221> SITE
<222> (72)
<223> Xaa equals stop translation

<400> 246
Met Val Gln Gly Pro Leu Thr His Leu Met Leu Val Leu Leu Ile Ser
1 5 10 15
Leu Ile Phe Leu Ser Arg Gly Ser Gly Arg Ala Trp Ala Phe Ser His
20 25 30
Ser Cys Phe Lys Thr Ser Asp Leu Leu Pro Cys Arg Asn Arg Trp Glu
35 40 45
Val Ile Glu Phe Leu His Tyr Ser Asn Leu His Ser His Ile Ser Leu
50 55 60
Ser Val Thr Lys Thr Phe Leu Xaa
65 70

<210> 247
<211> 57
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (57)
<223> Xaa equals stop translation

<400> 247
Met Arg Ser Tyr Phe Pro Phe Ser Val Cys Pro Phe Pro Phe Cys Ser
1 5 10 15
Pro Val Phe Phe Phe Val Phe Thr Asp Val Tyr Leu Cys Phe Phe Phe
20 25 30
Val Phe Ala Val Gly Arg His Leu Ser Asp Pro Phe Pro Ile Leu Phe
35 40 45
Phe Thr His Lys Cys Pro Asp Val Xaa
50 55

<210> 248
<211> 67
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (67)

<223> Xaa equals stop translation 132

<400> 248

Met Arg Ala Cys Gly Trp Asp Leu Ser Ile Leu Leu Val Gly Leu Val
1 5 10 15

Met Gly Arg Glu Gly Cys Tyr Ser Arg Leu Pro Pro Thr Glu Tyr Gln
20 25 30

Lys Gln Ala Gly Ser Ser Gly Val Cys Lys Asp Val Arg Pro Arg Asn
35 40 45

Gln Pro Ser Pro Ser Tyr Pro Cys Lys Ser Leu Ser Pro His Ala Pro
50 55 60

Leu Leu Xaa
65

<210> 249

<211> 46

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (46)

<223> Xaa equals stop translation

<400> 249

Met Tyr Leu Ile Leu Ser Trp Leu Phe Leu Cys Lys Leu Val Lys Cys
1 5 10 15

Tyr Phe Glu Ile Leu Leu Phe Ser Thr Ser Pro Gln Leu Leu Gln Trp
20 25 30

Thr Val Ile Val Thr Tyr Cys Gly Pro Leu Leu Arg Phe Xaa
35 40 45

<210> 250

<211> 54

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (54)

<223> Xaa equals stop translation

<400> 250

Met Leu Val Phe Leu Leu Leu Phe Ser Thr Val Thr Val Leu Cys Leu
1 5 10 15

Lys Val Val Phe Ser Leu Lys Ala Val Ala Tyr Ile Val Lys Asn Glu
20 25 30

Gly Leu Cys Leu Lys Phe Ile Ala Leu Gln Arg Val Val Ser Leu Lys

35

40

133

45

Ser Cys Thr Ile Lys Xaa
50

<210> 251

<211> 57

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (57)

<223> Xaa equals stop translation

<400> 251

Met Thr Phe Leu Leu Gln Trp Phe Pro Leu Gly Arg Ala Arg Val Val
1 5 10 15

Gly Asp Leu Cys Gly Phe Ser Thr Gln Ile His Pro Gly Val Ser Arg
20 25 30

Ala Gly Met Ala Asp Leu Glu Ser Pro Pro Phe Pro Arg Thr Cys Ser
35 40 45

Val Pro Arg Ala Ala Asn Lys Gly Xaa
50 55

<210> 252

<211> 42

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (42)

<223> Xaa equals stop translation

<400> 252

Met Phe Val Lys Tyr His Val Ile Met Val Ile Ile Phe Ile Phe Ile
1 5 10 15

Leu Ile Thr Ser Asp Lys His Gly Glu Ile Ile Tyr Ile Lys Tyr Ile
20 25 30

Asp Arg Val Ile Ile Thr Glu Arg Ile Xaa
35 40

<210> 253

<211> 161

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (161)

134

<223> Xaa equals stop translation

<400> 253

Met Gln Arg Val Ser Gly Leu Leu Ser Trp Thr Leu Ser Arg Val Leu
 1 5 10 15

Trp Leu Ser Gly Leu Ser Glu Pro Gly Ala Ala Arg Gln Pro Arg Ile
 20 25 30

Met Glu Glu Lys Ala Leu Glu Val Tyr Asp Leu Ile Arg Thr Ile Arg
 35 40 45

Asp Pro Glu Lys Pro Asn Thr Leu Glu Glu Leu Glu Val Val Ser Glu
 50 55 60

Ser Cys Val Glu Val Gln Glu Ile Asn Glu Glu Glu Tyr Leu Val Ile
 65 70 75 80

Ile Arg Phe Thr Pro Thr Val Pro His Cys Ser Leu Ala Thr Leu Ile
 85 90 95

Gly Leu Cys Leu Arg Val Lys Leu Gln Arg Cys Leu Pro Phe Lys His
 100 105 110

Lys Leu Glu Ile Tyr Ile Ser Glu Gly Thr His Ser Thr Glu Glu Asp
 115 120 125

Ile Asn Lys Gln Ile Asn Asp Lys Glu Arg Val Ala Ala Ala Met Glu
 130 135 140

Asn Pro Asn Leu Arg Glu Ile Val Glu Gln Cys Val Leu Glu Pro Asp
 145 150 155 160

Xaa

<210> 254

<211> 51

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (51)

<223> Xaa equals stop translation

<400> 254

Met Leu Phe Phe Ser Leu Lys Glu Ser Leu Tyr Ile Phe His Thr Ala
 1 5 10 15

Ile Leu Leu Val Val Cys Phe Ala Cys Ala Val Val Cys Gln Tyr Val
 20 25 30

Ile Val Arg Val Cys Ala Val Val Phe Cys Phe Ser Lys Ser Gln Ser
 35 40 45

Leu Ile Xaa
50

135

<210> 255
<211> 279
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (279)
<223> Xaa equals stop translation

<400> 255
Met Leu Ile Phe Gly Ala Ile Phe Gly Cys Leu Asp Pro Val Ala Thr
1 5 10 15
Leu Ala Ala Val Met Thr Glu Lys Ser Pro Phe Thr Thr Pro Ile Gly
20 25 30
Arg Lys Asp Glu Ala Asp Leu Ala Lys Ser Ala Leu Ala Met Ala Asp
35 40 45
Ser Asp His Leu Thr Ile Tyr Asn Ala Tyr Leu Gly Trp Lys Lys Ala
50 55 60
Arg Gln Glu Gly Gly Tyr Arg Ser Glu Ile Thr Tyr Cys Arg Arg Asn
65 70 75 80
Phe Leu Asn Arg Thr Ser Leu Leu Thr Leu Glu Asp Val Lys Gln Glu
85 90 95
Leu Ile Lys Leu Val Lys Ala Ala Gly Phe Ser Ser Ser Thr Thr Ser
100 105 110
Thr Ser Trp Glu Gly Asn Arg Ala Ser Gln Thr Leu Ser Phe Gln Glu
115 120 125
Ile Ala Leu Leu Lys Ala Val Leu Val Ala Gly Leu Tyr Asp Asn Val
130 135 140
Gly Lys Ile Ile Tyr Thr Lys Ser Val Asp Val Thr Glu Lys Leu Ala
145 150 155 160
Cys Ile Val Glu Thr Ala Gln Gly Lys Ala Gln Val His Pro Ser Ser
165 170 175
Val Asn Arg Asp Leu Gln Thr His Gly Trp Leu Leu Tyr Gln Glu Lys
180 185 190
Ile Arg Tyr Ala Arg Val Tyr Leu Arg Glu Thr Thr Leu Ile Thr Pro
195 200 205
Phe Pro Val Leu Leu Phe Gly Gly Asp Ile Glu Val Gln His Arg Glu
210 215 220
Arg Leu Leu Ser Ile Asp Gly Trp Ile Tyr Phe Gln Ala Pro Val Lys

<400> 257

Met Ile Met Ala Gln Lys Ile Gly Gly Leu Thr Trp Trp Ala Ile Met
1 5 10 15

Phe Ile Ile Leu Phe Glu Ile Thr Gly Thr Ser Ser Ser Phe Leu Arg
20 25 30

Ile Asn Ala Leu Pro His Phe Ser Met Asn Arg Cys Gly Glu Ala Tyr
35 40 45

137

Phe Pro Phe Ser Tyr Leu Tyr Thr Ser Leu Gln Lys Gln Phe Leu Met
 50 55 60

Lys Val Ser Gly Ile Val Lys Asn Leu Arg Gly Asn Asp Asp Trp Arg
 65 70 75 80

Cys Phe Gly Val Phe Phe Cys Ile His Phe Leu Met Arg Lys Val Leu
 85 90 95

Asn Val Val Gln Val Arg Pro Asn Tyr Tyr Leu Thr Ile Ile Gly Arg
 100 105 110

Phe Tyr Val Ser Val Lys Val Phe Lys Xaa
 115 120

<210> 258

<211> 59

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (59)

<223> Xaa equals stop translation

<400> 258

Met Gly Lys Ile Cys Lys Asn Trp Val Ser Phe Leu Asp Asn Val Leu
 1 5 10 15

Leu Leu Ile Leu Phe Leu Tyr Gly Leu Cys Leu Gly Trp Leu Cys Ile
 20 25 30

Tyr His Gln Ser Tyr Ser Thr Ala Cys Ile Cys Val Val Thr Asp Ala
 35 40 45

Glu Ile Gln Gln Lys Ser Leu His Ser Ile Xaa
 50 55

<210> 259

<211> 68

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (68)

<223> Xaa equals stop translation

<400> 259

Met Leu Val Leu Leu Trp Leu Gly Trp Ile Ser Ser Lys Ser Met Leu
 1 5 10 15

Ala Ala Tyr Phe Val Ala Pro Lys Tyr Pro Leu Lys Leu Ala Leu Val
 20 25 30

138

Ser Glu Pro Glu Ser Ser Ser Leu Ile Leu Lys Phe Leu Ser Leu Lys
 35 40 45

Asp Phe Leu Cys Cys Tyr Thr Thr Lys Leu Ser Val Asn Pro Pro Leu
 50 55 60

Leu Asn Asp Xaa
 65

<210> 260
 <211> 46
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (46)
 <223> Xaa equals stop translation

<400> 260
 Met Val Ser Phe His Phe Gln Cys Thr Ser Tyr Phe Val Arg Leu Phe
 1 5 10 15

Phe Gln Leu Gln Leu Phe Val Gly Leu Val Ile Val Leu Ala Leu Leu
 20 25 30

Ile Ser His Ser Leu Thr Tyr Ser Phe His Lys His Leu Xaa
 35 40 45

<210> 261
 <211> 110
 <212> PRT
 <213> Homo sapiens

<400> 261
 Phe Tyr Ile Ala Asp His Ser Phe Thr Ala Arg Pro Thr Leu Arg Met
 1 5 10 15

Phe Arg Ile Ser Ala Val Val Ala Thr Asp Lys Met Thr Phe Thr Ser
 20 25 30

Gly Gly Thr Leu Phe Gly Asp Gly Cys Ala Ser Ser Val Ala Gly Glu
 35 40 45

Val Met Asn Cys Gln Thr Val Leu Cys Ile Leu Trp Thr Pro Phe Val
 50 55 60

Phe Cys Pro Ser Ile Ala Val Ile Ile Ile Pro Cys Val Phe Thr Ser
 65 70 75 80

Lys Ala Leu Glu Ala Ile Trp Lys Trp Cys Arg Val Glu Arg Arg Pro
 85 90 95

His Ile Ile Glu Val Asp Val Leu Gly Lys Cys Pro Ala Phe
 100 105 110

<210> 262
 <211> 25
 <212> PRT
 <213> Homo sapiens

139

<400> 262
 Arg Pro Thr Leu Arg Met Phe Arg Ile Ser Ala Val Val Ala Thr Asp
 1 5 10 15

Lys Met Thr Phe Thr Ser Gly Gly Thr
 20 25

<210> 263
 <211> 28
 <212> PRT
 <213> Homo sapiens

<400> 263
 Pro Ser Ile Ala Val Ile Ile Ile Pro Cys Val Phe Thr Ser Lys Ala
 1 5 10 15

Leu Glu Ala Ile Trp Lys Trp Cys Arg Val Glu Arg
 20 25

<210> 264
 <211> 20
 <212> PRT
 <213> Homo sapiens

<400> 264
 Thr Ser Val Ser Phe His His Arg Tyr Lys Ser Ser Asp Arg Pro Ala
 1 5 10 15

His Lys Val Ser
 20

<210> 265
 <211> 115
 <212> PRT
 <213> Homo sapiens

<400> 265
 Arg His Asn Asp Phe Asn Lys Leu Ser Tyr Thr Glu Cys Asn Asn Met
 1 5 10 15

Asn Lys Arg Met Ala Lys Pro Glu Lys Lys Lys Gly Ser Val Lys Ser
 20 25 30

Ser Leu Gly Ile Phe Leu Gly Pro Asn Cys His Leu Ile Ser Ser Leu
 35 40 45

Phe Leu Phe Ser Val Ser Leu Tyr Pro Phe Ala Thr Gln Phe Pro Phe
 50 55 60

140

His Tyr Val Leu Ile Phe Ile Ile Gln Ala Phe Gly Leu Cys Leu Pro
 65 70 75 80

Leu Thr Glu Arg Gln Glu Ala Lys Ser Gly Leu Gly Gly Leu Cys Pro
 85 90 95

Asp Tyr Thr Trp Pro Cys Pro Cys Leu Leu Val Ser Cys Leu Ser Leu
 100 105 110

Leu Arg Leu
 115

<210> 266
 <211> 114
 <212> PRT
 <213> Homo sapiens

<400> 266
 Cys Glu Val Phe Ser Trp His Phe Pro Trp Ser Lys Leu Ser Pro His
 1 5 10 15

Leu Phe Leu Val Ser Phe Leu Cys Ile Pro Leu Ser Leu Cys His Thr
 20 25 30

Val Ser Phe Ser Leu Cys Ser Asn Ile Tyr Asn Pro Gly Leu Arg Thr
 35 40 45

Met Leu Ala Pro His Arg Glu Thr Gly Gly Gln Val Trp Ala Gly Trp
 50 55 60

Ala Leu Ser Arg Leu His Val Ala Leu Pro Met Ser Leu Gly Val Leu
 65 70 75 80

Ser Leu Pro Ala Pro Thr Val Thr Val Val Arg Met Glu Gly Gly Asp
 85 90 95

Trp Lys Val Cys Glu Gln Leu Gly Gln Cys Thr Tyr Ser His Arg Met
 100 105 110

Thr Lys

<210> 267
 <211> 23
 <212> PRT
 <213> Homo sapiens

<400> 267
 Lys Arg Met Ala Lys Pro Glu Lys Lys Lys Gly Ser Val Lys Ser Ser
 1 5 10 15

Leu Gly Ile Phe Leu Gly Pro
 20

<210> 268

<211> 31
 <212> PRT
 <213> Homo sapiens

141

<400> 268
 Tyr Asn Pro Gly Leu Arg Thr Met Leu Ala Pro His Arg Glu Thr Gly
 1 5 10 15

Gly Gln Val Trp Ala Gly Trp Ala Leu Ser Arg Leu His Val Ala
 20 25 30

<210> 269
 <211> 251
 <212> PRT
 <213> Homo sapiens

<400> 269
 Met Ser Pro Tyr Ala Ser Gln Gly Phe Pro Phe Leu Pro Pro Tyr Pro
 1 5 10 15

Pro Gln Glu Ala Asn Arg Ser Ile Thr Ser Leu Ser Val Ala Asp Thr
 20 25 30

Val Ser Ser Ser Thr Thr Ser His Thr Thr Ala Lys Pro Ala Ala Pro
 35 40 45

Ser Phe Gly Val Leu Ser Asn Leu Pro Leu Pro Ile Pro Thr Val Asp
 50 55 60

Ala Ser Ile Pro Thr Ser Gln Asn Gly Phe Gly Tyr Lys Met Pro Asp
 65 70 75 80

Val Pro Asp Ala Phe Pro Glu Leu Ser Glu Leu Ser Val Ser Gln Leu
 85 90 95

Thr Asp Met Asn Glu Gln Glu Glu Val Leu Leu Glu Gln Phe Leu Thr
 100 105 110

Leu Pro Gln Leu Lys Gln Ile Ile Thr Asp Lys Asp Asp Leu Val Lys
 115 120 125

Ser Ile Glu Glu Leu Ala Arg Lys Asn Leu Leu Leu Glu Pro Ser Leu
 130 135 140

Glu Ala Lys Arg Gln Thr Val Leu Asp Lys Tyr Glu Leu Leu Thr Gln
 145 150 155 160

Met Lys Ser Thr Phe Glu Lys Lys Met Gln Arg Gln His Glu Leu Ser
 165 170 175

Glu Ser Cys Ser Ala Ser Ala Leu Gln Ala Arg Leu Lys Val Ala Ala
 180 185 190

His Glu Ala Glu Glu Glu Ser Asp Asn Ile Ala Glu Asp Phe Leu Glu
 195 200 205

Gly Lys Met Glu Ile Asp Asp Phe Leu Ser Ser Phe Met Glu Lys Arg

210 215 192 220
 Thr Ile Cys His Cys Arg Arg Ala Lys Glu Glu Lys Leu Gln Gln Ala
 225 230 235 240

Ile Ala Met His Ser Gln Phe His Ala Pro Leu
 245 250

<210> 270
 <211> 23
 <212> PRT
 <213> Homo sapiens

<400> 270
 Leu Pro Pro Tyr Pro Pro Gln Glu Ala Asn Arg Ser Ile Thr Ser Leu
 1 5 10 15

Ser Val Ala Asp Thr Val Ser
 20

<210> 271
 <211> 27
 <212> PRT
 <213> Homo sapiens

<400> 271
 Thr Ala Lys Pro Ala Ala Pro Ser Phe Gly Val Leu Ser Asn Leu Pro
 1 5 10 15

Leu Pro Ile Pro Thr Val Asp Ala Ser Ile Pro
 20 25

<210> 272
 <211> 25
 <212> PRT
 <213> Homo sapiens

<400> 272
 Pro Asp Val Pro Asp Ala Phe Pro Glu Leu Ser Glu Leu Ser Val Ser
 1 5 10 15

Gln Leu Thr Asp Met Asn Glu Gln Glu
 20 25

<210> 273
 <211> 29
 <212> PRT
 <213> Homo sapiens

<400> 273
 Gln Phe Leu Thr Leu Pro Gln Leu Lys Gln Ile Ile Thr Asp Lys Asp
 1 5 10 15

Asp Leu Val Lys Ser Ile Glu Glu Leu Ala Arg Lys Asn

20

25

143

<210> 274

<211> 25

<212> PRT

<213> Homo sapiens

<400> 274

Arg Gln Thr Val Leu Asp Lys Tyr Glu Leu Leu Thr Gln Met Lys Ser
1 5 10 15

Thr Phe Glu Lys Lys Met Gln Arg Gln
20 25

<210> 275

<211> 28

<212> PRT

<213> Homo sapiens

<400> 275

Ala Ser Ala Leu Gln Ala Arg Leu Lys Val Ala Ala His Glu Ala Glu
1 5 10 15

Glu Glu Ser Asp Asn Ile Ala Glu Asp Phe Leu Glu
20 25

<210> 276

<211> 27

<212> PRT

<213> Homo sapiens

<400> 276

Met Glu Lys Arg Thr Ile Cys His Cys Arg Arg Ala Lys Glu Glu Lys
1 5 10 15

Leu Gln Gln Ala Ile Ala Met His Ser Gln Phe
20 25

<210> 277

<211> 69

<212> PRT

<213> Homo sapiens

<400> 277

Thr Arg Pro Val Phe Leu Ser Met Thr Pro Leu Lys Gly Ile Lys Ser
1 5 10 15

Val Ile Leu Pro Gln Val Phe Leu Cys Ala Tyr Met Ala Ala Phe Asn
20 25 30

Ser Ile Asn Gly Asn Arg Ser Tyr Thr Cys Lys Pro Leu Glu Arg Ser
35 40 45

Leu Leu Met Ala Gly Ala Val Ala Ser Ser Thr Phe Leu Gly Val Ile


```

50          55          144          60
Pro Gln Phe Val Gln
65

<210> 278
<211> 21
<212> PRT
<213> Homo sapiens

<400> 278
Pro Leu Lys Gly Ile Lys Ser Val Ile Leu Pro Gln Val Phe Leu Cys
1          5          10          15

Ala Tyr Met Ala Ala
20

<210> 279
<211> 21
<212> PRT
<213> Homo sapiens

<400> 279
Ala Phe Asn Ser Ile Asn Gly Asn Arg Ser Tyr Thr Cys Lys Pro Leu
1          5          10          15

Glu Arg Ser Leu Leu
20

<210> 280
<211> 129
<212> PRT
<213> Homo sapiens

<400> 280
Met Ser Asp Phe Glu Lys Val Asp Ile Ser Val His Gln His Ile His
1          5          10          15

Val Gly Pro Leu Leu Leu Met Thr Thr Glu Ser Trp Gly Pro Ser Cys
20          25          30

Ala Pro Ser Pro Ala Leu Leu Ser Gly His Thr Ala Ala Ser Phe Thr
35          40          45

His Thr Leu Gly Gly Val Leu Gly Cys Pro Pro Tyr His Lys Phe Tyr
50          55          60

Ser Ser Ala His Thr Ser Asp His Arg Lys Glu Thr Asn Lys Val Glu
65          70          75          80

Glu Gly Arg Trp Val Asp Val Thr Arg Ser Leu Gly Asn Phe Asn Phe
85          90          95

Arg Arg Lys Phe Phe Cys Val Ser Glu Leu Leu Ile Cys Gly Ile Phe
100         105         110

```

145

Leu	Asp	Ser	Ser	Trp	Lys	Leu	Gln	Ile	Asn	Ser	Asn	Asp	Cys	Lys	Val
		115					120					125			

Leu

<210> 281
 <211> 30
 <212> PRT
 <213> Homo sapiens

<400> 281

Val	Gly	Pro	Leu	Leu	Leu	Met	Thr	Thr	Glu	Ser	Trp	Gly	Pro	Ser	Cys
1				5					10					15	

Ala	Pro	Ser	Pro	Ala	Leu	Leu	Ser	Gly	His	Thr	Ala	Ala	Ser
			20					25					30

<210> 282
 <211> 27
 <212> PRT
 <213> Homo sapiens

<400> 282

Glu	Thr	Asn	Lys	Val	Glu	Glu	Gly	Arg	Trp	Val	Asp	Val	Thr	Arg	Ser
1				5					10					15	

Leu	Gly	Asn	Phe	Asn	Phe	Arg	Arg	Lys	Phe	Phe
			20					25		

<210> 283
 <211> 140
 <212> PRT
 <213> Homo sapiens

<400> 283

Gly	Arg	Gly	Asp	Lys	Pro	Arg	Gln	Asp	Arg	Pro	Ala	Ser	Leu	Arg	Leu
1				5					10					15	

Lys	Gly	Pro	Pro	Ser	Cys	Gln	Ala	Pro	Ala	Ser	His	Ser	Ser	Thr	Leu
			20					25					30		

Ser	Ser	His	Cys	Pro	Cys	Ser	Leu	Phe	Ala	Cys	Gly	Ser	Val	Trp	Pro
		35					40					45			

Gly	Ser	Leu	Gly	Ser	Gly	Ile	Phe	Ala	Arg	Leu	Ser	Gln	Leu	Leu	Pro
	50					55					60				

Ser	Pro	Ala	Ser	Trp	Gly	Trp	Asp	Phe	Leu	Thr	Leu	Arg	Gln	Ala	Gln
	65				70					75					80

Gln	Met	Leu	Gly	Pro	Ser	Leu	Cys	Pro	Gly	His	Ser	Thr	Ser	Ala	His
				85					90					95	

<400> 285
Ser Leu Arg Leu Lys Gly Pro Pro Ser Cys Gln Ala Pro Ala Ser His
1 5 10 15
Ser Ser Thr Leu Ser Ser His Cys Pro Cys Ser Leu Phe Ala
20 25 30

(47)

<210> 286
 <211> 30
 <212> PRT
 <213> Homo sapiens

<400> 286
 Gln Gln Met Leu Gly Pro Ser Leu Cys Pro Gly His Ser Thr Ser Ala
 1 5 10 15

His Gln His Tyr Gly Ala Tyr Val Leu Pro Arg Asp Leu Cys
 20 25 30

<210> 287
 <211> 31
 <212> PRT
 <213> Homo sapiens

<400> 287
 Asp Leu Gln Val Phe Val Ser Arg Asp Leu Pro His Ala Arg Pro Leu
 1 5 10 15

Pro Leu Thr Ala Ala Pro Phe Pro Leu Ile Val Pro Val Pro Phe
 20 25 30

<210> 288
 <211> 39
 <212> PRT
 <213> Homo sapiens

<400> 288
 Ala Gln Val His Ile Ser Thr Met Gly Pro Met Ser Cys Pro Glu Thr
 1 5 10 15

Ser Ala Pro Ser Cys Ser His Pro Gln Phe Arg Ala Arg Arg Pro Ser
 20 25 30

Arg Thr Pro Glu Ser Pro Val
 35

<210> 289
 <211> 17
 <212> PRT
 <213> Homo sapiens

<400> 289
 Gln Ala Pro Pro Arg Gln Thr Cys Lys Ser Ser Ser Gln Gly Thr Ser
 1 5 10 15

Leu

<210> 290

<211> 314
 <212> PRT
 <213> Homo sapiens

148

<220>
 <221> SITE
 <222> (27)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (111)
 <223> Xaa equals any of the naturally occurring L-amino acids

<400> 290
 Ala Ala Leu Arg Pro Ser Gly Ser Leu Ala Gly Pro Glu Trp Pro Trp
 1 5 10 15
 Gln His Trp Cys Gly Cys Trp Arg Glu His Xaa Val Lys Pro Gln Gln
 20 25 30
 Val Asp Leu His Ser Ala Arg Leu Trp Ala Ala Pro Ala Ala Val Gly
 35 40 45
 Pro Ala His Ala Gly Gly Ser Pro Gly Met Pro Pro Gly Gly Thr Ala
 50 55 60
 Pro His Ala Arg Arg His Ser Leu Pro Ser Pro Thr Ala Gln Ser His
 65 70 75 80
 Leu Trp His Val His Gly Leu Arg Gln Arg Gly Pro Lys Ala Val Pro
 85 90 95
 Leu Asp Leu Ala Gln Leu Val Thr Thr Thr Thr Pro Leu Phe Xaa Leu
 100 105 110
 Ala Leu Ser Ala Leu Leu Leu Gly Arg Arg His His Pro Leu Gln Leu
 115 120 125
 Ala Ala Met Gly Pro Leu Cys Leu Gly Ala Ala Cys Ser Leu Ala Gly
 130 135 140
 Glu Phe Arg Thr Pro Pro Thr Gly Cys Gly Phe Leu Leu Ala Ala Thr
 145 150 155 160
 Cys Leu Arg Gly Leu Lys Ser Val Gln Gln Ser Ala Leu Leu Gln Glu
 165 170 175
 Glu Arg Leu Asp Ala Val Thr Leu Leu Tyr Ala Thr Ser Leu Pro Ser
 180 185 190
 Phe Cys Leu Leu Ala Gly Ala Ala Leu Val Leu Glu Ala Gly Val Ala
 195 200 205
 Pro Pro Pro Thr Ala Gly Asp Ser Arg Leu Trp Ala Cys Ile Leu Leu
 210 215 220
 Ser Cys Leu Leu Ser Val Leu Tyr Asn Leu Ala Ser Phe Ser Leu Leu

```

<400> 291
Gly Gln Pro Ser Gly Pro Pro Ala Ala Trp Pro Gly Pro Ser Gly His
  1          5          10          15
Gly Ser Thr Gly Val Ala Ala Gly Gly Ser Thr Xaa Ser Ser Leu Asn
          20          25          30
Lys Trp Ile Phe Thr Val His Gly Phe Gly Arg Pro Leu Leu Ser
          35          40          45
Ala Leu His Met Leu Val Ala Ala Leu Ala Cys His Arg Gly Ala Arg
  50          55          60
Arg Pro
  65

```

<400> 292
Trp Pro Gly Pro Ser Gly His Gly Ser Thr Gly Val Ala Ala Gly Gly
1 5 10 15

150

Ser Thr Xaa Ser Ser
20

<210> 293
<211> 26
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (15)
<223> Xaa equals any of the naturally occurring L-amino acids

<400> 293
Glu Trp Pro Trp Gln His Trp Cys Gly Cys Trp Arg Glu His Xaa Val
1 5 10 15

Lys Pro Gln Gln Val Asp Leu His Ser Ala
20 25

<210> 294
<211> 28
<212> PRT
<213> Homo sapiens

<400> 294
Gln Gln Ser Ala Leu Leu Gln Glu Glu Arg Leu Asp Ala Val Thr Leu
1 5 10 15

Leu Tyr Ala Thr Ser Leu Pro Ser Phe Cys Leu Leu
20 25

<210> 295
<211> 27
<212> PRT
<213> Homo sapiens

<400> 295
Ala Cys Ile Leu Leu Ser Cys Leu Leu Ser Val Leu Tyr Asn Leu Ala
1 5 10 15

Ser Phe Ser Leu Leu Ala Leu Thr Ser Ala Leu
20 25

<210> 296
<211> 21
<212> PRT
<213> Homo sapiens

<400> 296
Ser Leu Asn Lys Trp Ile Phe Thr Val His Gly Phe Gly Arg Pro Leu
1 5 10 15

Leu Leu Ser Ala Leu
20

151

<210> 297
<211> 17
<212> PRT
<213> Homo sapiens

<400> 297
Lys Ser Thr Leu Ser Ala Ala Val Val Ala Thr Ile Leu Arg Thr Leu
1 5 10 15

Ala

<210> 298
<211> 100
<212> PRT
<213> Homo sapiens

<400> 298
Gly Asp His Ser Glu Gln Cys Leu Ile Lys Glu Met Gly Ala Arg Glu
1 5 10 15

Arg Arg Phe Cys Lys Ala Arg Gly Tyr Arg Asp Thr Gly Arg Glu Ala
20 25 30

Gln Ala Lys Ala Gly Gly Arg Arg Gly Ser Gln Trp Asn Glu Ser Gln
35 40 45

Cys Ser Ser Gln Arg Pro Arg Pro Ala Lys Glu Val Arg Lys Thr Arg
50 55 60

Pro Arg Ala Gly Val Gly Arg Gly Pro Ala Leu Leu Gln Leu Ser Leu
65 70 75 80

Leu Gln Gln Val Val Leu Tyr Val Arg Pro Ser Leu Arg Leu Val Trp
85 90 95

Leu Lys Ala Ser
100

<210> 299
<211> 84
<212> PRT
<213> Homo sapiens

<400> 299
Met Glu Arg Gly Glu Tyr Gly Gly Trp Gly Thr Tyr Gly Ser Leu Asp
1 5 10 15

Leu Gly Ser Gln Leu Cys Thr Val Arg Ser Ser Gly Pro Cys Gly Ser
20 25 30

Leu His Trp Gly Gln His Arg Ser Pro Ile Ser Gly Pro Asp Pro Asn

35

40

152

45

Pro Ser Ser Ser Arg Gly Gln Gln Ser Ile Gly Ser Lys Val Gly Ser
50 55 60

Pro Ser Arg Ser Gln Trp Arg Ser Trp Lys Glu Val Gly Arg Asp Pro
65 70 75 80

Glu Lys Gly Glu

<210> 300

<211> 23

<212> PRT

<213> Homo sapiens

<400> 300

Gln Ala Lys Ala Gly Gly Arg Arg Gly Ser Gln Trp Asn Glu Ser Gln
1 5 10 15

Cys Ser Ser Gln Arg Pro Arg
20

<210> 301

<211> 26

<212> PRT

<213> Homo sapiens

<400> 301

Val Gly Arg Gly Pro Ala Leu Leu Gln Leu Ser Leu Leu Gln Gln Val
1 5 10 15

Val Leu Tyr Val Arg Pro Ser Leu Arg Leu
20 25

<210> 302

<211> 22

<212> PRT

<213> Homo sapiens

<400> 302

Tyr Gly Ser Leu Asp Leu Gly Ser Gln Leu Cys Thr Val Arg Ser Ser
1 5 10 15

Gly Pro Cys Gly Ser Leu
20

<210> 303

<211> 20

<212> PRT

<213> Homo sapiens

<400> 303

Lys Val Gly Ser Pro Ser Arg Ser Gln Trp Arg Ser Trp Lys Glu Val

1

5

153
10

15

Gly Arg Asp Pro
20

<210> 304

<211> 214

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (18)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 304

Met Pro Gln Ser Leu Ser Ser Leu Ala Ser Ser Ser Ser Ser Phe Gln
1 5 10 15

Arg Xaa Lys Pro Cys Phe Gly Lys Lys Asn Asp Gly Glu Asn Gln Glu
20 25 30

His Ser Leu Gly Thr Glu Pro Ile Ile Thr Trp Lys Asp Phe Gln Lys
35 40 45

Thr Met Pro Trp Glu Ile Val Ile Leu Val Gly Gly Gly Tyr Ala Leu
50 55 60

Ala Ser Gly Ser Lys Ser Ser Gly Leu Ser Thr Trp Ile Gly Asn Gln
65 70 75 80

Met Leu Ser Leu Ser Ser Leu Pro Pro Trp Ala Val Thr Leu Leu Ala
85 90 95

Cys Ile Leu Val Ser Ile Val Thr Glu Phe Val Ser Asn Pro Ala Thr
100 105 110

Ile Thr Ile Phe Leu Pro Ile Leu Cys Ser Leu Ser Glu Thr Leu His
115 120 125

Ile Asn Pro Leu Tyr Thr Leu Ile Pro Val Thr Met Cys Ile Ser Phe
130 135 140

Ala Val Met Leu Pro Val Gly Asn Pro Pro Asn Ala Ile Val Phe Ser
145 150 155 160

Tyr Gly His Cys Gln Ile Lys Asp Met Val Lys Ala Gly Leu Gly Val
165 170 175

Asn Val Ile Gly Leu Val Ile Val Met Val Ala Ile Asn Thr Trp Gly
180 185 190

Val Ser Leu Phe His Leu Asp Thr Tyr Pro Ala Trp Ala Arg Val Ser
195 200 205

Asn Ile Thr Asp Gln Ala
210

154

<210> 305

<211> 23

<212> PRT

<213> Homo sapiens

<400> 305

Asn Asp Gly Glu Asn Gln Glu His Ser Leu Gly Thr Glu Pro Ile Ile
1 5 10 15

Thr Trp Lys Asp Phe Gln Lys
20

<210> 306

<211> 24

<212> PRT

<213> Homo sapiens

<400> 306

Ile Gly Asn Gln Met Leu Ser Leu Ser Ser Leu Pro Pro Trp Ala Val
1 5 10 15

Thr Leu Leu Ala Cys Ile Leu Val
20

<210> 307

<211> 27

<212> PRT

<213> Homo sapiens

<400> 307

Ala Thr Ile Thr Ile Phe Leu Pro Ile Leu Cys Ser Leu Ser Glu Thr
1 5 10 15

Leu His Ile Asn Pro Leu Tyr Thr Leu Ile Pro
20 25

<210> 308

<211> 26

<212> PRT

<213> Homo sapiens

<400> 308

Leu Pro Val Gly Asn Pro Pro Asn Ala Ile Val Phe Ser Tyr Gly His
1 5 10 15

Cys Gln Ile Lys Asp Met Val Lys Ala Gly
20 25

<210> 309

<211> 29

<212> PRT

<213> Homo sapiens

<400> 309

155

Leu Val Ile Val Met Val Ala Ile Asn Thr Trp Gly Val Ser Leu Phe
 1 5 10 15

His Leu Asp Thr Tyr Pro Ala Trp Ala Arg Val Ser Asn
 20 25

<210> 310

<211> 133

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (44)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (46)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 310

Glu Thr Cys Pro Ser Asn Gly Ile Glu Leu Arg Gln Ala Pro Thr Ser
 1 5 10 15

Leu Tyr Ile Leu Leu Leu His Ile Gln Pro Thr Pro Thr His Pro Met
 20 25 30

Leu Gly Arg Ser Tyr Val Leu Pro Ala Phe Ser Xaa Asn Xaa Glu His
 35 40 45

Gly Gly Leu Pro Asn Gln Ile Pro Lys Gly Asp Arg Asn Gly Asn Ile
 50 55 60

Arg His Ser Arg Ile Thr Phe Pro Cys Ser Ser Ser Thr Leu Gln Pro
 65 70 75 80

Glu Ser His Leu Gly Phe Ile Arg Ser Lys Leu His Gly Leu Val Arg
 85 90 95

Pro Gly Lys Asp Leu Arg Gly Arg Arg Ser Leu Gln Leu Ser Lys His
 100 105 110

Ser Leu Ser Thr Cys Tyr Met Leu Arg Trp Glu Thr Tyr Lys Gln Val
 115 120 125

Ser Tyr Thr Ala Val
 130

<210> 311

<211> 106

<212> PRT

<213> Homo sapiens

<400> 311

156

Gln Arg His Gln Glu Asn Asp Lys Arg Asn Val His Arg Phe Leu His
 1 5 10 15

Thr Cys Val His Met Pro Met Cys Thr His Thr His Thr Gln Ala Val
 20 25 30

Leu Ser Thr Trp Glu Gly Gln Phe Ser Asn Val Ala Ser Phe Thr Ser
 35 40 45

Leu Lys Arg Ile Pro Leu Ser Ile Ile Tyr Ile His Ser Ser His Ser
 50 55 60

Pro Arg Arg Phe Val Lys Val Cys Gln Leu Arg Gln Glu Lys Ala Leu
 65 70 75 80

Glu Leu Thr Glu Val Tyr Val Ser Ala Ser Leu Lys Leu Gln Leu Tyr
 85 90 95

His Leu His Cys His Phe His Thr Ala Val
 100 105

<210> 312

<211> 24

<212> PRT

<213> Homo sapiens

<400> 312

Arg Gln Ala Pro Thr Ser Leu Tyr Ile Leu Leu Leu His Ile Gln Pro
 1 5 10 15

Thr Pro Thr His Pro Met Leu Gly
 20

<210> 313

<211> 25

<212> PRT

<213> Homo sapiens

<400> 313

Ser His Leu Gly Phe Ile Arg Ser Lys Leu His Gly Leu Val Arg Pro
 1 5 10 15

Gly Lys Asp Leu Arg Gly Arg Arg Ser
 20 25

<210> 314

<211> 22

<212> PRT

<213> Homo sapiens

<400> 314

Arg Asn Val His Arg Phe Leu His Thr Cys Val His Met Pro Met Cys
 1 5 10 15

Thr His Thr His Thr Gln
20

157

<210> 315
<211> 25
<212> PRT
<213> Homo sapiens

<400> 315
Gln Leu Arg Gln Glu Lys Ala Leu Glu Leu Thr Glu Val Tyr Val Ser
1 5 10 15

Ala Ser Leu Lys Leu Gln Leu Tyr His
20 25

<210> 316
<211> 31
<212> PRT
<213> Homo sapiens

<400> 316
Pro Arg Val Arg Gly Arg Lys Glu Pro Gly Cys Leu Gly Pro Gly Arg
1 5 10 15

Ala Gly Gly Asp Ser Gln Lys Glu Ile Gly Ser Trp Gln Gln Met
20 25 30

<210> 317
<211> 296
<212> PRT
<213> Homo sapiens

<400> 317
Leu Ser Lys Gly Asn Arg Ile Met Ala Ala Asp Asp Asp Asn Gly Asp
1 5 10 15

Gly Thr Ser Leu Phe Asp Val Phe Ser Ala Ser Pro Leu Lys Asn Asn
20 25 30

Asp Glu Gly Ser Leu Asp Ile Tyr Ala Gly Leu Asp Ser Ala Val Ser
35 40 45

Asp Ser Ala Ser Lys Ser Cys Val Pro Ser Arg Asn Cys Leu Asp Leu
50 55 60

Tyr Glu Glu Ile Leu Thr Glu Glu Gly Thr Ala Lys Glu Ala Thr Tyr
65 70 75 80

Asn Asp Leu Gln Val Glu Tyr Gly Lys Cys Gln Leu Gln Met Lys Glu
85 90 95

Leu Met Lys Lys Phe Lys Glu Ile Gln Thr Gln Asn Phe Ser Leu Ile
100 105 110

Asn Glu Asn Gln Ser Leu Lys Lys Asn Ile Ser Ala Leu Ile Lys Thr

```
<210> 318
<211> 27
<212> PRT
<213> Homo sapiens
```

```
<210> 319
<211> 23
<212> PRT
<213> Homo sapiens
```

<400> 319
Cys Leu Asp Leu Tyr Glu Glu Ile Leu Thr Glu Glu Gly Thr Ala Lys
1 5 10 15

Glu Ala Thr Tyr Asn Asp Leu
20

159

<210> 320

<211> 26

<212> PRT

<213> Homo sapiens

<400> 320

Asp Glu Glu Ile Ser Asn Leu His Gln Lys Ile Val Leu Ser Phe His
1 5 10 15

Ile Phe Glu Ile Ile Ile Lys Leu Gln Gly
20 25

<210> 321

<211> 22

<212> PRT

<213> Homo sapiens

<400> 321

Glu Lys Glu Gly Lys Pro His Ser Asp Lys Arg Ser Thr Ser His Leu
1 5 10 15

Pro Thr Ser Val Glu Lys
20

<210> 322

<211> 26

<212> PRT

<213> Homo sapiens

<400> 322

Thr Glu Arg Val Arg Lys Asp Leu Ser Thr Gly Cys Gly Asp Gly Glu
1 5 10 15

Pro Arg Ile Leu Glu Ala Ser Gln Arg Leu
20 25

<210> 323

<211> 115

<212> PRT

<213> Homo sapiens

<400> 323

Lys Ser Tyr Phe Arg Thr Met Gly Gly Thr Lys Arg Gly Ile Lys Lys
1 5 10 15

Leu Val Asn Val Cys Leu Lys His Pro Lys Asn Thr Ser Leu Ser Gln
20 25 30

Gln Leu Val Phe Ala Lys Ile Asn Lys Ile Leu Ile Ser Lys Thr Thr
35 40 45

160

Lys	Ser	Thr	Asn	Leu	Lys	Gly	Leu	Lys	Cys	Leu	Pro	Pro	Leu	Ser	Val
50						55					60				
Ser	Ile	His	Pro	Thr	Phe	Ile	Tyr	Tyr	Lys	His	Asn	Thr	Thr	Leu	Arg
65					70					75					80
Ile	Val	Phe	Gly	Thr	Tyr	Phe	Asp	Phe	Phe	Pro	Tyr	Arg	Lys	Asn	Lys
				85					90					95	
Asp	Gln	Ala	Phe	Glu	Gly	Glu	Asp	Trp	Glu	Ser	Ser	Leu	Asn	Val	Ser
		100						105					110		
Asp	Ala	Trp													
		115													

<210> 324

<211> 22

<212> PRT

<213> Homo sapiens

<400> 324

Thr	Lys	Arg	Gly	Ile	Lys	Lys	Leu	Val	Asn	Val	Cys	Leu	Lys	His	Pro
1				5					10					15	

Lys	Asn	Thr	Ser	Leu	Ser
				20	

<210> 325

<211> 26

<212> PRT

<213> Homo sapiens

<400> 325

Ser	Ile	His	Pro	Thr	Phe	Ile	Tyr	Tyr	Lys	His	Asn	Thr	Thr	Leu	Arg
1				5					10					15	

Ile	Val	Phe	Gly	Thr	Tyr	Phe	Asp	Phe	Phe
			20					25	

<210> 326

<211> 70

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (9)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (44)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

161

<222> (45)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 326

Gln Arg Pro His Pro Gln Pro Trp Xaa Pro Met Thr Leu Met Gly Thr
 1 5 10 15

Gly Ile Pro Val Phe Ala His Lys Met Leu Pro Phe Asp Pro Pro Cys
 20 25 30

His Leu Ser Cys Thr His Ile Asn Pro Lys Pro Xaa Xaa Pro Gln Gly
 35 40 45

Asp Glu Gln Lys Ser Gln Gly Thr Glu Glu Trp Cys Asp Arg Glu Gly
 50 55 60

Lys Lys Arg Arg Ser Ile
 65 70

<210> 327

<211> 21

<212> PRT

<213> Homo sapiens

<400> 327

Pro Met Thr Leu Met Gly Thr Gly Ile Pro Val Phe Ala His Lys Met
 1 5 10 15

Leu Pro Phe Asp Pro
 20

<210> 328

<211> 21

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (15)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (16)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 328

Pro Pro Cys His Leu Ser Cys Thr His Ile Asn Pro Lys Pro Xaa Xaa
 1 5 10 15

Pro Gln Gly Asp Glu
 20

<210> 329

<211> 21
 <212> PRT
 <213> Homo sapiens

162

<400> 329
 Glu Gln Lys Ser Gln Gly Thr Glu Glu Trp Cys Asp Arg Glu Gly Lys
 1 5 10 15
 Lys Arg Arg Ser Ile
 20

<210> 330
 <211> 70
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (64)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (65)
 <223> Xaa equals any of the naturally occurring L-amino acids

<400> 330
 Asp Glu Trp Gly Ala Gly Arg Arg Met Glu Trp Glu Asp Asn Leu Pro
 1 5 10 15
 Leu Glu Phe Ser Cys Pro Val Thr Lys Leu Leu Ser Val Pro Ser Trp
 20 25 30
 Thr Pro Leu Asp Ala Gln Met Leu Leu Leu Phe Phe Pro Ser Leu Ser
 35 40 45
 His His Ser Ser Val Pro Trp Leu Phe Cys Ser Ser Pro Cys Gly Xaa
 50 55 60
 Xaa Gly Leu Gly Phe Ile
 65 70

<210> 331
 <211> 21
 <212> PRT
 <213> Homo sapiens

<400> 331
 Glu Trp Glu Asp Asn Leu Pro Leu Glu Phe Ser Cys Pro Val Thr Lys
 1 5 10 15
 Leu Leu Ser Val Pro
 20

<210> 332

<211> 21
 <212> PRT
 <213> Homo sapiens

163

<400> 332
 Pro Ser Trp Thr Pro Leu Asp Ala Gln Met Leu Leu Leu Phe Phe Pro
 1 5 10 15
 Ser Leu Ser His His
 20

<210> 333
 <211> 21
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (15)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (16)
 <223> Xaa equals any of the naturally occurring L-amino acids

<400> 333
 His Ser Ser Val Pro Trp Leu Phe Cys Ser Ser Pro Cys Gly Xaa Xaa
 1 5 10 15

Gly Leu Gly Phe Ile
 20

<210> 334
 <211> 15
 <212> PRT
 <213> Homo sapiens

<400> 334
 Gln Gly Leu Ser His Ile Phe Trp Met Asn Glu Gln Thr Leu Lys
 1 5 10 15

<210> 335
 <211> 32
 <212> PRT
 <213> Homo sapiens

<400> 335
 Thr Leu Val Cys Leu Gly Val Ser Ser Glu Glu Gly Ser Cys Pro Arg
 1 5 10 15

Asp Val Thr Gly Pro Gly Cys Cys Phe Ser Leu Thr Leu Thr Gly Phe
 20 25 30

164

<210> 336
 <211> 233
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (57)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (62)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (78)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (79)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (80)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (231)
 <223> Xaa equals any of the naturally occurring L-amino acids

<400> 336
 Ala Asp Leu Ile Val Leu Trp His His His Pro Leu Trp Pro Gln His
 1 5 10 15

Leu Ala Leu Pro Ser Ser Gly Ala Ser His Asp His Val Glu Leu Thr
 20 25 30

Val Tyr Pro Lys Thr Val Ala Ala Ser Trp Leu Leu Glu Leu Ser Arg
 35 40 45

Pro Pro Ile Phe Cys Leu Phe Thr Xaa Pro Ala Leu Thr Xaa His Gly
 50 55 60

Leu Asp Arg Val Ala Ala Leu Val Glu Cys Thr Ile Trp Xaa Xaa Xaa
 65 70 75 80

Gly Met Trp Tyr Arg Arg Arg Tyr Ser Cys Cys Gln Phe Arg Asp Arg
 85 90 95

Ser Ile Arg Asp Val Phe Pro Glu Ala Val Met Leu Gln Gln His Leu

100 105 110
 Arg His Leu Ala Val Ala Thr Tyr Arg Cys Arg Arg Arg Ser Pro Cys
 115 120 125
 Lys Ala Pro Thr Val Glu Glu Ala Glu Gly Gly Lys Pro Arg Ala Val
 130 135 140
 Pro Ser Gly Thr Gly Phe Gln Lys His Gly Gln Glu Pro Gly Gly Ser
 145 150 155 160
 Thr Ser Pro His Trp Phe Trp Gly His Leu Gln Leu Leu Val Leu Ser
 165 170 175
 Val Asn Asn Arg Gln Leu Phe Val Gln Gly Arg Ala Gly Tyr Leu Glu
 180 185 190
 Met Thr Gly Leu Pro Cys Pro Lys Leu Leu Leu Thr Leu Leu Arg Gly
 195 200 205
 Leu Thr Pro Gly Val Gly His Gly Leu Cys Ala Tyr Arg Arg Gly Cys
 210 215 220
 Leu Ala Trp Arg Leu Asp Xaa Ala Ser
 225 230

<210> 337

<211> 176

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (70)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (71)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (92)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 337

Ile Leu Trp Arg Gln Ala Pro Glu Ala Pro His Cys Ser Gln Asp Ser
 1 5 10 15

Val Ser Ser Ser Pro Arg Leu Gln Glu Asp Leu Ala His Val Thr Gln
 20 25 30

Val Thr Arg His Pro His Phe Arg Ser Leu Pro Ser Ala Trp Cys Ser
 35 40 45

His Ser Ser Leu Leu Pro Val Ser Leu Pro Arg His Ala Leu Ala Thr

50 55 60
 Lys Ser Pro Asn Met Xaa Xaa Ser Ser Pro Ile Leu His Leu Ile Gln
 65 70 75 80
 Phe Thr Gly Gln Ile Ser Ser Pro Leu Gly Gly Xaa Val Gln Pro Pro
 85 90 95
 Gly Gln Thr Ala Ser Pro Ile Cys Thr Gln Pro Met Ser His Pro Arg
 100 105 110
 Arg Gln Ala Ser Gln Gln Cys Glu Gln Gln Leu Trp Thr Gly Gln Thr
 115 120 125
 Ser His Leu Gln Ile Pro Cys Pro Ala Leu Asn Lys Glu Leu Pro Val
 130 135 140
 Val Asp Thr Gln Asp Lys Glu Leu Gln Met Ser Pro Glu Pro Met Trp
 145 150 155 160
 Gly Cys Gly Pro Ser Arg Leu Leu Pro Met Leu Leu Glu Ser Cys Ala
 165 170 175

<210> 338
 <211> 34
 <212> PRT
 <213> Homo sapiens

<400> 338
 Met Leu Gln Gln His Leu Arg His Leu Ala Val Ala Thr Tyr Arg Cys
 1 5 10 15
 Arg Arg Arg Ser Pro Cys Lys Ala Pro Thr Val Glu Glu Ala Glu Gly
 20 25 30
 Gly Lys

<210> 339
 <211> 29
 <212> PRT
 <213> Homo sapiens

<400> 339
 Val Thr Gln Val Thr Arg His Pro His Phe Arg Ser Leu Pro Ser Ala
 1 5 10 15
 Trp Cys Ser His Ser Ser Leu Leu Pro Val Ser Leu Pro
 20 25

<210> 340
 <211> 28

<212> PRT
<213> Homo sapiens

167

<400> 340
Gly Gln Thr Ala Ser Pro Ile Cys Thr Gln Pro Met Ser His Pro Arg
1 5 10 15
Arg Gln Ala Ser Gln Gln Cys Glu Gln Gln Leu Trp
20 25

<210> 341
<211> 79
<212> PRT
<213> Homo sapiens

<400> 341
Phe Ile Thr Leu Arg Leu Gly Pro Lys Asn Met Ala Gly Val Leu Trp
1 5 10 15
Arg His Ser Asn Leu Gln Thr Pro His Tyr Ile Ser Trp Cys Pro Leu
20 25 30
Leu Asn Tyr Arg Glu Thr Gly Asn Cys Leu Leu His Val Ser Gly Phe
35 40 45
Leu Asn Ser Arg Leu Leu Ala Asn Cys Ser Gly Glu Ala Ser Gly Lys
50 55 60
Val Ile Gln Thr Leu Leu Trp Pro Gly Glu Ile Ser Ala Val Ala
65 70 75

<210> 342
<211> 82
<212> PRT
<213> Homo sapiens

<400> 342
Lys Ile Arg Thr Phe Leu Phe Ser Gly His Arg Leu Phe Ser Thr Gln
1 5 10 15
Gly Gln Ser Leu Thr Val Lys Ala His Thr Ala Phe Met Leu Ile Val
20 25 30
Lys Asn Leu Arg Tyr Phe Ile Ala Phe Lys Phe Leu Met Gly Ile Ser
35 40 45
Asp Ser Ser Glu Ile Gly Leu Val Met Gln Pro Leu Gln Lys Pro His
50 55 60
Thr Val Ile Leu Ile Arg Gly Ile Glu Phe Leu Ser Pro Gly Gly Val
65 70 75 80
Leu Pro

<210> 343
 <211> 26
 <212> PRT
 <213> Homo sapiens

168

<400> 343
 Met Ala Gly Val Leu Trp Arg His Ser Asn Leu Gln Thr Pro His Tyr
 1 5 10 15
 Ile Ser Trp Cys Pro Leu Leu Asn Tyr Arg
 20 25

<210> 344
 <211> 29
 <212> PRT
 <213> Homo sapiens

<400> 344
 Tyr Phe Ile Ala Phe Lys Phe Leu Met Gly Ile Ser Asp Ser Ser Glu
 1 5 10 15
 Ile Gly Leu Val Met Gln Pro Leu Gln Lys Pro His Thr
 20 25

<210> 345
 <211> 31
 <212> PRT
 <213> Homo sapiens

<400> 345
 Asp Val Leu Leu Pro Leu Leu Tyr Leu Leu Val Arg Lys His Ile Asn
 1 5 10 15
 Arg Ala Gly Ile Gly Asn Thr Phe Gln Gly Gly Ala Asn Cys Ile
 20 25 30

<210> 346
 <211> 99
 <212> PRT
 <213> Homo sapiens

<400> 346
 Met Cys Cys Cys Leu Cys Cys Thr Ser Trp Ser Gly Ser Thr Ser Thr
 1 5 10 15
 Glu Arg Val Ser Gly Thr Arg Phe Arg Glu Val Pro Thr Ala Ser Cys
 20 25 30
 Ser Ser Ser Ala Pro Ala Pro Ser Glu Leu Gly Ser Ser Leu Ser Val
 35 40 45
 Ala Ala Ala Ala Leu Leu Ser Leu Pro Pro Arg Ala Arg Leu Ala Leu
 50 55 60
 Pro Arg Leu Pro Arg Leu Pro Ser Gln Glu Asn Leu Arg Asn Pro Lys

Leu Pro Pro Arg Ser Ser Gly Ser Lys Phe Xaa Arg Tyr Gln Arg Pro

145 150 170 155 160
 Gly Pro Arg Val Gly Val His Leu His Lys Gly Ser Ser Glu Ile Arg
 165 170 175
 Glu Ala Gly Gly Pro Gln Leu Trp Pro Gln Cys Pro His Pro Val Asp
 180 185 190
 Leu Asp Val Leu Arg Thr Thr Gln His Cys Leu Gln Ser Glu Gly Pro
 195 200 205
 Thr Ser Val His Leu Ser Ser Val
 210 215

<210> 348

<211> 147

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (34)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (39)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 348

Glu Val Glu Glu Ala Glu Leu Ala Ala Ala Leu Pro Met Glu Pro Arg
 1 5 10 15

Ala Ser Ile Ala Gly Ala Ser Gly Ala Ala Asp Met His Phe Cys Pro
 20 25 30

Ala Xaa Gly Thr His Arg Xaa Ala Tyr Pro Gln Glu Gly Ser Thr Tyr
 35 40 45

Ala Thr Glu Leu Glu Arg Thr Lys Ala Pro Gly Ala Trp Lys Phe Pro
 50 55 60

Trp Gly Pro Leu Gly Phe Leu Arg Phe Ser Trp Leu Gly Arg Arg Gly
 65 70 75 80

Ser Leu Gly Ser Ala Ser Arg Ala Leu Gly Gly Arg Leu Arg Arg Ala
 85 90 95

Ala Ala Ala Thr Glu Arg Glu Glu Pro Ser Ser Asp Gly Ala Gly Ala
 100 105 110

Glu Asp Glu His Asp Ala Val Gly Thr Ser Leu Lys Arg Val Pro Asp
 115 120 125

Thr Arg Ser Val Asp Val Leu Pro Asp Gln Glu Val Gln Gln Arg Gln
 130 135 140

Gln His Ile
145

171

<210> 349
<211> 31
<212> PRT
<213> Homo sapiens

<400> 349
Arg Arg Ile Ser Gly Ile Pro Arg Asp Pro Arg Gly Thr Ser Lys His
1 5 10 15

Leu Glu Leu Leu Ser Phe Leu Val Leu Trp His Arg Cys Cys Leu
20 25 30

<210> 350
<211> 29
<212> PRT
<213> Homo sapiens

<400> 350
Arg Thr Lys Ala Pro Gly Ala Trp Lys Phe Pro Trp Gly Pro Leu Gly
1 5 10 15

Phe Leu Arg Phe Ser Trp Leu Gly Arg Arg Gly Ser Leu
20 25

<210> 351
<211> 11
<212> PRT
<213> Homo sapiens

<400> 351
Pro Arg Leu Ala Gln Leu Arg Leu Leu Ser Leu
1 5 10

<210> 352
<211> 178
<212> PRT
<213> Homo sapiens

<400> 352
Gln Ser Asp Phe Arg Glu Met Asn Gln Thr Asn Ser Thr Ser Asn Ala
1 5 10 15

Ala Lys Ala Arg Glu Ala Gln Gln Gly Arg Gly Arg Asp Arg Glu Ala
20 25 30

Ile Phe Ser Ser Ser Ala Leu Glu His Leu Val Cys Tyr Leu Gln Ala
35 40 45

Tyr Lys His Thr Leu Leu Phe Ile Arg Ser Leu Asn Glu His Gly Leu
50 55 60

172

Gln	Gln	Leu	Leu	Phe	Gln	Trp	Arg	Asp	Gly	Leu	Phe	Gly	Asn	Trp	Tyr
65					70					75					80
Phe	Arg	Ile	Pro	Ile	Leu	Leu	Phe	Phe	Thr	Gly	Phe	His	Cys	Tyr	His
				85					90					95	
Leu	Ser	Cys	Pro	His	Leu	Pro	Cys	Ala	Gln	Arg	Gln	Ser	Ser	Arg	Gly
			100					105					110		
Thr	Val	Pro	Tyr	Val	Leu	Cys	Pro	His	Pro	His	His	His	Leu	His	His
		115					120					125			
Tyr	Ser	Trp	Phe	Pro	Phe	Leu	Ile	Pro	Val	Leu	His	Thr	Leu	Pro	Lys
	130					135					140				
Leu	Gln	Pro	Lys	Phe	His	Gly	Arg	Pro	Glu	Gln	Pro	Leu	Asn	Leu	Leu
145					150					155					160
Gln	Val	Lys	Pro	Thr	Ser	Gly	Thr	Ile	Ala	Ser	Ala	Glu	Gln	Val	Trp
				165					170					175	

Val Lys

<210> 353

<211> 29

<212> PRT

<213> Homo sapiens

<400> 353

Val	Cys	Tyr	Leu	Gln	Ala	Tyr	Lys	His	Thr	Leu	Leu	Phe	Ile	Arg	Ser
1				5					10					15	

Leu	Asn	Glu	His	Gly	Leu	Gln	Gln	Leu	Leu	Phe	Gln	Trp
		20					25					

<210> 354

<211> 32

<212> PRT

<213> Homo sapiens

<400> 354

Val	Pro	Tyr	Val	Leu	Cys	Pro	His	Pro	His	His	His	Leu	His	His	Tyr
1				5					10				15		

Ser	Trp	Phe	Pro	Phe	Leu	Ile	Pro	Val	Leu	His	Thr	Leu	Pro	Lys	Leu
			20					25					30		

<210> 355

<211> 31

<212> PRT

<213> Homo sapiens

(73)

<400> 355

Glu Ser Glu Arg Ala Val Val Tyr Leu Ile Thr Gly Ala Leu Phe Ile
 1 5 10 15

Val Ser Ser Cys Val Leu Cys Phe Leu Pro Ser Ser Arg Arg Glu
 20 25 30

<210> 356

<211> 249

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (4)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (221)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 356

Met Trp Val Xaa Gly Glu Glu Val Leu Gly Ser His Ala Ala Ser Pro
 1 5 10 15

Ala Phe Leu His Arg Cys Phe Ser Glu Glu Ser Cys Val Ser Ile Pro
 20 25 30

Glu Val Glu Gly Tyr Val Val Val Leu Gln Pro Asp Ala Pro Gln Ile
 35 40 45

Leu Leu Ser Gly Thr Ala His Phe Ala Arg Pro Ala Val Asp Phe Glu
 50 55 60

Gly Thr Asn Gly Val Pro Leu Phe Pro Asp Leu Gln Ile Thr Cys Ser
 65 70 75 80

Ile Ser His Gln Val Glu Ala Lys Lys Asp Glu Ser Trp Gln Gly Thr
 85 90 95

Val Thr Asp Thr Arg Met Ser Asp Glu Ile Val His Asn Leu Asp Gly
 100 105 110

Cys Glu Ile Ser Leu Val Gly Asp Asp Leu Asp Pro Glu Arg Glu Ser
 115 120 125

Leu Leu Leu Asp Thr Thr Ser Leu Gln Gln Arg Gly Leu Glu Leu Thr
 130 135 140

Asn Thr Ser Ala Tyr Leu Thr Ile Ala Gly Val Glu Ser Ile Thr Val
 145 150 155 160

Tyr Glu Glu Ile Leu Arg Gln Ala Arg Tyr Arg Leu Arg His Gly Ala
 165 170 175

174

Ala Leu Tyr Thr Arg Lys Phe Arg Leu Ser Cys Ser Glu Met Asn Gly
 180 185 190

Arg Tyr Ser Ser Asn Glu Phe Ile Val Glu Val Asn Val Leu His Ser
 195 200 205

Met Asn Arg Val Ala His Pro Ser His Val Leu Ser Xaa Gln Gln Phe
 210 215 220

Leu His Arg Gly His Gln Pro Pro Pro Glu Met Ala Gly His Ser Leu
 225 230 235 240

Ala Ser Ser His Arg Asn Ser Ser Thr
 245

<210> 357
 <211> 23
 <212> PRT
 <213> Homo sapiens

<400> 357
 Leu Gly Ser His Ala Ala Ser Pro Ala Phe Leu His Arg Cys Phe Ser
 1 5 10 15

Glu Glu Ser Cys Val Ser Ile
 20

<210> 358
 <211> 29
 <212> PRT
 <213> Homo sapiens

<400> 358
 Gly Tyr Val Val Val Leu Gln Pro Asp Ala Pro Gln Ile Leu Leu Ser
 1 5 10 15

Gly Thr Ala His Phe Ala Arg Pro Ala Val Asp Phe Glu
 20 25

<210> 359
 <211> 26
 <212> PRT
 <213> Homo sapiens

<400> 359
 Ile Thr Cys Ser Ile Ser His Gln Val Glu Ala Lys Lys Asp Glu Ser
 1 5 10 15

Trp Gln Gly Thr Val Thr Asp Thr Arg Met
 20 25

<210> 360
 <211> 29
 <212> PRT

<213> Homo sapiens

175

<400> 360

Asn Leu Asp Gly Cys Glu Ile Ser Leu Val Gly Asp Asp Leu Asp Pro
1 5 10 15

Glu Arg Glu Ser Leu Leu Leu Asp Thr Thr Ser Leu Gln
20 25

<210> 361

<211> 23

<212> PRT

<213> Homo sapiens

<400> 361

Ser Ala Tyr Leu Thr Ile Ala Gly Val Glu Ser Ile Thr Val Tyr Glu
1 5 10 15

Glu Ile Leu Arg Gln Ala Arg
20

<210> 362

<211> 26

<212> PRT

<213> Homo sapiens

<400> 362

Arg Leu Ser Cys Ser Glu Met Asn Gly Arg Tyr Ser Ser Asn Glu Phe
1 5 10 15

Ile Val Glu Val Asn Val Leu His Ser Met
20 25

<210> 363

<211> 25

<212> PRT

<213> Homo sapiens

<400> 363

Gln Gln Phe Leu His Arg Gly His Gln Pro Pro Pro Glu Met Ala Gly
1 5 10 15

His Ser Leu Ala Ser Ser His Arg Asn
20 25

<210> 364

<211> 299

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (52)

<223> Xaa equals any of the naturally occurring L-amino acids

176

<400> 364

```

Met Ala Asp Ser Glu Thr Phe Ile Ser Leu Glu Glu Cys Arg Gly His
 1              5              10              15

Lys Arg Ala Arg Lys Arg Thr Ser Met Glu Thr Ala Leu Ala Leu Glu
      20              25              30

Lys Leu Phe Pro Lys Gln Cys Gln Val Leu Gly Ile Val Thr Pro Gly
      35              40              45

Ile Val Val Xaa Pro Met Gly Ser Gly Ser Asn Arg Pro Gln Glu Ile
      50              55              60

Glu Ile Gly Glu Ser Gly Phe Ala Leu Leu Phe Pro Gln Ile Glu Gly
      65              70              75              80

Ile Lys Ile Gln Pro Phe His Phe Ile Lys Asp Pro Lys Asn Leu Thr
      85              90              95

Leu Glu Arg His Gln Leu Thr Glu Val Gly Leu Leu Asp Asn Pro Glu
      100             105             110

Leu Arg Val Val Leu Val Phe Gly Tyr Asn Cys Cys Lys Val Gly Ala
      115             120             125

Ser Asn Tyr Leu Gln Gln Val Val Ser Thr Phe Ser Asp Met Asn Ile
      130             135             140

Ile Leu Ala Gly Gly Gln Val Asp Asn Leu Ser Ser Leu Thr Ser Glu
      145             150             155             160

Lys Asn Pro Leu Asp Ile Asp Ala Ser Gly Val Val Gly Leu Ser Phe
      165             170             175

Ser Gly His Arg Ile Gln Ser Ala Thr Val Leu Leu Asn Glu Asp Val
      180             185             190

Ser Asp Glu Lys Thr Ala Glu Ala Ala Met Gln Arg Leu Lys Ala Ala
      195             200             205

Asn Ile Pro Glu His Asn Thr Ile Gly Phe Met Phe Ala Cys Val Gly
      210             215             220

Arg Gly Phe Gln Tyr Tyr Arg Ala Lys Gly Asn Val Glu Ala Asp Ala
      225             230             235             240

Phe Arg Lys Phe Phe Pro Ser Val Pro Leu Phe Gly Phe Phe Gly Asn
      245             250             255

Gly Glu Ile Gly Cys Asp Arg Ile Val Thr Gly Asn Phe Ile Leu Arg
      260             265             270

Lys Cys Asn Glu Val Lys Asp Asp Asp Leu Phe His Ser Tyr Thr Thr
      275             280             285

Ile Met Ala Leu Ile His Leu Gly Ser Ser Lys
      290             295

```

177

<210> 365
<211> 21
<212> PRT
<213> Homo sapiens

<400> 365
His Lys Arg Ala Arg Lys Arg Thr Ser Met Glu Thr Ala Leu Ala Leu
1 5 10 15
Glu Lys Leu Phe Pro
20

<210> 366
<211> 24
<212> PRT
<213> Homo sapiens

<400> 366
Met Gly Ser Gly Ser Asn Arg Pro Gln Glu Ile Glu Ile Gly Glu Ser
1 5 10 15
Gly Phe Ala Leu Leu Phe Pro Gln
20

<210> 367
<211> 22
<212> PRT
<213> Homo sapiens

<400> 367
Phe His Phe Ile Lys Asp Pro Lys Asn Leu Thr Leu Glu Arg His Gln
1 5 10 15
Leu Thr Glu Val Gly Leu
20

<210> 368
<211> 23
<212> PRT
<213> Homo sapiens

<400> 368
Phe Gly Tyr Asn Cys Cys Lys Val Gly Ala Ser Asn Tyr Leu Gln Gln
1 5 10 15
Val Val Ser Thr Phe Ser Asp
20

<210> 369
<211> 20
<212> PRT
<213> Homo sapiens

<400> 369

178

Thr Ser Glu Lys Asn Pro Leu Asp Ile Asp Ala Ser Gly Val Val Gly
 1 5 10 15

Leu Ser Phe Ser
 20

<210> 370

<211> 26

<212> PRT

<213> Homo sapiens

<400> 370

Asn Glu Asp Val Ser Asp Glu Lys Thr Ala Glu Ala Ala Met Gln Arg
 1 5 10 15

Leu Lys Ala Ala Asn Ile Pro Glu His Asn
 20 25

<210> 371

<211> 25

<212> PRT

<213> Homo sapiens

<400> 371

Tyr Tyr Arg Ala Lys Gly Asn Val Glu Ala Asp Ala Phe Arg Lys Phe
 1 5 10 15

Phe Pro Ser Val Pro Leu Phe Gly Phe
 20 25

<210> 372

<211> 26

<212> PRT

<213> Homo sapiens

<400> 372

Ile Gly Cys Asp Arg Ile Val Thr Gly Asn Phe Ile Leu Arg Lys Cys
 1 5 10 15

Asn Glu Val Lys Asp Asp Asp Leu Phe His
 20 25

<210> 373

<211> 341

<212> PRT

<213> Homo sapiens

<400> 373

Met Pro Lys Arg Lys Val Thr Phe Gln Gly Val Gly Asp Glu Glu Asp
 1 5 10 15

Glu Asp Glu Ile Ile Val Pro Lys Lys Lys Leu Val Asp Pro Val Ala

20				25				30							
Gly	Ser	Gly	Gly	Pro	Gly	Ser	Arg	Phe	Lys	Gly	Lys	His	Ser	Leu	Asp
35				40				45							
Ser	Asp	Glu	Glu	Glu	Asp	Asp	Asp	Asp	Gly	Gly	Ser	Ser	Lys	Tyr	Asp
50				55				60							
Ile	Leu	Ala	Ser	Glu	Asp	Val	Glu	Gly	Gln	Glu	Ala	Ala	Thr	Leu	Pro
65				70				75				80			
Ser	Glu	Gly	Gly	Val	Arg	Ile	Thr	Pro	Phe	Asn	Leu	Gln	Glu	Glu	Met
				85				90				95			
Glu	Glu	Gly	His	Phe	Asp	Ala	Asp	Gly	Asn	Tyr	Phe	Leu	Asn	Arg	Asp
100								105				110			
Ala	Gln	Ile	Arg	Asp	Ser	Trp	Leu	Asp	Asn	Ile	Asp	Trp	Val	Lys	Ile
115				120				125							
Arg	Glu	Arg	Pro	Pro	Gly	Gln	Arg	Gln	Ala	Ser	Asp	Ser	Glu	Glu	Glu
130				135				140							
Asp	Ser	Leu	Gly	Gln	Thr	Ser	Met	Ser	Ala	Gln	Ala	Leu	Leu	Glu	Gly
145				150				155				160			
Leu	Leu	Glu	Leu	Leu	Leu	Pro	Arg	Glu	Thr	Val	Ala	Gly	Ala	Leu	Arg
				165				170				175			
Arg	Leu	Gly	Ala	Arg	Gly	Gly	Gly	Lys	Gly	Arg	Lys	Gly	Pro	Gly	Gln
180								185				190			
Pro	Ser	Ser	Pro	Gln	Arg	Leu	Asp	Arg	Leu	Ser	Gly	Leu	Ala	Asp	Gln
195				200				205							
Met	Val	Ala	Arg	Gly	Asn	Leu	Gly	Val	Tyr	Gln	Glu	Thr	Arg	Glu	Arg
210				215				220							
Leu	Ala	Met	Arg	Leu	Lys	Gly	Leu	Gly	Cys	Gln	Thr	Leu	Gly	Pro	His
225				230				235				240			
Asn	Pro	Thr	Pro	Pro	Pro	Ser	Leu	Asp	Met	Phe	Ala	Glu	Glu	Leu	Ala
245				250				255							
Glu	Glu	Glu	Leu	Glu	Thr	Pro	Thr	Pro	Thr	Gln	Arg	Gly	Glu	Ala	Glu
260				265				270							
Ser	Arg	Gly	Asp	Gly	Leu	Val	Asp	Val	Met	Trp	Glu	Tyr	Lys	Trp	Glu
275				280				285							
Asn	Thr	Gly	Asp	Ala	Glu	Leu	Tyr	Gly	Pro	Phe	Thr	Ser	Ala	Gln	Met
290				295				300							
Gln	Thr	Trp	Val	Ser	Glu	Gly	Tyr	Phe	Pro	Asp	Gly	Val	Tyr	Cys	Arg
305				310				315				320			
Lys	Leu	Asp	Pro	Pro	Gly	Gly	Gln	Phe	Tyr	Asn	Ser	Lys	Arg	Ile	Asp
325				330				335							

(80

Phe Asp Leu Tyr Thr
340

<210> 374
<211> 24
<212> PRT
<213> Homo sapiens

<400> 374
Thr Phe Gln Gly Val Gly Asp Glu Glu Asp Glu Ile Ile Val
1 5 10 15

Pro Lys Lys Lys Leu Val Asp Pro
20

<210> 375
<211> 27
<212> PRT
<213> Homo sapiens

<400> 375
Pro Gly Ser Arg Phe Lys Gly Lys His Ser Leu Asp Ser Asp Glu Glu
1 5 10 15

Glu Asp Asp Asp Asp Gly Gly Ser Ser Lys Tyr
20 25

<210> 376
<211> 25
<212> PRT
<213> Homo sapiens

<400> 376
Glu Ala Ala Thr Leu Pro Ser Glu Gly Gly Val Arg Ile Thr Pro Phe
1 5 10 15

Asn Leu Gln Glu Glu Met Glu Glu Gly
20 25

<210> 377
<211> 29
<212> PRT
<213> Homo sapiens

<400> 377
Phe Leu Asn Arg Asp Ala Gln Ile Arg Asp Ser Trp Leu Asp Asn Ile
1 5 10 15

Asp Trp Val Lys Ile Arg Glu Arg Pro Pro Gly Gln Arg
20 25

<210> 378

<211> 26
<212> PRT
<213> Homo sapiens

(8)

<400> 378
Ser Leu Gly Gln Thr Ser Met Ser Ala Gln Ala Leu Leu Glu Gly Leu
1 5 10 15

Leu Glu Leu Leu Leu Pro Arg Glu Thr Val
20 25

<210> 379
<211> 28
<212> PRT
<213> Homo sapiens

<400> 379
Arg Gly Gly Gly Lys Gly Arg Lys Gly Pro Gly Gln Pro Ser Ser Pro
1 5 10 15

Gln Arg Leu Asp Arg Leu Ser Gly Leu Ala Asp Gln
20 25

<210> 380
<211> 24
<212> PRT
<213> Homo sapiens

<400> 380
Gln Glu Thr Arg Glu Arg Leu Ala Met Arg Leu Lys Gly Leu Gly Cys
1 5 10 15

Gln Thr Leu Gly Pro His Asn Pro
20

<210> 381
<211> 28
<212> PRT
<213> Homo sapiens

<400> 381
Asp Met Phe Ala Glu Glu Leu Ala Glu Glu Glu Leu Glu Thr Pro Thr
1 5 10 15

Pro Thr Gln Arg Gly Glu Ala Glu Ser Arg Gly Asp
20 25

<210> 382
<211> 30
<212> PRT
<213> Homo sapiens

<400> 382
Glu Leu Tyr Gly Pro Phe Thr Ser Ala Gln Met Gln Thr Trp Val Ser

1 5 10 15
 Glu Gly Tyr Phe Pro Asp Gly Val Tyr Cys Arg Lys Leu Asp
 20 25 30

<210> 383
 <211> 14
 <212> PRT
 <213> Homo sapiens

<400> 383
 Pro His Ser Ser Arg Val Ser Phe Leu Gln Ser Leu Ser Phe
 1 5 10

<210> 384
 <211> 141
 <212> PRT
 <213> Homo sapiens

<400> 384
 Arg Gly Gln Pro Arg Pro Cys Val Ser Gly Val Cys Leu Ser Pro His
 1 5 10 15

Ser Arg Phe Trp Glu Cys Cys Ser Phe Tyr Leu Gln Gly Leu Pro Ala
 20 25 30

Leu Arg Cys Ser Arg Thr Pro Pro Gly Cys His Phe Phe Arg Val Phe
 35 40 45

Pro Ser Cys Pro Phe Ser Ser Ser Arg Ser Pro Ser Cys Phe Thr His
 50 55 60

Ile Cys Pro Val Val Arg Ile Gln Phe Ser Arg Ala Leu Trp Val Ser
 65 70 75 80

Thr Cys Leu Val Leu Ala Ile Thr Pro Gly Lys Trp Leu Leu Pro Glu
 85 90 95

Asp Arg Ala Leu Ser Leu Met Leu Leu Ala Ser Leu Gln Cys Cys Pro
 100 105 110

Pro Pro Phe Gly Ala Trp Trp Met Gln Val Leu Thr His Lys Gly Arg
 115 120 125

Gln Ala Gly Leu Gly Pro Gly Val Ser Ser Arg Pro Leu
 130 135 140

<210> 385
 <211> 133
 <212> PRT
 <213> Homo sapiens

<400> 385
 Ser Asn Ile Lys Ser Leu Pro Pro Thr Asn Ser Leu Ser Leu Leu Arg
 1 5 10 15

123

Ala Gln Thr Gly Thr Asp Cys Ala Val Ser Pro Gly Leu Ala Gly Pro
 20 25 30

Cys His Gln Arg Gly Leu Glu Asp Thr Pro Gly Pro Arg Pro Ala Cys
 35 40 45

Leu Pro Leu Cys Val Ser Thr Cys Ile His Gln Ala Pro Lys Gly Gly
 50 55 60

Gly Gln His Trp Arg Glu Ala Ser Ser Ile Arg Asp Arg Ala Leu Ser
 65 70 75 80

Ser Gly Arg Ser His Phe Pro Gly Val Met Ala Lys Thr Lys His Val
 85 90 95

Asp Thr His Asn Ala Arg Glu Asn Trp Ile Arg Thr Thr Gly Gln Met
 100 105 110

Trp Val Lys His Glu Gly Glu Arg Glu Glu Glu Lys Gly His Glu Gly
 115 120 125

Lys Thr Leu Lys Lys
 130

<210> 386

<211> 25

<212> PRT

<213> Homo sapiens

<400> 386

Val Cys Leu Ser Pro His Ser Arg Phe Trp Glu Cys Cys Ser Phe Tyr
 1 5 10 15

Leu Gln Gly Leu Pro Ala Leu Arg Cys
 20 25

<210> 387

<211> 27

<212> PRT

<213> Homo sapiens

<400> 387

Gln Phe Ser Arg Ala Leu Trp Val Ser Thr Cys Leu Val Leu Ala Ile
 1 5 10 15

Thr Pro Gly Lys Trp Leu Leu Pro Glu Asp Arg
 20 25

<210> 388

<211> 27

<212> PRT

<213> Homo sapiens

<400> 388

184

Ser	Leu	Ser	Leu	Leu	Arg	Ala	Gln	Thr	Gly	Thr	Asp	Cys	Ala	Val	Ser
1				5					10					15	

Pro	Gly	Leu	Ala	Gly	Pro	Cys	His	Gln	Arg	Gly
		20						25		

<210> 389
<211> 28
<212> PRT
<213> Homo sapiens

<400> 389
Ser Gly Arg Ser His Phe Pro Gly Val Met Ala Lys Thr Lys His Val
1 5 10 15

Asp	Thr	His	Asn	Ala	Arg	Glu	Asn	Trp	Ile	Arg	Thr
		20					25				

<210> 390
<211> 20
<212> PRT
<213> Homo sapiens

<400> 390
Ala Arg Val Glu Val Gln Gly Gln Gly Pro Gly Ala Lys Val Asp Ala
1 5 10 15

Gly	Glu	Gly	Gln
		20	

<210> 391
<211> 121
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (46)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (66)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (98)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (121)
<223> Xaa equals any of the naturally occurring L-amino acids

<400> 391

Trp Val Val Leu Ser Gln Leu Gln Ala Gln Gly Val Ala Gly Met Met
 1 5 10 15

Cys Ser Tyr Pro Glu Gly Gln Lys Lys Gly Lys Glu Ala Thr Arg Ser
 20 25 30

His Arg Trp Val Pro Arg Ser Leu Pro Gly Met Gly Ser Xaa Leu Ala
 35 40 45

Ala Pro His Ser Asn Pro Trp Leu Ala Pro Leu Ala Leu Leu Glu Ile
 50 55 60

Pro Xaa Pro Val Leu Cys Glu Trp Lys Arg Lys Leu Ile Ala Leu Glu
 65 70 75 80

Glu Val Ser Glu Cys Arg Pro Gly Val Gly Gly Gly Gly Gly Phe Leu
 85 90 95

Ser Xaa Cys Arg Arg Gly His Leu Ser Phe Leu Ser Gly Ala Pro Tyr
 100 105 110

Pro Leu Phe Pro Ile Ser Pro Leu Xaa
 115 120

<210> 392

<211> 206

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (105)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (127)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (131)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (180)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 392

Glu Leu Arg His Gly Gly Pro Arg Gln Val Lys Asp Ser Phe Leu Asp
 1 5 10 15

Tyr Met Gly Tyr Pro Asp Glu Asp Arg Ala Gly Pro Pro Ser Arg Trp
 20 25 30

186

Phe Pro Arg Glu Arg Phe Leu Ser Pro Pro Thr Val Val Pro Leu Cys
 35 40 45
 Val Glu Leu Arg Leu Gly Phe Glu Ser Gly Met Gly Trp Gly Val Pro
 50 55 60
 Gly Ser Ser His Ser Glu Gly Gly Pro Glu Ala Arg Trp Pro Leu Ile
 65 70 75 80
 Ala Pro Met Tyr Thr Val Thr Gln Trp Phe Gln Arg Pro Asn Ser Gly
 85 90 95
 Arg Gly Pro Gln Pro Pro Pro Gln Xaa Arg Gly Glu Ile Gly Lys Arg
 100 105 110
 Gly Tyr Gly Ala Pro Glu Arg Lys Leu Arg Trp Pro Leu Leu Xaa Trp
 115 120 125
 Glu Arg Xaa Pro Pro Pro Pro Pro Thr Pro Gly Arg His Ser Glu Thr
 130 135 140
 Ser Ser Ser Ala Ile Ser Phe Leu Phe His Ser Gln Arg Thr Gly Trp
 145 150 155 160
 Gly Ile Ser Ser Ser Ala Asn Gly Ala Ser Gln Gly Leu Leu Trp Gly
 165 170 175
 Ala Ala Arg Xaa Leu Pro Ile Pro Gly Arg Asp Leu Gly Thr His Leu
 180 185 190
 Trp Asp Leu Val Ala Ser Phe Pro Phe Phe Cys Pro Ser Gly
 195 200 205

<210> 393

<211> 24

<212> PRT

<213> Homo sapiens

<400> 393

Pro Glu Gly Gln Lys Lys Gly Lys Glu Ala Thr Arg Ser His Arg Trp
 1 5 10 15

Val Pro Arg Ser Leu Pro Gly Met
 20

<210> 394

<211> 26

<212> PRT

<213> Homo sapiens

<400> 394

Leu Arg Leu Gly Phe Glu Ser Gly Met Gly Trp Gly Val Pro Gly Ser
 1 5 10 15

Ser His Ser Glu Gly Gly Pro Glu Ala Arg
 20 25

187

<210> 395
 <211> 24
 <212> PRT
 <213> Homo sapiens

<400> 395
 His Ser Gln Arg Thr Gly Trp Gly Ile Ser Ser Ser Ala Asn Gly Ala
 1 5 10 15
 Ser Gln Gly Leu Leu Trp Gly Ala
 20

<210> 396
 <211> 54
 <212> PRT
 <213> Homo sapiens

<400> 396
 Phe Ile Met Lys Leu Leu Tyr Gln Leu Leu Met Leu Thr Thr Ser Ser
 1 5 10 15
 Ser Tyr Ser Leu Ile Thr His Leu Cys Tyr Ser Ile Phe Leu Cys Ser
 20 25 30
 Phe Tyr Phe His Phe Pro Cys Asn Val Ser Leu Phe Val Leu Ile Ser
 35 40 45
 Glu Glu Phe Ile Tyr Asp
 50

<210> 397
 <211> 21
 <212> PRT
 <213> Homo sapiens

<400> 397
 Leu Met Leu Thr Thr Ser Ser Ser Tyr Ser Leu Ile Thr His Leu Cys
 1 5 10 15
 Tyr Ser Ile Phe Leu
 20

<210> 398
 <211> 21
 <212> PRT
 <213> Homo sapiens

<400> 398
 Leu Cys Ser Phe Tyr Phe His Phe Pro Cys Asn Val Ser Leu Phe Val
 1 5 10 15
 Leu Ile Ser Glu Glu
 20

188

<210> 399
<211> 53
<212> PRT
<213> Homo sapiens

<400> 399
Met Arg Lys Asn Ile Phe Ala Ile Leu Asp Lys Met Leu Thr Cys Leu
1 5 10 15
Ile Ile Asn Glu Leu Phe Arg Asn Gln Tyr Lys Glu Thr Asn Ile Thr
20 25 30
Arg Glu Val Lys Ile Lys Gly Thr Glu Glu Asn Gly Ile Ala Gln Met
35 40 45
Ser Tyr Lys Ala Ile
50

<210> 400
<211> 21
<212> PRT
<213> Homo sapiens

<400> 400
Asp Lys Met Leu Thr Cys Leu Ile Ile Asn Glu Leu Phe Arg Asn Gln
1 5 10 15
Tyr Lys Glu Thr Asn
20

<210> 401
<211> 21
<212> PRT
<213> Homo sapiens

<400> 401
Asn Ile Thr Arg Glu Val Lys Ile Lys Gly Thr Glu Glu Asn Gly Ile
1 5 10 15
Ala Gln Met Ser Tyr
20

<210> 402
<211> 7
<212> PRT
<213> Homo sapiens

<400> 402
Gly Ile Ser Glu Arg Lys Pro
1 5

<210> 403

<211> 25
 <212> PRT
 <213> Homo sapiens

189

<400> 403
 Gln Ser Pro Ala Val Ser Tyr Thr Val Thr Ser Gln Val Pro Trp Gly
 1 5 10 15
 Leu Gly Leu Leu Ala Gly Glu Lys Arg
 20 25

<210> 404
 <211> 100
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (96)
 <223> Xaa equals any of the naturally occurring L-amino acids

<400> 404
 Leu Pro Ser His Pro Leu Arg Pro Leu Thr Phe Ser Ser Ala Met Cys
 1 5 10 15
 Met His Leu Pro Pro Pro Leu Cys Arg Arg Ala Ala Leu Ser Ala Pro
 20 25 30
 Phe Ala Thr Gln His Arg Pro Trp Ser Val Ala Ala Ala Cys Leu Pro
 35 40 45
 Arg Ile His Gln Asn Pro Leu Asp Ala Glu Tyr Pro Ser Gly Cys Cys
 50 55 60
 Arg Met Ser Phe Leu Pro Ala Ala Cys Ser Asn Ile Tyr Ser Gln Glu
 65 70 75 80
 Cys His Tyr Thr Leu Met Ser His Ser Glu Ala Ser Thr Leu Gln Xaa
 85 90 95
 Ala Gln Leu Leu
 100

<210> 405
 <211> 76
 <212> PRT
 <213> Homo sapiens

<400> 405
 Met Leu Leu Gln Ala Ala Gly Arg Lys Leu Met Arg Gln Gln Pro Asp
 1 5 10 15
 Gly Tyr Ser Ala Ser Arg Gly Phe Trp Trp Met Arg Gly Arg Gln Ala
 20 25 30
 Ala Ala Thr Leu His Gly Arg Cys Trp Val Ala Lys Gly Ala Asp Ser

35

40

190

45

Ala Ala Leu Arg Gln Arg Gly Gly Gly Arg Cys Met His Ile Ala Asp
 50 55 60

Glu Lys Val Arg Gly Leu Ser Gly Cys Asp Gly Ser
 65 70 75

<210> 406

<211> 25

<212> PRT

<213> Homo sapiens

<400> 406

Leu Cys Arg Arg Ala Ala Leu Ser Ala Pro Phe Ala Thr Gln His Arg
 1 5 10 15

Pro Trp Ser Val Ala Ala Ala Cys Leu
 20 25

<210> 407

<211> 24

<212> PRT

<213> Homo sapiens

<400> 407

Arg Gly Phe Trp Trp Met Arg Gly Arg Gln Ala Ala Ala Thr Leu His
 1 5 10 15

Gly Arg Cys Trp Val Ala Lys Gly
 20

<210> 408

<211> 23

<212> PRT

<213> Homo sapiens

<400> 408

Gln Arg Gly Gly Gly Arg Cys Met His Ile Ala Asp Glu Lys Val Arg
 1 5 10 15

Gly Leu Ser Gly Cys Asp Gly
 20

<210> 409

<211> 106

<212> PRT

<213> Homo sapiens

<400> 409

Thr His Pro Ser His Pro Ser Ile Val Ile Gln Ser Thr Val Ser Leu
 1 5 10 15

Cys Leu Thr Ala Ser Ser Arg Arg Lys Lys Ser Asp Cys Leu Ser Leu

20 25 30
 Cys Gln Val Ser Cys Ser Gln Arg Pro Gly Ser His Lys Thr Asn Val
 35 40 45
 Ala Trp Gly Phe Leu Met Ser Arg Val His Phe Ser Val Arg Trp Val
 50 55 60
 Ser Gly Gly Arg Gly Ile Thr Gly Ala Ile Cys Lys Glu Ser Ser Leu
 65 70 75 80
 Pro Cys Lys Glu Ile Gln Gly Lys Ala Cys Tyr Phe Cys His His Pro
 85 90 95
 Ala Gln Gln Ser Thr Pro Phe Ser His Ile
 100 105

<210> 410
 <211> 27
 <212> PRT
 <213> Homo sapiens

<400> 410
 Val Ile Gln Ser Thr Val Ser Leu Cys Leu Thr Ala Ser Ser Arg Arg
 1 5 10 15

Lys Lys Ser Asp Cys Leu Ser Leu Cys Gln Val
 20 25

<210> 411
 <211> 26
 <212> PRT
 <213> Homo sapiens

<400> 411
 Ile Cys Lys Glu Ser Ser Leu Pro Cys Lys Glu Ile Gln Gly Lys Ala
 1 5 10 15

Cys Tyr Phe Cys His His Pro Ala Gln Gln
 20 25

<210> 412
 <211> 188
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (140)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (149)
 <223> Xaa equals any of the naturally occurring L-amino acids

<400> 412

(92)

Ser Leu Gln Val Leu Arg Thr Leu Gly Ser Lys Cys Gly Asp Phe Leu
 1 5 10 15

Arg Ser Arg Phe Cys Lys Asp Val Leu Pro Lys Leu Ala Gly Ser Leu
 20 25 30

Val Thr Gln Ala Pro Ile Ser Ala Arg Ala Gly Pro Val Tyr Ser His
 35 40 45

Thr Leu Ala Phe Lys Leu Gln Leu Ala Val Leu Gln Gly Leu Gly Pro
 50 55 60

Leu Cys Glu Arg Leu Asp Leu Gly Glu Gly Asp Leu Asn Lys Val Ala
 65 70 75 80

Asp Ala Cys Leu Ile Tyr Leu Ser Val Lys Gln Pro Val Lys Leu Gln
 85 90 95

Glu Ala Ala Arg Ser Val Phe Leu His Leu Met Lys Val Asp Pro Asp
 100 105 110

Ser Thr Trp Phe Leu Leu Asn Glu Leu Tyr Cys Pro Val Gln Phe Thr
 115 120 125

Pro Pro His Pro Ser Leu His Pro Val Gln Leu Xaa Gly Ala Ser Gly
 130 135 140

Gln Gln Asn Pro Xaa His Asp Gln Arg Ala Pro Ala Ala Gln Gly Ala
 145 150 155 160

Ala Val Thr Leu Leu Pro His His Arg Gly His Arg Ser Leu Pro Tyr
 165 170 175

Cys Gln Pro Glu Ala Gly Leu Thr Pro Pro Arg Pro
 180 185

<210> 413

<211> 138

<212> PRT

<213> Homo sapiens

<400> 413

Gly Ala Asp Gly Asn Val Ser Asp Phe Asp Asn Glu Glu Glu Glu Gln
 1 5 10 15

Ser Val Pro Pro Lys Val Asp Glu Asn Asp Thr Arg Pro Asp Val Glu
 20 25 30

Pro Pro Leu Pro Leu Gln Ile Gln Ile Ala Met Asp Val Met Glu Arg
 35 40 45

Cys Ile His Leu Leu Ser Asp Lys Asn Leu Gln Ile Arg Leu Lys Val
 50 55 60

Leu Asp Val Leu Asp Leu Cys Val Val Val Leu Gln Ser His Lys Asn

```
<210> 417
<211> 26
<212> PRT
<213> Homo sapiens
```

194

<400> 417

Trp Pro Ser Leu Val His Arg Leu Thr Arg Asp Ala Pro Leu Ala Val
 1 5 10 15

Leu Arg Ala Phe Lys Phe Tyr Val Pro Trp
 20 25

<210> 418

<211> 58

<212> PRT

<213> Homo sapiens

<400> 418

Ser Leu Gly Ile Ser Thr Phe Gly Ile Met Val Phe Ser Val Tyr Phe
 1 5 10 15

Gly Gly Ile Met Ile Ser Ile Pro Tyr Ser Gly Ile Ser Phe Gly Asn
 20 25 30

Lys Lys Glu Leu Asn Ile Asp Ser Cys Tyr Asn Met Val Asn Leu Lys
 35 40 45

Asn Ile Met Phe Ser Glu Arg Ser Gln Thr
 50 55

<210> 419

<211> 15

<212> PRT

<213> Homo sapiens

<400> 419

His Ala Ser Gly Asn Asn Asp Pro Leu Trp Phe Leu Thr Tyr Leu
 1 5 10 15

<210> 420

<211> 21

<212> PRT

<213> Homo sapiens

<400> 420

Met Val Phe Ser Val Tyr Phe Gly Gly Ile Met Ile Ser Ile Pro Tyr
 1 5 10 15

Ser Gly Ile Ser Phe
 20

<210> 421

<211> 20

<212> PRT

<213> Homo sapiens

<400> 421

Phe Gly Asn Lys Lys Glu Leu Asn Ile Asp Ser Cys Tyr Asn Met Val

1

5

10⁵
10

15

Asn Leu Lys Asn
20

<210> 422
<211> 75
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (48)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (49)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (50)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (72)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (74)
<223> Xaa equals any of the naturally occurring L-amino acids

<400> 422
Met Asn Ser Phe Ser Val Ile Ala Ser Ile Val Val Leu Leu Pro Phe
1 5 10 15

Pro Gly Leu Ser Val Ser Ala Cys Leu Pro Ser His Ser His Gln Cys
20 25 30

Lys Thr Phe Ile Leu Leu Phe Leu Pro Ser Ser Glu Lys Thr Leu Xaa
35 40 45

Xaa Xaa Pro Pro Ser His Ser Ser Thr Leu Gly Gly Gln Gly Gly Gln
50 55 60

Ile Met Arg Ser Gly Asp Arg Xaa His Xaa Gly
65 70 75

<210> 423
<211> 81
<212> PRT
<213> Homo sapiens

196

<220>

<221> SITE

<222> (5)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (6)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (60)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (74)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 423

Val	Val	Phe	Phe	Xaa	Xaa	Phe	Phe	Glu	Met	Glu	Ser	His	Ser	Val	Ala
1				5					10					15	

Gln	Ala	Gly	Val	Gln	Trp	Arg	Asn	Leu	Gly	Ser	Leu	Gln	Ala	Leu	Pro
			20					25					30		

Pro	Gly	Phe	Met	Pro	Phe	Ser	Cys	Leu	Ser	Leu	Pro	Gly	Ser	Trp	Asp
		35					40						45		

Tyr	Arg	Arg	Pro	Pro	Pro	Ser	Pro	Ala	Asn	Leu	Xaa	Cys	Ile	Phe	Ser
	50					55					60				

Arg	Asp	Gly	Gly	His	His	Val	Ser	Gln	Xaa	Gly	Leu	Asp	Leu	Leu	Thr
65					70					75					80

Ser

<210> 424

<211> 28

<212> PRT

<213> Homo sapiens

<400> 424

Ile	Val	Val	Leu	Leu	Pro	Phe	Pro	Gly	Leu	Ser	Val	Ser	Ala	Cys	Leu
1				5					10					15	

Pro	Ser	His	Ser	His	Gln	Cys	Lys	Thr	Phe	Ile	Leu
		20					25				

<210> 425

<211> 26

<212> PRT

<213> Homo sapiens

<400> 425

197

Pro Gly Phe Met Pro Phe Ser Cys Leu Ser Leu Pro Gly Ser Trp Asp
 1 5 10 15

Tyr Arg Arg Pro Pro Pro Ser Pro Ala Asn
 20 25

<210> 426

<211> 16

<212> PRT

<213> Homo sapiens

<400> 426

Tyr Arg Phe Lys Asn Pro Lys Cys Arg Leu Phe Ser Val Pro Cys Arg
 1 5 10 15

<210> 427

<211> 128

<212> PRT

<213> Homo sapiens

<400> 427

Thr Gln Asn Arg Glu Leu Leu Ala Trp Lys Pro Lys Gly Thr Asp Asp
 1 5 10 15

Ile Cys Thr Ser His Asn Thr Thr His Ile Gln Lys Met Pro Gly Glu
 20 25 30

Ala Asn Ser Cys Cys Pro Arg Gly Ala Lys Ser Tyr His Ile Asp Cys
 35 40 45

Trp Pro Pro Ala Leu Phe Pro Arg Cys Val Ala Tyr Leu Phe Leu Asn
 50 55 60

Lys Pro Ala Thr Leu Arg Lys Lys Tyr Tyr Cys Lys Pro Tyr His Thr
 65 70 75 80

Gln Leu His Pro Ala Trp His Arg Glu Lys Ser Ala Phe Trp Ile Phe
 85 90 95

Glu Thr Val Ser Gln Ser Lys Gln Ser Leu Thr Ser Leu Val Tyr Ser
 100 105 110

Val Asn Glu Leu Leu Val Leu Ser Asn Leu Ala Gln Trp Ala Leu Gly
 115 120 125

<210> 428

<211> 23

<212> PRT

<213> Homo sapiens

198

<400> 428

Ala Trp Lys Pro Lys Gly Thr Asp Asp Ile Cys Thr Ser His Asn Thr
1 5 10 15

Thr His Ile Gln Lys Met Pro
20

<210> 429

<211> 25

<212> PRT

<213> Homo sapiens

<400> 429

Cys Pro Arg Gly Ala Lys Ser Tyr His Ile Asp Cys Trp Pro Pro Ala
1 5 10 15

Leu Phe Pro Arg Cys Val Ala Tyr Leu
20 25

<210> 430

<211> 26

<212> PRT

<213> Homo sapiens

<400> 430

Ser Tyr His Ile Asp Cys Trp Pro Pro Ala Leu Phe Pro Arg Cys Val
1 5 10 15

Ala Tyr Leu Phe Leu Asn Lys Pro Ala Thr
20 25

<210> 431

<211> 29

<212> PRT

<213> Homo sapiens

<400> 431

Arg Lys Lys Tyr Tyr Cys Lys Pro Tyr His Thr Gln Leu His Pro Ala
1 5 10 15

Trp His Arg Glu Lys Ser Ala Phe Trp Ile Phe Glu Thr
20 25

<210> 432

<211> 28

<212> PRT

<213> Homo sapiens

<400> 432

Ile Cys Leu Asp Ser Cys Ser Gln Val Ser Val Thr Ser Leu Trp Ser
1 5 10 15

199

Phe Leu Arg Val His Ser Leu Val Gln Thr Leu Trp
 20 25

<210> 433
 <211> 75
 <212> PRT
 <213> Homo sapiens

<400> 433
 His Tyr Cys Cys Asp Phe Gly Thr Ser Leu Leu Gly Phe Tyr Val Pro
 1 5 10 15

Phe His Tyr Tyr Val His Met Val Asn Ile Ile Leu Thr Thr Ile Asp
 20 25 30

Phe Tyr His Tyr Lys Phe Cys Cys Ser Gln Asn Ala Asn Lys His Cys
 35 40 45

Phe Lys His Phe Gln Ile Met Thr Thr Val Pro Tyr Leu Asn Ile Asn
 50 55 60

Lys Glu Asn Leu Arg Phe Lys Asn Ile Phe Lys
 65 70 75

<210> 434
 <211> 27
 <212> PRT
 <213> Homo sapiens

<400> 434
 Thr Ser Leu Leu Gly Phe Tyr Val Pro Phe His Tyr Tyr Val His Met
 1 5 10 15

Val Asn Ile Ile Leu Thr Thr Ile Asp Phe Tyr
 20 25

<210> 435
 <211> 22
 <212> PRT
 <213> Homo sapiens

<400> 435
 Phe Gln Ile Met Thr Thr Val Pro Tyr Leu Asn Ile Asn Lys Glu Asn
 1 5 10 15

Leu Arg Phe Lys Asn Ile
 20

<210> 436
 <211> 106
 <212> PRT
 <213> Homo sapiens

<400> 436

200

Ile	Ser	Glu	Ser	Met	Ser	Leu	Val	Arg	Ser	Leu	Gln	Phe	Tyr	Arg	Gly
1				5					10					15	
Lys	Asn	Arg	Ala	Glu	Arg	Thr	Val	Ile	Ser	Ser	Ser	Ser	His	Ser	Cys
			20					25					30		
His	Leu	Ile	Asp	Leu	Glu	Phe	Gln	Pro	Arg	Ser	Asp	Gly	Glu	Val	Ser
		35					40					45			
Ile	Ser	Phe	Leu	Glu	Lys	Gly	Val	Glu	Leu	Arg	Trp	Gly	Met	Gly	Leu
	50					55					60				
Glu	Asp	Leu	Ile	Gly	Leu	Gly	Leu	Gly	Val	Ser	Thr	Arg	Arg	Ser	Thr
65					70					75					80
Val	Arg	Arg	Lys	Glu	Pro	Thr	Lys	Ala	Gly	Met	His	Thr	Ala	Cys	Ser
				85					90					95	
Glu	Glu	Met	Glu	Pro	Glu	Asn	Arg	Glu	Asn						
			100					105							

<210> 437

<211> 143

<212> PRT

<213> Homo sapiens

<400> 437

Asp	Gly	Ser	Arg	Ser	Val	Ala	Gln	Ala	Arg	Val	Gln	Trp	His	His	Arg
1				5					10					15	
Gly	Ser	Leu	Pro	Pro	Leu	Pro	Pro	Arg	Phe	Lys	Gln	Phe	Pro	Leu	Arg
			20					25					30		
His	Leu	Arg	Val	Gly	Gly	Ile	Thr	Gly	Ala	Cys	Arg	His	Thr	Gln	Ile
		35					40					45			
Ile	Phe	Val	Val	Leu	Val	Gln	Met	Gly	Phe	His	His	Val	Gly	Gln	Ala
	50					55				60					
Gly	Leu	Glu	Leu	Leu	Thr	Ser	Gly	Asp	Pro	Pro	Ala	Leu	Ala	Ser	Gln
65					70					75					80
Ser	Ala	Gly	Ile	Thr	Gly	Val	Ser	His	Ser	Thr	Arg	Pro	Lys	Leu	Leu
				85					90					95	
Ser	Trp	Leu	Pro	Ser	Asp	Asn	Leu	Leu	Gly	Met	Ala	Leu	Tyr	Ser	Ile
			100					105					110		
Gln	Trp	Ala	Leu	Leu	Ala	Asn	Ser	Leu	Tyr	Phe	Gln	Val	Pro	Ser	Pro
		115					120					125			
Leu	Ser	Met	Leu	Cys	Ala	Phe	Leu	Pro	Leu	Trp	Val	Pro	Ser	Ala	
	130					135					140				

<210> 438

<211> 27

<212> PRT

29

<213> Homo sapiens

<400> 438

Arg Gly Lys Asn Arg Ala Glu Arg Thr Val Ile Ser Ser Ser Ser His
1 5 10 15

Ser Cys His Leu Ile Asp Leu Glu Phe Gln Pro
20 25

<210> 439

<211> 32

<212> PRT

<213> Homo sapiens

<400> 439

Leu Gly Leu Gly Val Ser Thr Arg Arg Ser Thr Val Arg Arg Lys Glu
1 5 10 15

Pro Thr Lys Ala Gly Met His Thr Ala Cys Ser Glu Glu Met Glu Pro
20 25 30

<210> 440

<211> 24

<212> PRT

<213> Homo sapiens

<400> 440

Gly Asp Pro Pro Ala Leu Ala Ser Gln Ser Ala Gly Ile Thr Gly Val
1 5 10 15

Ser His Ser Thr Arg Pro Lys Leu
20

<210> 441

<211> 25

<212> PRT

<213> Homo sapiens

<400> 441

Ala Leu Tyr Ser Ile Gln Trp Ala Leu Leu Ala Asn Ser Leu Tyr Phe
1 5 10 15

Gln Val Pro Ser Pro Leu Ser Met Leu
20 25

<210> 442

<211> 35

<212> PRT

<213> Homo sapiens

<400> 442

Asp Arg Ile Leu Leu Phe Tyr Ser Arg Asp Gly Gln Thr Thr Ser Lys
 1 5 10 15

Gly Pro Asn Pro Ala Cys Cys Leu Phe Leu Leu Lys Lys Phe Tyr Trp
 20 25 30

Asn Thr Ala
 35

<210> 443

<211> 21

<212> PRT

<213> Homo sapiens

<400> 443

Asp Gly Gln Thr Thr Ser Lys Gly Pro Asn Pro Ala Cys Cys Leu Phe
 1 5 10 15

Leu Leu Lys Lys Phe
 20

<210> 444

<211> 24

<212> PRT

<213> Homo sapiens

<400> 444

Asp Pro Arg Val Arg Arg Thr Leu Asp Leu Gly Ile Thr Leu Tyr Leu
 1 5 10 15

Phe Leu Tyr Ile Phe Leu Ser Leu
 20

<210> 445

<211> 244

<212> PRT

<213> Homo sapiens

<400> 445

Pro Ala Leu Gly Glu Cys Cys Leu Asp Ala Phe Leu Phe Leu Leu Gly
 1 5 10 15

Lys Gln Leu Lys Lys Ser Gly Glu Lys Pro Leu Leu Gly Gly Ser Leu
 20 25 30

Met Glu Tyr Ala Ile Leu Ser Ala Ile Ala Ala Met Asn Glu Pro Lys
 35 40 45

Thr Cys Ser Thr Thr Ala Leu Lys Lys Tyr Val Leu Glu Asn His Pro
 50 55 60

Gly Thr Asn Ser Asn Tyr Gln Met His Leu Leu Lys Lys Thr Leu Gln
 65 70 75 80

203

Lys Cys Glu Lys Asn Gly Trp Met Glu Gln Ile Ser Gly Lys Gly Phe
 85 90 95
 Ser Gly Thr Phe Gln Leu Cys Phe Pro Tyr Tyr Pro Ser Pro Gly Val
 100 105 110
 Leu Phe Pro Lys Lys Glu Pro Asp Asp Ser Arg Asp Glu Asp Glu Asp
 115 120 125
 Glu Asp Glu Ser Ser Glu Glu Asp Ser Glu Asp Glu Glu Pro Pro Pro
 130 135 140
 Lys Arg Arg Leu Gln Lys Lys Thr Pro Ala Lys Ser Pro Gly Lys Ala
 145 150 155 160
 Ala Ser Val Lys Gln Arg Gly Ser Lys Pro Ala Pro Lys Val Ser Ala
 165 170 175
 Ala Gln Arg Gly Lys Ala Arg Pro Leu Pro Lys Lys Ala Pro Pro Lys
 180 185 190
 Ala Lys Thr Pro Ala Lys Lys Thr Arg Pro Ser Ser Thr Val Ile Lys
 195 200 205
 Lys Pro Ser Gly Gly Ser Ser Lys Lys Pro Ala Thr Ser Ala Arg Lys
 210 215 220
 Glu Val Lys Leu Pro Gly Lys Gly Lys Ser Thr Met Lys Lys Ser Phe
 225 230 235 240
 Arg Val Lys Lys

<210> 446

<211> 152

<212> PRT

<213> Homo sapiens

<400> 446

Asp Phe Glu Phe His His Asp Thr Leu Phe Ser Tyr Lys Ile Tyr Phe
 1 5 10 15
 Phe Thr Leu Lys Asp Phe Phe Met Val Asp Leu Pro Leu Pro Gly Asn
 20 25 30
 Phe Thr Ser Phe Leu Ala Leu Val Ala Gly Phe Phe Glu Glu Pro Pro
 35 40 45
 Leu Gly Phe Leu Met Thr Val Asp Glu Gly Leu Val Phe Leu Ala Gly
 50 55 60
 Val Leu Ala Leu Gly Gly Ala Phe Leu Gly Lys Gly Leu Ala Phe Pro
 65 70 75 80
 Arg Trp Ala Ala Glu Thr Leu Gly Ala Gly Leu Asp Pro Leu Cys Phe
 85 90 95

204

Thr Asp Ala Ala Phe Pro Gly Asp Leu Ala Gly Val Phe Phe Cys Asn
 100 105 110

Leu Leu Leu Gly Gly Gly Ser Ser Ser Ser Glu Ser Ser Ser Asp Asp
 115 120 125

Ser Ser Ser Ser Ser Ser Ser Ser Leu Glu Ser Ser Gly Ser Phe Phe
 130 135 140

Gly Asn Arg Thr Pro Gly Leu Gly
 145 150

<210> 447
 <211> 28
 <212> PRT
 <213> Homo sapiens

<400> 447
 Cys Leu Asp Ala Phe Leu Phe Leu Leu Gly Lys Gln Leu Lys Lys Ser
 1 5 10 15

Gly Glu Lys Pro Leu Leu Gly Gly Ser Leu Met Glu
 20 25

<210> 448
 <211> 30
 <212> PRT
 <213> Homo sapiens

<400> 448
 Tyr Gln Met His Leu Leu Lys Lys Thr Leu Gln Lys Cys Glu Lys Asn
 1 5 10 15

Gly Trp Met Glu Gln Ile Ser Gly Lys Gly Phe Ser Gly Thr
 20 25 30

<210> 449
 <211> 28
 <212> PRT
 <213> Homo sapiens

<400> 449
 Lys Thr Pro Ala Lys Ser Pro Gly Lys Ala Ala Ser Val Lys Gln Arg
 1 5 10 15

Gly Ser Lys Pro Ala Pro Lys Val Ser Ala Ala Gln
 20 25

<210> 450
 <211> 28
 <212> PRT
 <213> Homo sapiens

<400> 450

205

Ser Ser Lys Lys Pro Ala Thr Ser Ala Arg Lys Glu Val Lys Leu Pro
1 5 10 15

Gly Lys Gly Lys Ser Thr Met Lys Lys Ser Phe Arg
20 25

<210> 451

<211> 23

<212> PRT

<213> Homo sapiens

<400> 451

Val Asp Glu Gly Leu Val Phe Leu Ala Gly Val Leu Ala Leu Gly Gly
1 5 10 15

Ala Phe Leu Gly Lys Gly Leu
20

<210> 452

<211> 25

<212> PRT

<213> Homo sapiens

<400> 452

Gly Leu Asp Pro Leu Cys Phe Thr Asp Ala Ala Phe Pro Gly Asp Leu
1 5 10 15

Ala Gly Val Phe Phe Cys Asn Leu Leu
20 25

<210> 453

<211> 59

<212> PRT

<213> Homo sapiens

<400> 453

Thr Met Leu Phe Tyr Leu Ser Ser Gln Pro Asp Trp Gln Leu Asp Phe
1 5 10 15

Phe Arg Val Ser Phe Asn Gly Pro Val Phe Phe Ile Ile Ile Phe Asn
20 25 30

Asp Arg Ala Gly Phe Arg Met Gln Ala Leu Val Ser Gln Ala Ala Cys
35 40 45

Arg Arg Ser Arg Tyr Lys Leu Ser Val Val Tyr
50 55

<210> 454

<211> 23

<212> PRT

<213> Homo sapiens

<400> 454

206

Asp	Arg	Ala	Gly	Phe	Arg	Met	Gln	Ala	Leu	Val	Ser	Gln	Ala	Ala	Cys
1				5					10					15	

Arg	Arg	Ser	Arg	Tyr	Lys	Leu
			20			

<210> 455
 <211> 22
 <212> PRT
 <213> Homo sapiens

Leu	Ala	Ala	Gly	Ile	Leu	Asn	Ser	Ser	Leu	Pro	Ala	Leu	Tyr	His	Ser
1				5					10					15	

Val	Glu	Glu	Ile	Ser	Gln
			20		

<210> 456
 <211> 45
 <212> PRT
 <213> Homo sapiens

Xaa	Tyr	Arg	Met	Asn	Thr	Lys	Phe	Leu	Glu	Ser	Tyr	Lys	Met	Ser	Thr
1				5					10					15	

Thr	Leu	Ser	Arg	Arg	His	Gln	Asn	Val	Ser	Leu	Cys	Lys	Asp	Met	Lys
			20					25					30		

Thr	Pro	Ala	Gly	Thr	Asp	Thr	Lys	Ile	Ala	Phe	Leu	Glu
			35				40					45

<210> 457
 <211> 21
 <212> PRT
 <213> Homo sapiens

Ser	Tyr	Lys	Met	Ser	Thr	Thr	Leu	Ser	Arg	Arg	His	Gln	Asn	Val	Ser
1				5					10					15	

Leu	Cys	Lys	Asp	Met
			20	

<210> 458
 <211> 57
 <212> PRT
 <213> Homo sapiens

Ile	Cys	Ile	Glu	Ser	Leu	Met	Leu	His	Tyr	Ile	Ala	Leu	Val	Phe	Glu
1				5					10					15	

107

Met Ala Phe Met Phe Pro Leu Val Tyr His Glu Met Gly Ser Asp Ser
 20 25 30

Ile Arg Phe His Leu Cys Gln Val Asp Ser Cys Leu Pro Ser Met Met
 35 40 45

Arg Phe Phe Phe Ser Phe Pro Phe Leu
 50 55

<210> 459
 <211> 21
 <212> PRT
 <213> Homo sapiens

<400> 459
 Tyr Ile Ala Leu Val Phe Glu Met Ala Phe Met Phe Pro Leu Val Tyr
 1 5 10 15

His Glu Met Gly Ser
 20

<210> 460
 <211> 21
 <212> PRT
 <213> Homo sapiens

<400> 460
 Ser Asp Ser Ile Arg Phe His Leu Cys Gln Val Asp Ser Cys Leu Pro
 1 5 10 15

Ser Met Met Arg Phe
 20

<210> 461
 <211> 115
 <212> PRT
 <213> Homo sapiens

<400> 461
 Gly Gly Val Ser Val Gln Asp Gly Ser Leu Arg Glu Glu Thr Asp Val
 1 5 10 15

Gly Glu Gly Gly Arg Pro Arg Gly Gly Gln Ser Glu Gly Ala Arg Val
 20 25 30

Thr Arg Arg Pro Ser Pro Pro Asp Ser Asn Ala Ser Ala Phe Asp Leu
 35 40 45

Asp Leu Asp Phe Ser Pro Phe Cys Ile Trp Cys Tyr Arg Leu Glu Thr
 50 55 60

Pro Ala Glu Val Val Phe Ser Pro Ala Pro Leu Arg Leu Ser Gly Pro
 65 70 75 80

Gly Leu Ala Pro Val Val Phe Val Ser Thr Leu Pro Ser Leu Gln Pro

85

208
90

95

Ser Ser Phe Cys Gly Trp Asp Leu Pro Ala Arg Pro Arg Gly Leu Ser
 100 105 110

Gly Phe Arg
 115

<210> 462

<211> 111

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (82)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 462

Phe Thr Asn Lys Ser Cys Ser Lys Met Ser Ser Thr His Leu Tyr Lys
 1 5 10 15

Gly Ser Asp Val Leu Cys Tyr Ala Arg Ser Ser Glu Ser Met Ser Leu
 20 25 30

Ser Cys Gly Asp Val Ala Asn Ala Gly Arg Leu Thr Pro Arg Leu His
 35 40 45

Leu Ala Arg Ser Ala Ser Gln Gly Pro Pro Thr Leu Pro Arg Val Pro
 50 55 60

Pro Arg Gly Ser Arg Pro Pro Thr Ala Gly Glu Ser Pro Ala Pro Arg
 65 70 75 80

Thr Xaa Ser Leu Glu Asn His Lys Asn Ile Asp His Leu Ser Ser Asn
 85 90 95

Ser His Gly Lys Phe Arg Ile Tyr Gly Gln Asn Asp Ile Lys Ile
 100 105 110

<210> 463

<211> 80

<212> PRT

<213> Homo sapiens

<400> 463

Gln Asp Val Ile Tyr Thr Phe Val Gln Arg Phe Arg Arg Pro Met Leu
 1 5 10 15

Cys Thr Ile Leu Arg Lys Tyr Glu Pro Val Val Arg Gly Arg Arg Lys
 20 25 30

Arg Trp Gln Ala His Pro Ser Ser Ala Phe Gly Lys Lys Arg Leu Pro
 35 40 45

Arg Pro Pro His Pro Ala Gln Gly Ala Pro Gln Arg Glu Gln Ala Ser

50

55

309

60

His Ser Trp Arg Glu Pro Gly Pro Gln Asn Thr Phe Pro Arg Lys Pro
 65 70 75 80

<210> 464

<211> 22

<212> PRT

<213> Homo sapiens

<400> 464

Arg Glu Glu Thr Asp Val Gly Glu Gly Gly Arg Pro Arg Gly Gly Gln
 1 5 10 15

Ser Glu Gly Ala Arg Val
 20

<210> 465

<211> 27

<212> PRT

<213> Homo sapiens

<400> 465

Gly Pro Gly Leu Ala Pro Val Val Phe Val Ser Thr Leu Pro Ser Leu
 1 5 10 15

Gln Pro Ser Ser Phe Cys Gly Trp Asp Leu Pro
 20 25

<210> 466

<211> 24

<212> PRT

<213> Homo sapiens

<400> 466

Met Ser Ser Thr His Leu Tyr Lys Gly Ser Asp Val Leu Cys Tyr Ala
 1 5 10 15

Arg Ser Ser Glu Ser Met Ser Leu
 20

<210> 467

<211> 28

<212> PRT

<213> Homo sapiens

<400> 467

Ser Gln Gly Pro Pro Thr Leu Pro Arg Val Pro Pro Arg Gly Ser Arg
 1 5 10 15

Pro Pro Thr Ala Gly Glu Ser Pro Ala Pro Arg Thr

20

25 710

<210> 468
 <211> 25
 <212> PRT
 <213> Homo sapiens

<400> 468
 Arg Phe Arg Arg Pro Met Leu Cys Thr Ile Leu Arg Lys Tyr Glu Pro
 1 5 10 15
 Val Val Arg Gly Arg Arg Lys Arg Trp
 20 25

<210> 469
 <211> 24
 <212> PRT
 <213> Homo sapiens

<400> 469
 Arg Leu Pro Arg Pro Pro His Pro Ala Gln Gly Ala Pro Gln Arg Glu
 1 5 10 15
 Gln Ala Ser His Ser Trp Arg Glu
 20

<210> 470
 <211> 81
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (43)
 <223> Xaa equals any of the naturally occurring L-amino acids

<400> 470
 Arg Gly Met Arg Gly Arg Trp Leu Val Ser Ser Gly Ala Ala Phe Pro
 1 5 10 15
 Ile Pro Leu Asn Gly Phe Cys Glu Ser Arg Glu Phe Phe Pro Asp Ser
 20 25 30
 Gly Ser Val Leu Leu His Trp Arg Pro Asn Xaa Val Leu Ile Glu Ile
 35 40 45
 Lys Val Phe Gly Ser Arg Ser Gln Ser Leu Ile Ser Ser Lys Asn Leu
 50 55 60
 Lys Thr Ser Leu Thr Phe Ile Tyr Gly Lys Val Glu Glu Val Leu Asn
 65 70 75 80

Asn

211

<210> 471
 <211> 81
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (62)
 <223> Xaa equals any of the naturally occurring L-amino acids

<400> 471
 Leu Lys Leu Ser Ser Ala Asp Ser Gln Ala Ile Met Asn Ile Phe Ser
 1 5 10 15
 Ala Asp Cys Met Pro Arg Leu His Ile Ala Leu Gln Thr Glu Met Ile
 20 25 30
 Pro Asn Arg Ala Pro Gln Gly Gly Ala Ala Ala Asn Leu Trp His Glu
 35 40 45
 Ala Gln Tyr Arg Arg Leu Pro Phe Ser Arg Ala Pro Glu Xaa Thr Asp
 50 55 60
 Ala His Gln Ala Ser Ala Gln Arg Gly Ala Ala Gln Leu Pro Arg Glu
 65 70 75 80
 Gln

<210> 472
 <211> 28
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (28)
 <223> Xaa equals any of the naturally occurring L-amino acids

<400> 472
 Pro Ile Pro Leu Asn Gly Phe Cys Glu Ser Arg Glu Phe Phe Pro Asp
 1 5 10 15
 Ser Gly Ser Val Leu Leu His Trp Arg Pro Asn Xaa
 20 25

<210> 473
 <211> 29
 <212> PRT
 <213> Homo sapiens

<400> 473
 Asn Ile Phe Ser Ala Asp Cys Met Pro Arg Leu His Ile Ala Leu Gln
 1 5 10 15

212

Thr Glu Met Ile Pro Asn Arg Ala Pro Gln Gly Gly Ala
 20 25

<210> 474

<211> 37

<212> PRT

<213> Homo sapiens

<400> 474

Thr Phe Arg Leu Val Ser Ala His Leu Lys Thr Arg Lys Leu Ile Asn
 1 5 10 15

Pro Glu Ala Ala Glu Arg Arg Trp Arg Asp Trp Asp Ser Arg Gln Gly
 20 25 30

Trp Leu Ser Val Lys
 35

<210> 475

<211> 21

<212> PRT

<213> Homo sapiens

<400> 475

Lys Thr Arg Lys Leu Ile Asn Pro Glu Ala Ala Glu Arg Arg Trp Arg
 1 5 10 15

Asp Trp Asp Ser Arg
 20

<210> 476

<211> 83

<212> PRT

<213> Homo sapiens

<400> 476

Trp Asn Tyr Thr Val Asn Asn Leu Tyr Leu Phe Ser Phe Ser Ile Val
 1 5 10 15

Ser Met Lys Phe Met His Val Leu Ser Ile Asn Ile Phe Phe Gly Arg
 20 25 30

Ala Arg Trp Leu Thr Pro Val Ile Pro Ala Leu Leu Glu Ala Glu Ala
 35 40 45

Gly Gly Ser Leu Gly Gln Glu Phe Lys Thr Ser Leu Gly Lys Asp Gly
 50 55 60

Glu Thr Pro Ser Leu Leu Lys Ile Gln Lys Leu Ala Gly His Gly Gly
 65 70 75 80

Arg Arg Leu

<210> 477
 <211> 76
 <212> PRT
 <213> Homo sapiens

713

<400> 477
 Asp Gln Pro Gly Lys His Gly Glu Thr Leu Ser Leu Leu Lys Met Gln
 1 5 10 15
 Lys Leu Thr Trp Cys Gly Gly Met Pro Phe Val Ile Pro Ser Tyr Ser
 20 25 30
 Arg Ser Pro Arg Pro Glu Asn Arg Leu Asn Leu Gly Asp Arg Gly Cys
 35 40 45
 Thr Glu Leu Leu His Ser Ser Leu Gly Asn Arg Val Arg Leu Ser Lys
 50 55 60
 Lys Lys Glu Val Tyr Met Met Glu Leu Tyr Ser Lys
 65 70 75

<210> 478
 <211> 28
 <212> PRT
 <213> Homo sapiens

<400> 478
 Val Ile Pro Ala Leu Leu Glu Ala Glu Ala Gly Gly Ser Leu Gly Gln
 1 5 10 15
 Glu Phe Lys Thr Ser Leu Gly Lys Asp Gly Glu Thr
 20 25

<210> 479
 <211> 29
 <212> PRT
 <213> Homo sapiens

<400> 479
 Asn Arg Leu Asn Leu Gly Asp Arg Gly Cys Thr Glu Leu Leu His Ser
 1 5 10 15
 Ser Leu Gly Asn Arg Val Arg Leu Ser Lys Lys Lys Glu
 20 25

<210> 480
 <211> 17
 <212> PRT
 <213> Homo sapiens

<400> 480
 His Ala Ser Glu His Leu Ala Ala Leu Pro Val Asn Val Lys Ile Gly
 1 5 10 15

Lys

214

<210> 481
 <211> 77
 <212> PRT
 <213> Homo sapiens

<400> 481
 Leu Val Cys Ile Leu Leu Val His Trp Ile Pro Pro Leu Gly Ala Trp
 1 5 10 15
 Gly Leu Ser Leu Met Leu Phe Leu Ile Leu Glu Gln Arg Cys Gly Lys
 20 25 30
 Gly Lys Trp Arg Asn Ala Leu Leu Ser Val Ser Phe Ser Val Pro Gln
 35 40 45
 Leu Gln Met Gln Lys Val Ser Leu Asp Ser Thr Pro Leu Asn Val Asn
 50 55 60
 His Asp Lys Met Asp Ile Trp Lys Leu Thr Pro Lys Leu
 65 70 75

<210> 482
 <211> 57
 <212> PRT
 <213> Homo sapiens

<400> 482
 Ile Met Ile Lys Trp Ile Phe Gly Asn Leu Leu Leu Ser Cys Asp Leu
 1 5 10 15
 Gly Cys Ile Ser Thr Ser Gly Leu Pro Gln Tyr Gln Gly Leu Arg Leu
 20 25 30
 Leu Asn Phe Glu Tyr Ser Leu Gly Phe Met Leu Arg Ser Leu Trp Ser
 35 40 45
 Arg Ser Ala Ile Gln Cys Phe Phe Ser
 50 55

<210> 483
 <211> 21
 <212> PRT
 <213> Homo sapiens

<400> 483
 Leu Leu Leu Ser Cys Asp Leu Gly Cys Ile Ser Thr Ser Gly Leu Pro
 1 5 10 15
 Gln Tyr Gln Gly Leu
 20

<210> 484

<211> 21
 <212> PRT
 <213> Homo sapiens

MS

<400> 484
 Leu Arg Leu Leu Asn Phe Glu Tyr Ser Leu Gly Phe Met Leu Arg Ser
 1 5 10 15

Leu Trp Ser Arg Ser
 20

<210> 485
 <211> 78
 <212> PRT
 <213> Homo sapiens

<400> 485
 Ala Ser Pro His Leu Phe Ile Glu Lys Trp Gly Arg Ala Phe Ile Leu
 1 5 10 15

Arg Lys Leu Leu Leu Val Pro Val Ile Ser Lys Arg Ile Ile Asn Ile
 20 25 30

Met Ala His Gln Val Lys Pro Pro Ile Phe Cys Ala Met Ile Met Cys
 35 40 45

Asn Leu Phe Cys Ser Gly Tyr Glu His Leu Leu Phe Thr Leu Met Arg
 50 55 60

Phe Phe Ser Phe Glu Gln Ile Phe Asp Glu Val Val Phe His
 65 70 75

<210> 486
 <211> 25
 <212> PRT
 <213> Homo sapiens

<400> 486
 Lys Leu Leu Leu Val Pro Val Ile Ser Lys Arg Ile Ile Asn Ile Met
 1 5 10 15

Ala His Gln Val Lys Pro Pro Ile Phe
 20 25

<210> 487
 <211> 358
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (352)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

246

<222> (356)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 487

Phe Ala Val Ile Arg Phe Glu Ser Ile Ile His Glu Phe Asp Pro Trp
 1 5 10 15

Phe Asn Tyr Arg Ser Thr His His Leu Ala Ser His Gly Phe Tyr Glu
 20 25 30

Phe Leu Asn Trp Phe Asp Glu Arg Ala Trp Tyr Pro Leu Gly Arg Ile
 35 40 45

Val Gly Gly Thr Val Tyr Pro Gly Leu Met Ile Thr Ala Gly Leu Ile
 50 55 60

His Trp Ile Leu Asn Thr Leu Asn Ile Thr Val His Ile Arg Asp Val
 65 70 75 80

Cys Val Phe Leu Ala Pro Thr Phe Ser Gly Leu Thr Ser Ile Ser Thr
 85 90 95

Phe Leu Leu Thr Arg Glu Leu Trp Asn Gln Gly Ala Gly Leu Leu Ala
 100 105 110

Ala Cys Phe Ile Ala Ile Val Pro Gly Tyr Ile Ser Arg Ser Val Ala
 115 120 125

Gly Ser Phe Asp Asn Glu Gly Ile Ala Ile Phe Ala Leu Gln Phe Thr
 130 135 140

Tyr Tyr Leu Trp Val Lys Ser Val Lys Thr Gly Ser Val Phe Trp Thr
 145 150 155 160

Met Cys Cys Cys Leu Ser Tyr Phe Tyr Met Val Ser Ala Trp Gly Gly
 165 170 175

Tyr Val Phe Ile Ile Asn Leu Ile Pro Leu His Val Phe Val Leu Leu
 180 185 190

Leu Met Gln Arg Tyr Ser Lys Arg Val Tyr Ile Ala Tyr Ser Thr Phe
 195 200 205

Tyr Ile Val Gly Leu Ile Leu Ser Met Gln Ile Pro Phe Val Gly Phe
 210 215 220

Gln Pro Ile Arg Thr Ser Glu His Met Ala Ala Ala Gly Val Phe Ala
 225 230 235 240

Leu Leu Gln Ala Tyr Ala Phe Leu Gln Tyr Leu Arg Asp Arg Leu Thr
 245 250 255

Lys Gln Glu Phe Gln Thr Leu Phe Phe Leu Gly Val Ser Leu Ala Ala
 260 265 270

Gly Ala Val Phe Leu Ser Val Ile Tyr Leu Thr Tyr Thr Gly Tyr Ile
 275 280 285

217

Ala Pro Trp Ser Gly Arg Phe Tyr Ser Leu Trp Asp Thr Gly Tyr Ala
 290 295 300

Lys Ile His Ile Pro Ile Ile Ala Ser Val Ser Glu His Gln Pro Thr
 305 310 315 320

Thr Trp Val Ser Phe Phe Phe Asp Leu His Ile Leu Val Cys Thr Phe
 325 330 335

Pro Ala Gly Leu Trp Phe Cys Ile Lys Asn Ile Asn Asp Glu Arg Xaa
 340 345 350

Phe Gly Lys Xaa Gly Phe
 355

<210> 488

<211> 27

<212> PRT

<213> Homo sapiens

<400> 488

Glu Phe Asp Pro Trp Phe Asn Tyr Arg Ser Thr His His Leu Ala Ser
 1 5 10 15

His Gly Phe Tyr Glu Phe Leu Asn Trp Phe Asp
 20 25

<210> 489

<211> 23

<212> PRT

<213> Homo sapiens

<400> 489

Thr Arg Glu Leu Trp Asn Gln Gly Ala Gly Leu Leu Ala Ala Cys Phe
 1 5 10 15

Ile Ala Ile Val Pro Gly Tyr
 20

<210> 490

<211> 22

<212> PRT

<213> Homo sapiens

<400> 490

Thr Tyr Tyr Leu Trp Val Lys Ser Val Lys Thr Gly Ser Val Phe Trp
 1 5 10 15

Thr Met Cys Cys Cys Leu
 20

<210> 491

<211> 25

<212> PRT
 <213> Homo sapiens

218

<400> 491
 Gly Val Phe Ala Leu Leu Gln Ala Tyr Ala Phe Leu Gln Tyr Leu Arg
 1 5 10 15

Asp Arg Leu Thr Lys Gln Glu Phe Gln
 20 25

<210> 492
 <211> 27
 <212> PRT
 <213> Homo sapiens

<400> 492
 Tyr Ser Leu Trp Asp Thr Gly Tyr Ala Lys Ile His Ile Pro Ile Ile
 1 5 10 15

Ala Ser Val Ser Glu His Gln Pro Thr Thr Trp
 20 25

<210> 493
 <211> 408
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (20)
 <223> Xaa equals any of the naturally occurring L-amino acids

<400> 493
 Met Gly His Met Leu Tyr Leu Leu Gly Asn Ile Asn Lys Arg Thr Met
 1 5 10 15

His Lys Tyr Xaa Gln Glu Ser Lys Lys Ala Gly Lys Ala Ser Phe Ala
 20 25 30

Tyr Ala Trp Val Leu Asp Glu Thr Gly Glu Glu Arg Glu Arg Gly Val
 35 40 45

Thr Met Asp Val Gly Met Thr Lys Phe Glu Thr Thr Thr Lys Val Ile
 50 55 60

Thr Leu Met Asp Ala Pro Gly His Lys Asp Phe Ile Pro Asn Met Ile
 65 70 75 80

Thr Gly Ala Ala Gln Ala Asp Val Ala Val Leu Val Val Asp Ala Ser
 85 90 95

Arg Gly Glu Phe Glu Ala Gly Phe Glu Thr Gly Gly Gln Thr Arg Glu
 100 105 110

His Gly Leu Leu Val Arg Ser Leu Gly Val Thr Gln Leu Ala Val Ala
 115 120 125

219

Val Asn Lys Met Asp Gln Val Asn Trp Gln Gln Glu Arg Phe Gln Glu
 130 135 140
 Ile Thr Gly Lys Leu Gly His Phe Leu Lys Gln Ala Gly Phe Lys Glu
 145 150 155 160
 Ser Asp Val Gly Phe Ile Pro Thr Ser Gly Leu Ser Gly Glu Asn Leu
 165 170 175
 Ile Thr Arg Ser Gln Ser Ser Glu Leu Thr Lys Trp Tyr Lys Gly Leu
 180 185 190
 Cys Leu Leu Glu Gln Ile Asp Ser Phe Lys Pro Pro Gln Arg Ser Ile
 195 200 205
 Asp Lys Pro Phe Arg Leu Cys Val Ser Asp Val Phe Lys Asp Gln Gly
 210 215 220
 Ser Gly Phe Cys Ile Thr Gly Lys Ile Glu Ala Gly Tyr Ile Gln Thr
 225 230 235 240
 Gly Asp Arg Leu Leu Ala Met Pro Pro Asn Glu Thr Cys Thr Val Lys
 245 250 255
 Gly Ile Thr Leu His Asp Glu Pro Val Asp Trp Ala Ala Ala Gly Asp
 260 265 270
 His Val Ser Leu Thr Leu Val Gly Met Asp Ile Ile Lys Ile Asn Val
 275 280 285
 Gly Cys Ile Phe Cys Gly Pro Lys Val Pro Ile Lys Ala Cys Thr Arg
 290 295 300
 Phe Arg Ala Arg Ile Leu Ile Phe Asn Ile Glu Ile Pro Ile Thr Lys
 305 310 315 320
 Gly Phe Pro Val Leu Leu His Tyr Gln Thr Val Ser Glu Pro Ala Val
 325 330 335
 Ile Lys Arg Leu Ile Ser Val Leu Asn Lys Ser Thr Gly Glu Val Thr
 340 345 350
 Lys Lys Lys Pro Lys Phe Leu Thr Lys Gly Gln Asn Ala Leu Val Glu
 355 360 365
 Leu Gln Thr Gln Arg Pro Ile Ala Leu Glu Leu Tyr Lys Asp Phe Lys
 370 375 380
 Glu Leu Gly Arg Phe Met Leu Arg Tyr Gly Gly Ser Thr Ile Ala Ala
 385 390 395 400
 Gly Val Val Thr Glu Ile Lys Glu
 405

<210> 494

<211> 21

<212> PRT
 <213> Homo sapiens

270

<220>
 <221> SITE
 <222> (16)
 <223> Xaa equals any of the naturally occurring L-amino acids

<400> 494
 Leu Tyr Leu Leu Gly Asn Ile Asn Lys Arg Thr Met His Lys Tyr Xaa
 1 5 10 15

Gln Glu Ser Lys Lys
 20

<210> 495
 <211> 23
 <212> PRT
 <213> Homo sapiens

<400> 495
 Leu Asp Glu Thr Gly Glu Glu Arg Glu Arg Gly Val Thr Met Asp Val
 1 5 10 15

Gly Met Thr Lys Phe Glu Thr
 20

<210> 496
 <211> 22
 <212> PRT
 <213> Homo sapiens

<400> 496
 Gly His Lys Asp Phe Ile Pro Asn Met Ile Thr Gly Ala Ala Gln Ala
 1 5 10 15

Asp Val Ala Val Leu Val
 20

<210> 497
 <211> 23
 <212> PRT
 <213> Homo sapiens

<400> 497
 Gly Phe Glu Thr Gly Gly Gln Thr Arg Glu His Gly Leu Leu Val Arg
 1 5 10 15

Ser Leu Gly Val Thr Gln Leu
 20

<210> 498
 <211> 23
 <212> PRT

<213> Homo sapiens

221

<400> 498

Trp Gln Gln Glu Arg Phe Gln Glu Ile Thr Gly Lys Leu Gly His Phe
1 5 10 15

Leu Lys Gln Ala Gly Phe Lys
20

<210> 499

<211> 22

<212> PRT

<213> Homo sapiens

<400> 499

Thr Ser Gly Leu Ser Gly Glu Asn Leu Ile Thr Arg Ser Gln Ser Ser
1 5 10 15

Glu Leu Thr Lys Trp Tyr
20

<210> 500

<211> 23

<212> PRT

<213> Homo sapiens

<400> 500

Pro Gln Arg Ser Ile Asp Lys Pro Phe Arg Leu Cys Val Ser Asp Val
1 5 10 15

Phe Lys Asp Gln Gly Ser Gly
20

<210> 501

<211> 22

<212> PRT

<213> Homo sapiens

<400> 501

Leu Ile Ser Val Leu Asn Lys Ser Thr Gly Glu Val Thr Lys Lys Lys
1 5 10 15

Pro Lys Phe Leu Thr Lys
20

<210> 502

<211> 25

<212> PRT

<213> Homo sapiens

<400> 502

Gln Arg Pro Ile Ala Leu Glu Leu Tyr Lys Asp Phe Lys Glu Leu Gly
1 5 10 15

WO 99/24836

PCT/US98/23435


Arg Phe Met Leu Arg Tyr Gly Gly Ser
20 25

???

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>183</u> , line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input checked="" type="checkbox"/>	
Name of depositary institution American Type Culture Collection ("ATCC")	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit <div style="text-align: center;">16 OCTOBER 1997</div>	Accession Number <div style="text-align: center;">209368</div>
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

<div style="text-align: center; border-bottom: 1px solid black; margin-bottom: 5px;">For receiving Office use only</div> <div style="display: flex; align-items: center; margin-bottom: 10px;"> <input checked="" type="checkbox"/> This sheet was received with the international application </div> <div style="border-top: 1px solid black; padding-top: 10px;"> Authorized officer <div style="text-align: center; font-family: cursive;">  </div> </div>	<div style="text-align: center; border-bottom: 1px solid black; margin-bottom: 5px;">For International Bureau use only</div> <div style="display: flex; align-items: center; margin-bottom: 10px;"> <input type="checkbox"/> This sheet was received by the International Bureau on: </div> <div style="border-top: 1px solid black; padding-top: 10px;"> Authorized officer </div>
---	---

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>183</u> , line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input checked="" type="checkbox"/>	
Name of depositary institution <u>American Type Culture Collection ("ATCC")</u>	
Address of depositary institution (including postal code and country) <u>10801 University Boulevard</u> <u>Manassas, Virginia 20110-2209</u> <u>United States of America</u>	
Date of deposit <u>23 OCTOBER 1997</u>	Accession Number <u>209407</u>
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only
<input checked="" type="checkbox"/> This sheet was received with the international application
Authorized officer <u>Melvin S. Swoboda, Jr.</u>

For International Bureau use only
<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>183</u> , line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution <u>American Type Culture Collection ("ATCC")</u>	
Address of depositary institution <i>(including postal code and country)</i> <u>10801 University Boulevard</u> <u>Manassas, Virginia 20110-2209</u> <u>United States of America</u>	
Date of deposit <u>30 OCTOBER 1997</u>	Accession Number <u>209423</u>
C. ADDITIONAL INDICATIONS <i>(leave blank if not applicable)</i> This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE <i>(if the indications are not for all designated States)</i>	
E. SEPARATE FURNISHING OF INDICATIONS <i>(leave blank if not applicable)</i>	
The indications listed below will be submitted to the International Bureau later <i>(specify the general nature of the indications e.g., "Accession Number of Deposit")</i>	

For receiving Office use only	
<input checked="" type="checkbox"/> This sheet was received with the international application	
Authorized officer <u>Melvin S. Brooker, Jr.</u>	

For International Bureau use only	
<input type="checkbox"/> This sheet was received by the International Bureau on:	
Authorized officer	

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/23435

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : Please See Extra Sheet.

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 536/23.5, 23.1; 435/320.1, 440, 252.3, 69.1, 7.1; 530/350, 387.1; 514/12

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	SAPPERSTEIN, S.K. et al, p115 is a general vesicular transport factor related to the yeast endoplasmic reticulum to Golgi transport factor Usolp. Proceedings of the National Academy of Sciences USA. 17 January 1995, Vol. 92, No. 2, pages 522-526, see entire document.	1, 2, 5, and 7-10
X	BARROSO et al, Transcytosis-associated protein (TAP)/p115 is a general fusion factor required for binding of vesicles to acceptor membranes. Proceedings of the National Academy of Sciences USA. 17 January 1995, Vol. 92, No. 2, pages 527-531, see entire document.	11-13 and 16

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principles or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

29 JANUARY 1999

Date of mailing of the international search report

25 FEB 1999

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

JAMES MARTINELL

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/23435

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	SOHADA et al, Phosphorylation of the vesicle docking protein p115 regulates its association with the Golgi membrane. Journal of Biological Chemistry. 27 February 1998, Vol. 273, No. 9, pages 5385-5388, see entire document.	1, 2, 5-13, and 16
A	ADAMS et al, complementary DNA sequencing: Expressed sequence tags and the human genome project. Science. 21 June 1991, Vol. 252, pages 1651-1656, see entire document.	1-22

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/23435

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☒ Claims Nos.: 23
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

Claim 23 is directed to a product of the process of claim 20. Claim 20 is not a process for the production of a product, but a process for the detection of a substance. Hence, no meaningful search can be carried out.

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.

2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.

3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/23435

A. CLASSIFICATION OF SUBJECT MATTER:
IPC (6):

G01N 33/68, 33/53; C07K 16/00; C12N 15/11, 15/12, 15/00, 15/63; A61K 38/17, 38/16; C12P 21/02

A. CLASSIFICATION OF SUBJECT MATTER:
US CL :

536/23.5, 23.1; 435/320.1, 440, 252.3, 69.1, 7.1; 530/350, 387.1; 514/12

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

APS, STN, MPSRCH (SEQ ID NOs 11 and 136 only). One nucleotide sequence and one amino acid sequence have been searched. It is not clear which sequences are embraced by the claims because the claims refer to sequences X and Y. The table beginning at page 183 contains many sequences X and Y, yet the claims refer to X and Y in the singular. If the claims are to embrace more than one X and more than one Y, it is not clear whether each X always requires the corresponding sequence Y. Additionally, the claims are in improper format in referring to the description (see PCT Rule 6.2(a)). Accordingly, the first X nucleotide sequence disclosed and the first Y amino acid sequence corresponding to the first X sequence disclosed in the table were searched.